



# Mark Scheme (Results)

January 2020

Pearson Edexcel International GCSE in  
Chemistry (4CH1)  
Paper 1C

## **Edexcel and BTEC Qualifications**

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at [www.edexcel.com](http://www.edexcel.com) or [www.btec.co.uk](http://www.btec.co.uk). Alternatively, you can get in touch with us using the details on our contact us page at [www.edexcel.com/contactus](http://www.edexcel.com/contactus).

## **Pearson: helping people progress, everywhere**

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: [www.pearson.com/uk](http://www.pearson.com/uk)

January 2020

Publications Code 4CH1\_1C\_msc\_2020305

All the material in this publication is copyright

© Pearson Education Ltd 2020

### **General Marking Guidance**

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

Question number	Answer	Notes	Marks
1 (a)	(i) argon / helium	ACCEPT Ar / He	1
	(ii) nitrogen	ACCEPT N <sub>2</sub>	1
	(iii) carbon dioxide	ACCEPT CO <sub>2</sub>	1
	(iv) carbon dioxide	ACCEPT CO <sub>2</sub>	1
(b)	(i) $S + O_2 \rightarrow SO_2$		1
	(ii) acid rain	ACCEPT an adverse effect of acid rain e.g. kills fish, damages plants, corrodes limestone/marble buildings/statues etc.  IGNORE toxic/pollutant	1
			<b>Total 6</b>

Question number	Answer	Notes	Marks										
2 (a)	<table border="1" data-bbox="419 320 1046 813"> <tr> <td data-bbox="419 320 895 421">name of the part of the atom labelled Z</td> <td data-bbox="895 320 1046 421">nucleus</td> </tr> <tr> <td data-bbox="419 421 895 521">number of protons in this atom</td> <td data-bbox="895 421 1046 521">12</td> </tr> <tr> <td data-bbox="419 521 895 622">number of the group that contains this element</td> <td data-bbox="895 521 1046 622">2</td> </tr> <tr> <td data-bbox="419 622 895 723">number of the period that contains this element</td> <td data-bbox="895 622 1046 723">3</td> </tr> <tr> <td data-bbox="419 723 895 813">charge on the ion formed from this atom</td> <td data-bbox="895 723 1046 813">2+</td> </tr> </table>	name of the part of the atom labelled Z	nucleus	number of protons in this atom	12	number of the group that contains this element	2	number of the period that contains this element	3	charge on the ion formed from this atom	2+	<p data-bbox="1098 779 1315 813"><b>ACCEPT</b> +2 / Mg<sup>2+</sup></p>	5
name of the part of the atom labelled Z	nucleus												
number of protons in this atom	12												
number of the group that contains this element	2												
number of the period that contains this element	3												
charge on the ion formed from this atom	2+												
(b)	<ul data-bbox="435 987 1023 1115" style="list-style-type: none"> <li>• calculate sum of mass numbers multiplied by percentage abundances</li> <li>• divide answer by 100</li> <li>• give answer to one decimal place</li> </ul> <p data-bbox="379 1144 624 1178">Example calculation</p> <p data-bbox="379 1205 1015 1238"><b>M1</b> <math>(24 \times 79.2) + (25 \times 10.0) + (26 \times 10.8)</math> <b>OR</b> 2431.6</p> <p data-bbox="379 1335 762 1368"><b>M2</b> <math>2431.6 \div 100</math> <b>OR</b> 24.316</p> <p data-bbox="379 1603 496 1637"><b>M3</b> 24.3</p>	<p data-bbox="1098 1200 1358 1294"><b>REJECT</b> if correct working given but incorrectly evaluated</p> <p data-bbox="1098 1357 1342 1391"><b>ALLOW</b> ECF from M1</p> <p data-bbox="1098 1417 1350 1570"><math>(24 \times 0.792) + (25 \times 0.100) + (26 \times 0.108)</math> <b>OR</b> 24.316 with or without working scores <b>M1</b> and <b>M2</b></p> <p data-bbox="1098 1603 1374 1697"><b>ALLOW</b> ECF from M2 if calculated answer is to 1dp</p>	3										
			<b>Total 8</b>										

Question number	Answer	Notes	Marks
3 (a)	galvanising	ACCEPT galvanisation	1
(b) (i)	rust		1
(ii)	M1 oxygen / air  M2 water	ACCEPT O <sub>2</sub> IGNORE O  ACCEPT H <sub>2</sub> O/moisture  ACCEPT in either order	2
(c) (i)	(a reaction which) gives out / produces / releases heat (energy) / thermal energy	IGNORE energy without mention of heat or thermal	1
(ii)	An explanation that links the following two points  M1 aluminium/Al is more reactive than iron/Fe  M2 (because) aluminium/Al displaces iron/Fe (from its oxide)	ACCEPT aluminium/Al is higher in reactivity series than iron/Fe  ACCEPT reverse argument  ALLOW replaces/takes place of	2
(iii)	An explanation that links the following three points  M1 aluminium is oxidised and iron/iron oxide is reduced  M2 aluminium gains oxygen  M3 iron oxide/iron loses oxygen	ALLOW both oxidation and reduction occur  ALLOW aluminium/Al loses electrons  ALLOW iron <u>ions</u> /Fe <sup>3+</sup> gains electrons  ALLOW correct references to changes in oxidation number for M2 and M3	3
			<b>Total 10</b>



Question number	Answer	Notes	Marks
5 (a) (i)	S		1
	(ii) T and U		1
	(iii) U		1
(b)	<p>A description that makes reference to the following three points</p> <p><b>M1</b> (add) bromine water</p> <p><b>M2</b> no change / stays orange</p> <p><b>M3</b> (bromine water) decolourised / changes (from orange) to colourless</p>	<p><b>ACCEPT</b> Br<sub>2</sub> (aq)</p> <p><b>ALLOW</b> no reaction</p> <p>If initial colour of bromine water is given in M2 or M3 it must be correct -<b>ALLOW</b> any combination of orange/yellow/brown - but penalise once only</p> <p>If bromine given for M1 then in M2 and M3 allow any combination of red/orange/brown/yellow</p> <p><b>M2</b> and <b>M3</b> dep on bromine water/bromine in M1</p> <p>If no reagent and correct M2 and M3 - score 1</p> <p>if incorrect reagent and correct M2 and M3 score 0</p> <p><b>IGNORE</b> clear</p> <p><b>REJECT</b> discoloured</p> <p><b>ALLOW M1</b> acidified potassium manganate(VII)</p> <p><b>M2</b> no change/stays purple</p> <p><b>M3</b> decolourised / goes colourless</p>	3

Question number	Answer	Notes	Marks
5 (c)	<p>Any <b>two</b> of the following points</p> <p><b>M1</b> (can be represented by a) general formula</p> <p><b>M2</b> each member differs from the next by a CH<sub>2</sub> group OWTTE</p> <p><b>M3</b> (each member has) same functional group</p> <p><b>M4</b> (each member has) similar/same chemical properties / similar/same (chemical) reactions</p> <p><b>M5</b> trend in physical properties (between successive members)</p>	<p><b>ACCEPT</b> react in similar/same way</p> <p><b>ACCEPT</b> named physical property, e.g. boiling point</p> <p><b>REJECT</b> similar/same physical properties</p>	2
(d) (i)	<p>but-1-ene</p>	<b>ALLOW</b> 1-butene	1
(ii)	<p><b>Either</b></p> $  \begin{array}{cccc}  & \text{H} & \text{H} & \text{H} & \text{H} \\  &   &   &   &   \\  \text{H} & - \text{C} & - \text{C} & = \text{C} & - \text{C} - \text{H} \\  &   & & &   \\  & \text{H} & & & \text{H}  \end{array}  $ <p><b>Or</b></p> $  \begin{array}{c}  \text{H} & & & \text{H} \\    & & & / \\  \text{H} - \text{C} & - & \text{C} = & \text{C} \\    & & & \backslash \\  \text{H} & & & \text{H} \\    & & & \\  \text{H} - \text{C} & - & \text{H} \\    \\  \text{H}  \end{array}  $	<p><b>ACCEPT</b> cis or trans isomer</p> <p><b>REJECT</b> displayed formulae of cyclic alkanes</p>	1

Question number	Answer	Notes	Marks												
5 (e) (i)	<ul style="list-style-type: none"> <li>• Divide percentages by relative atomic masses</li> <li>• Divide results by smallest value to obtain ratio</li> </ul> <p>Example calculation</p> <p><b>M1</b></p> <table style="margin-left: 40px;"> <tr> <td>C</td> <td>H</td> <td>F</td> </tr> <tr> <td><math>\frac{36.36}{12}</math></td> <td><math>\frac{6.06}{1}</math></td> <td><math>\frac{57.58}{19}</math></td> </tr> </table> <p><b>M2</b></p> <table style="margin-left: 40px;"> <tr> <td><math>\frac{3.03}{3.03}</math></td> <td><math>\frac{6.06}{3.03}</math></td> <td><math>\frac{3.03}{3.03}</math></td> </tr> </table> <p><b>OR</b></p> <table style="margin-left: 40px;"> <tr> <td>1</td> <td>2</td> <td>1</td> </tr> </table>	C	H	F	$\frac{36.36}{12}$	$\frac{6.06}{1}$	$\frac{57.58}{19}$	$\frac{3.03}{3.03}$	$\frac{6.06}{3.03}$	$\frac{3.03}{3.03}$	1	2	1	0 marks if division by atomic numbers or upside down calculation	2
C	H	F													
$\frac{36.36}{12}$	$\frac{6.06}{1}$	$\frac{57.58}{19}$													
$\frac{3.03}{3.03}$	$\frac{6.06}{3.03}$	$\frac{3.03}{3.03}$													
1	2	1													
(ii)	<ul style="list-style-type: none"> <li>• divide relative molecular mass by empirical formula mass</li> <li>• correct molecular formula</li> </ul> <p>Example calculation</p> <p><b>M1</b></p> <table style="margin-left: 40px;"> <tr> <td><math>\frac{66}{12 + 2 + 19}</math></td> <td><b>OR</b></td> <td><math>\frac{66}{33}</math></td> <td><b>OR</b></td> <td>2</td> </tr> </table> <p><b>M2</b></p> <p>C<sub>2</sub>H<sub>4</sub>F<sub>2</sub></p>	$\frac{66}{12 + 2 + 19}$	<b>OR</b>	$\frac{66}{33}$	<b>OR</b>	2	<p><b>ACCEPT</b> symbols in any order</p> <p>correct answer without working scores 2 marks.</p> <p>2CH<sub>2</sub>F scores 1</p>	2							
$\frac{66}{12 + 2 + 19}$	<b>OR</b>	$\frac{66}{33}$	<b>OR</b>	2											
<b>Total 14</b>															

Question number	Answer	Notes	Marks						
6 (a)	zinc + hydrochloric acid → zinc chloride + hydrogen	<b>ACCEPT</b> fully correct chemical equation	1						
(b)	<table border="1" data-bbox="419 383 1011 640"> <tr> <td data-bbox="419 383 858 483">temperature in °C after adding zinc</td> <td data-bbox="858 383 1011 483">22.4</td> </tr> <tr> <td data-bbox="419 483 858 584">temperature in °C before adding zinc</td> <td data-bbox="858 483 1011 584">17.7</td> </tr> <tr> <td data-bbox="419 584 858 640">temperature change in °C</td> <td data-bbox="858 584 1011 640">4.7</td> </tr> </table> <p data-bbox="379 674 480 707">M1 22.4</p> <p data-bbox="379 734 480 768">M2 17.7</p> <p data-bbox="379 795 501 828">M3 (+)4.7</p>	temperature in °C after adding zinc	22.4	temperature in °C before adding zinc	17.7	temperature change in °C	4.7	<p data-bbox="1110 383 1393 506">If readings are correct but in wrong order award 1 mark for M1 and M2</p> <p data-bbox="1110 864 1377 1043"><b>ALLOW</b> ECF for M3 if M1 and/or M2 incorrect If answers not given to nearest 0.1 °C penalise once only</p>	3
temperature in °C after adding zinc	22.4								
temperature in °C before adding zinc	17.7								
temperature change in °C	4.7								
(c) (i)	<p data-bbox="379 1081 1070 1115">An explanation that links any <b>two</b> of the following points</p> <p data-bbox="379 1144 743 1178">M1 polystyrene is an insulator</p> <p data-bbox="379 1328 687 1361">M2 (so) reduces heat loss</p> <p data-bbox="379 1451 1038 1507">M3 temperature rise/change/reading will be closer to true value OWTTE</p>	<p data-bbox="1110 1144 1361 1290"><b>ALLOW</b> is not a (good) conductor of heat <b>ALLOW</b> is a poor conductor of heat</p> <p data-bbox="1110 1328 1385 1417"><b>ALLOW</b> prevents heat loss <b>ALLOW</b> keeps heat in</p> <p data-bbox="1110 1451 1369 1574"><b>ALLOW</b> temperature rise/change/reading will be more accurate/valid</p>	2						
(ii)	<p data-bbox="379 2047 568 2080">Any <b>three</b> from</p> <p data-bbox="379 2114 699 2148">M1 amount/mass of metal</p> <p data-bbox="379 2175 683 2208">M2 concentration of acid</p> <p data-bbox="379 2235 603 2240">M3 volume of acid</p> <p data-bbox="379 2296 711 2240">M4 (speed/time of) stirring</p> <p data-bbox="379 2356 778 2240">M5 external / room temperature</p>	<p data-bbox="1110 2114 1382 2181"><b>ALLOW</b> size / surface area of metal</p> <p data-bbox="1110 2235 1342 2240"><b>ALLOW</b> amount of acid</p> <p data-bbox="1110 2356 1385 2240"><b>ALLOW</b> initial /starting temperature</p>	3						

Question number	Answer	Notes	Marks
6 (d)	(i) no reaction (occurred between copper and hydrochloric acid)	<b>IGNORE</b> copper is unreactive <b>ALLOW</b> copper is less reactive than hydrogen	1
	(ii) Any value between 1.5 and 5.0 °C inclusive		1
	(iii) most reactive magnesium zinc iron tin copper	<b>ACCEPT</b> symbols	1
			<b>Total</b> 12

Question number	Answer	Marks
7 (a)	(i) <b>B</b> bromine	1
	(ii) A is incorrect as astatine is a solid C is incorrect as chlorine is a gas D is incorrect as iodine is a solid	1
	(iii) <b>C</b> chlorine (as it is pale green)  A is incorrect as astatine is black B is incorrect as bromine is brown C is incorrect as iodine is dark grey  <b>A</b> astatine  B is incorrect as bromine is more reactive than astatine C is incorrect as chlorine is more reactive than astatine D is incorrect as iodine is more reactive than astatine	1

(b) (i)	<p><b>M1</b> (colourless solution turns) brown</p> <p><b>M2</b> (solution stays) brown / no change</p>	<b>ALLOW</b> no reaction	2
(ii)	bromine would not react with (sodium) bromide / bromine cannot displace itself OWTTE	<p><b>ALLOW</b> bromine cannot react with itself</p> <p><b>ALLOW</b> both contain bromine/same element/same halogen</p> <p><b>ALLOW</b> because no reaction would occur</p> <p><b>REJECT</b> bromine cannot displace bromide</p>	1
(iii)	$\text{Br}_2 + 2\text{NaI} \rightarrow 2\text{NaBr} + \text{I}_2$	<p><b>ACCEPT</b> correct ionic equation</p> $\text{Br}_2 + 2\text{I}^- \rightarrow 2\text{Br}^- + \text{I}_2$ <p><b>ALLOW</b> multiples and fractions</p>	1

Question number	Answer	Notes	Marks
7 (c)	<p>A description that makes reference to the following 6 points</p> <p>Test for cation</p> <p><b>M1</b> add sodium hydroxide (solution)</p> <p><b>M2</b> if blue precipitate forms solution contains copper(II) ion(s) / contains <math>\text{Cu}^{2+}</math> / is a copper compound</p> <p><b>M3</b> if green precipitate forms solution contains iron(II) ion(s) / contains <math>\text{Fe}^{2+}</math> / is an iron compound</p>	<p><b>ALLOW</b> ammonia solution</p> <p><b>IGNORE</b> qualifiers <b>REJECT</b> other colours</p> <p><b>IGNORE</b> qualifiers <b>REJECT</b> other colours</p> <p>If no reagent or incorrect reagent but correct M2 and M3 score 1</p> <p><b>ALLOW</b> <b>M1</b> flame test or description of flame test</p> <p><b>M2</b> if blue-green (flame) solution contains copper(II) ion(s) / contains <math>\text{Cu}^{2+}</math> / is a copper compound</p> <p>No <b>M3</b> for this test</p> <p><b>ALLOW</b> <b>M1</b> addition of suitable metal above Cu in reactivity series</p> <p><b>M2</b> brown/pink/pink-brown solid forms</p> <p>No <b>M3</b> for this test</p>	6


	<p>Test for anion</p> <p><b>M4</b> add silver nitrate (solution)</p> <p><b>M5</b> if white precipitate forms solution contains chloride ion(s) / contains <math>\text{Cl}^-</math> / is a chloride</p> <p><b>M6</b> if cream precipitate forms solution contains bromide ion(s) / contains <math>\text{Br}^-</math> / is a bromide</p>	<p><b>IGNORE</b> addition of nitric acid</p> <p><b>REJECT</b> addition of hydrochloric acid for <b>M4</b></p> <p>If no reagent or incorrect reagent but correct M5 and M6 score 1</p> <p><b>ALLOW</b> <b>M4</b> add chlorine water (to solution)</p> <p><b>M5</b> if turns orange/yellow/brown solution contains bromide ion(s) / contains <math>\text{Br}^-</math> / is a bromide</p> <p>No <b>M6</b> for this test</p>	<p><b>Total 13</b></p>
--	--	--	------------------------

Question number	Answer	Notes	Marks
8 (a) (i)	sublimation / subliming		1
	(ii) <b>M1</b> (add to/bubble into) limewater <b>M2</b> (limewater) turns cloudy/milky	<b>ACCEPT</b> forms white precipitate <b>M2 DEP M1</b>	2
	(b) An explanation that links the following two points  <b>M1</b> weak forces (of attraction) between molecules / weak intermolecular forces (of attraction)  <b>M2</b> little energy needed to overcome the (intermolecular) forces	<b>ALLOW</b> weak intermolecular bonds <b>ALLOW</b> weak intermolecular attractions  <b>IGNORE</b> less energy  <b>ALLOW</b> little energy needed to separate the molecules  <b>M2 DEP M1</b> correct or missing	2

(c)	<p>Any explanation that links any <b>three</b> of the following points for diamond</p> <p><b>M1</b> each (carbon) atom is (covalently) bonded to four other (carbon) atoms</p> <p><b>M2</b> in a (giant) tetrahedral lattice /network / structure</p> <p><b>M3</b> the (covalent) bonds are (very) strong</p> <p><b>M4</b> (therefore) diamond is (very) hard (and so good for cutting tools)</p> <p>Any explanation that links any <b>three</b> of the following points for graphite</p> <p><b>M5</b> each (carbon) atom is (covalently) bonded to three other (carbon) atoms</p> <p><b>M6</b> (the structure is) in layers</p> <p><b>M7</b> weak forces (between layers)</p> <p><b>M8</b> (the layers can) slide over each other/ rub off</p> <p><b>M9</b> this makes graphite soft (so it can make marks on paper)</p>	<p><b>ALLOW</b> each carbon has four bonds</p> <p><b>ALLOW</b> 3D/rigid in place of tetrahedral</p> <p><b>ALLOW</b> reference to lot of energy needed to break the (covalent) bonds</p> <p><b>ALLOW</b> there are lots of/many (covalent) bonds</p> <p><b>ALLOW</b> diamond is (very) strong</p> <p>If mention of intermolecular forces in diamond MAX 2 for diamond</p> <p>If mention of ions in diamond only M4 can be scored</p> <p><b>ALLOW</b> sheets</p> <p><b>ALLOW</b> slippery</p> <p>If mention of intermolecular forces in graphite MAX 2 for graphite</p> <p>If mention of ions in graphite only M9 can be scored</p>	<p>6</p> <p><b>Total 11</b></p>
-----	---	---	---------------------------------

Question number	Answer	Notes	Marks
9 (a)	to minimise/prevent (mass loss by) evaporation of the (liquid) fuel OWTTE	<b>ALLOW</b> to find mass of fuel used/burned	1
(b) (i)	soot/carbon	<b>REJECT</b> copper oxide	1
(ii)	An explanation that links the following two points.  M1 incomplete combustion (occurs)  M2 (because) the air/oxygen supply is limited OWTTE	<b>ALLOW</b> mark for soot/carbon if not seen in (i), unless copper oxide is mentioned in (i)  If copper oxide in (i) <b>ALLOW</b> 1 mark for (because) copper reacts with oxygen (in air)	2
(c) (i)	<ul style="list-style-type: none"> <li>• substitution into <math>Q = mc\Delta T</math></li> <li>• calculation of heat energy in Joules</li> <li>• conversion to kJ</li> </ul> <p>Example calculation</p> <p>M1 <math>Q = 100 \times 4.2 \times 30</math></p> <p>M2 = 12600 (J)</p> <p>M3 = 12.6 kJ</p>	<p>12600 (J) with no working scores M1 and M2 M2 ECF M1</p> <p><b>ALLOW</b> approximately = 13 kJ</p> <p>12.6 kJ with no working scores 3</p>	3

(ii)	<ul style="list-style-type: none"><li>• calculate the amount, in moles, of methanol</li><li>• divide Q by the amount in moles</li><li>• give the answer with the correct sign</li></ul> <p>Example calculation</p> <p><b>M1</b> <math>0.96 \div 32</math> <b>OR</b> <math>0.03</math></p> <p><b>M2</b> <math>12.6 \div 0.03</math> <b>OR</b> <math>420</math> (kJ/mol)</p> <p><b>M3</b> <math>- 420</math> (kJ/mol)</p>	<p><b>ACCEPT</b> <math>13 \div 0.03</math> <b>OR</b> <math>430/433</math> for <b>M2</b></p> <p><b>AND</b> <math>- 430 / - 433</math> for <b>M3</b></p>	3
------	---	--	---

Question number	Answer	Notes	Marks
9 (d) (i)	<p><b>M1</b> all points plotted correctly</p> <p><b>M2</b> line of best fit drawn with a ruler</p> 	does not need to start at (0,0)	2
(ii)	<p><b>M1</b> straight line extrapolated up to 6 carbon atoms</p> <p><b>M2</b> value of <math>\Delta H</math> read from their graph</p>	<p><b>ALLOW</b> extra point shown at 6 carbon atoms</p> <p>negative sign needed</p>	2
(iii)	The greater the number of carbon atoms (per molecule) the greater (the magnitude/ value of) $\Delta H$	<p><b>ALLOW</b> <math>\Delta H</math> is (directly) proportional to the number of carbon atoms per molecule</p> <p><b>ALLOW</b> The greater the number of carbon atoms (per molecule) the more exothermic the <math>\Delta H</math> value</p>	1
			<b>Total 15</b>

Question number	Answer	Notes	Marks
10 (a) (i)	$4\text{NH}_3 + 5\text{O}_2 \rightleftharpoons 4\text{NO} + 6\text{H}_2\text{O}$	<b>ACCEPT</b> multiples and fractions	1
(ii)	reversible (reaction)	<b>ACCEPT</b> reaction that goes both ways / both forwards and backwards reactions occur  <b>IGNORE</b> references to equilibrium	1
(iii)	to increase the rate of the reaction / to speed up the reaction OWTTE	<b>IGNORE</b> references to lowering the activation energy	1
(b)	$2\text{NO} + \text{O}_2 \rightarrow 2\text{NO}_2$	<b>ACCEPT</b> multiples and fractions	1
(c) (i)	<ul style="list-style-type: none"> <li>• calculate <math>M_r</math> of <math>\text{NO}_2</math> and <math>\text{HNO}_3</math></li> <li>• calculate the amount, in moles, of <math>\text{NO}_2</math></li> <li>• calculate the amount, in moles, of <math>\text{HNO}_3</math></li> <li>• calculate the mass in tonnes of <math>\text{HNO}_3</math></li> </ul> <p>Example calculation</p> <p><b>M1</b> <math>M_r</math> of <math>\text{NO}_2 = 46</math> <math>M_r</math> of <math>\text{HNO}_3 = 63</math></p> <p><b>M2</b> <math>n(\text{NO}_2) = 11.5 \times 10^6 \div 46</math> OR 250 000 (mol)</p> <p><b>M3</b> <math>n(\text{HNO}_3) = \frac{2 \times 25\,0000}{3}</math> OR 167 000 / 170 000</p> <p><b>M4</b> <math>(167\,000 \times 63 \text{ g}) = 10.5</math> (tonnes)</p>	<p><b>ALLOW</b> working in megamoles i.e. <math>11.5 \div 46</math> OR 0.25</p> <p><b>ALLOW</b> ECF from incorrect <math>M_r</math> of <math>\text{NO}_2</math></p> <p>calculator answer 166666.66 <b>ALLOW</b> working in megamoles i.e. <math>\frac{2 \times 0.25}{3}</math> OR 0.167 / 0.17</p> <p><b>ALLOW</b> ECF from M2</p> <p>10.5 (tonnes) with no working scores 4</p> <p><b>ACCEPT</b> 10.7 (if 170 000 used)</p> <p><b>ALLOW</b> ECF from M3 <b>ALLOW</b> ECF from incorrect <math>M_r</math> of <math>\text{HNO}_3</math></p>	4
(ii)	can be (re)used in stage 2 / to make more nitrogen dioxide (in stage 2) / can be used to make more nitric acid	<b>IGNORE</b> can be recycled/reused unless qualified	1

Question number	Answer	Notes	Marks
10 (d)	<ul style="list-style-type: none"> <li>• calculate the amount, in moles, of copper(II) nitrate</li> <li>• calculate the theoretical yield, in moles, of copper(II) nitrate</li> <li>• calculate the percentage yield</li> </ul> <p>Example calculation</p> <p><b>M1</b> <math>n\text{Cu}(\text{NO}_3)_2</math> formed = <math>15.3 \div 187.5</math> <b>OR</b> 0.0816</p> <p><b>M2</b> theoretical <math>n\text{Cu}(\text{NO}_3)_2</math> = <math>0.200 \div 2</math> <b>OR</b> 0.100</p> <p><b>M3</b> (% yield) = <math>\frac{(0.0816 \times 100)}{(0.100)} = 81.6</math> (%)</p> <p>Alternative method</p> <ul style="list-style-type: none"> <li>• calculate the theoretical yield, in moles, of copper(II) nitrate</li> <li>• calculate the theoretical mass of copper nitrate that should be formed</li> <li>• calculate the percentage yield</li> </ul> <p>Example calculation</p> <p><b>M1</b> theoretical <math>n\text{Cu}(\text{NO}_3)_2</math> = <math>0.200 \div 2</math> <b>OR</b> 0.100</p> <p><b>M2</b> theoretical mass of copper nitrate = <math>0.1 \times 187.5 = 18.75</math></p> <p><b>M3</b> (% yield) = <math>\frac{15.3}{18.75} \times 100 = 81.6</math> (%)</p>	<p><b>ALLOW</b> 0.082</p> <p><b>ACCEPT</b> 82 (%)</p> <p>Mark M3 CSQ on M1 and M2</p> <p>40.8 scores 2</p> <p><b>ALLOW</b> 18.8</p> <p><b>ACCEPT</b> 82 (%)</p> <p>Mark M3 CSQ on M1 and M2</p> <p>40.8 scores 2</p> <p>81.6(%) with no working scores 3 marks</p>	<p>3</p> <p><b>Total 12</b></p>

