



Mark Scheme (Results)

January 2014

International GCSE
Physics (4PH0) Paper 2P

Edexcel Level 1/Level 2 Certificates
Physics (KPH0) Paper 2P

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

January 2014

Publications Code UG037827

All the material in this publication is copyright

© Pearson Education Ltd 2014

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

Question number	Answer	Notes	Marks
1 (a) (i)	B (53)		1
	(ii) D (131)		1
1 (b)	Any two of - MP1 Beta is (moderately) ionising; MP2 Beta has a short range; MP3 idea that I-131 has a short half-life; MP4 idea that iodine is absorbed (easily) by the thyroid; MP5 (hence) reduces damage to healthy cells; MP6 (hence) does not penetrate out of the body; MP7 (therefore) kills (only) tumour cells;	Ignore I-131 is radioactive, it emits beta	2

Total 4 marks


Question number	Answer	Notes	Marks																					
2 (a)	Similarity - both have magnitude/OWTTE; Difference - vector has direction OR scalar does not have direction;		2																					
(b)	<table border="1" style="margin-left: auto; margin-right: auto;"> <thead> <tr> <th>Quantity</th> <th>Scalar</th> <th>Vector</th> </tr> </thead> <tbody> <tr> <td>Density</td> <td></td> <td></td> </tr> <tr> <td>energy</td> <td></td> <td></td> </tr> <tr> <td>force</td> <td></td> <td></td> </tr> <tr> <td>momentum</td> <td></td> <td></td> </tr> <tr> <td>speed</td> <td></td> <td></td> </tr> <tr> <td>velocity</td> <td></td> <td></td> </tr> </tbody> </table> <p>One or two correct ticks = 1 mark Three or four correct ticks = 2 marks All five correct ticks = 3 marks</p>	Quantity	Scalar	Vector	Density			energy			force			momentum			speed			velocity			Ignore density (already completed)	3
Quantity	Scalar	Vector																						
Density																								
energy																								
force																								
momentum																								
speed																								
velocity																								

Total 5 marks

Question number	Answer	Notes	Marks
3 (a) (i)	smoke particles in air (in smoke cell) OR pollen on water OR dust particles in air;	Accept correct description of Brownian motion applied to unspecified particles in a suitable medium	1
	(ii) Any two of - MP1 Idea that tiny/smaller particles are hitting; MP2 Larger (observed) particles are moved; MP3 Idea of random motion of larger particles;	Allow zig-zag movement	2
3 (b)	Any six ideas about arrangement and motion of particles Max 2 for each state Solid – Regular pattern OR close packed; Vibration in position; Little space between particles; Liquid – Irregular pattern; Able to move over/pace other particles; Little space between particles; Gas – No pattern; Able to move freely/faster; Larger space between particles;	Accept same ideas shown in labelled diagrams Condone fixed position Condone no fixed position Ignore vibration relating to liquid Condone no fixed position Ignore vibration relating to gas	max 6

Total 9 marks

Question number	Answer	Notes	Marks
4 (a)	Any three of - MP1 use a stirrer / stir with thermometer; MP2 centralise / spread heat source; MP3 move thermistor and thermometer to same level; MP4 move thermistor and thermometer closer together; MP5 Use thermometer with finer scale / digital thermometer;	Ignore repeat readings Assume horizontal separation meant	Max 3
(b)	(milli)Ammeter;	Allow ampmeter	1
(c) (i)	Scale; (at least half the grid) Axes labelled including units; Plotting $\pm \frac{1}{2}$ small square;; Line of best fit;	Accept axes reversed -1 each plotting error, minimum 0 for plotting Curve through either (80, 0.2) or (100, 0.4) Allow line bisecting these two points	5
(c) (ii)	DOP (80, 0.2) circled (if supported by line of best fit)	Allow (100, 0.4) circled if supported by line of best fit	1



Temperature in °C	Voltage in V
20	6.0
40	2.2
60	1.1
80	0.2
100	0.4

Question number	Answer	Notes	Marks
4 (d) (i)	voltage = current x resistance;	Accept rearrangements and symbols e.g. current = voltage ÷ resistance, $V=IR$, $R=V/I$	1
(ii)	Substitution into correctly rearranged equation; Conversion between amps and milliamps; Calculation yielding value correct to at least 2 s.f.; e.g. $I = 5.9 \div 680$ $= 0.00868 \text{ (A)}$ $= 8.7 \text{ (mA)}$	Accept x 1000 in calculation Allow 1 mark max if response is only a successful reverse argument leading to 5.8 V or 5.78 V	3

Total 14 marks

Question number	Answer	Notes	Marks
5	any four from - MP1 momentum reduced; MP2 by same amount; MP3 over longer time; MP4 so force reduced; MP5 use of "force = rate of change of momentum"; MP6 less force means less damage/injuries;	Responses should be in the context of momentum ignore "momentum absorbed" ignore "impact reduced" simple mention of eqn is insufficient	max 4

Total 4 marks

Question number	Answer	Notes	Marks
6 (a) (i)	set-up showing any two from-clear indication of equipment needed; correct refraction at one surface of glass block shown; protractor shown in use;	ray-box or pins Allow ruler for apparent depth method	2
(ii)	angle of incidence; angle of refraction; OR critical angle; idea of grazing emergence;	Allow apparent depth method, i.e. real depth; apparent depth;	2
(iii)	find $\sin i$ and $\sin r$; refractive index is the ratio of sines; OR find $\sin c$; refractive index is $\frac{1}{\sin c}$;	Accept for two marks <ul style="list-style-type: none"> • $(n =) \sin i / \sin r$ • $(n =) 1 / \sin c$ • graph of $\sin i$ vs $\sin r$ Allow refractive index = real depth \div apparent depth for two marks	2
(b) (i)	Diagram – reflection at first back surface; reflection at second back surface;	judge by eye <ul style="list-style-type: none"> • straightness of ray and correctness of angle • emergent ray parallel to incident ray 	2
(ii)	Refracted / slows down / wavelength decreases	Ignore: direction change ideas it does nothing / nothing happens	1

Total 9 marks

Question number	Answer	Notes	Marks
7 (a)	Rods magnetised; And repel;	Reject ideas of charge for one mark only	2
(b)	MP1. A named magnetic material e.g.(soft) iron; MP2. because the material is capable of being magnetised; MP3. DOP (iron only) but does not retain its magnetism;	ACCEPT steel, mu-metal, nickel, cobalt accept RA steel would stay magnetised/apart	3
(c)	any two from- MP1. field (in coil) switches polarity; MP2. field (in rods) weaker; MP3. (since) field alternates with current or at 50 Hz; MP4. rods may not have time to become fully magnetised;	allow <ul style="list-style-type: none"> • 100 times a second or mains frequency • hysteresis ideas • domain theory • reluctance ideas 	2

Total 7 marks

Question number	Answer	Notes	Marks
8 (a)	weight of (the) plank		1
8 (b)	(i) moment = force x (perpendicular) distance (from pivot) (ii) substitution; final value; e.g. 1200×0.75 900 (Nm)		1 2
(c)	principle of moments (stated or implied); correct calculation of distance from hand to pivot; calculation of total anticlockwise moment; final value; e.g. $(F \times 2.25) + (200 \times 0.75) = (1200 \times 0.75)$ $F = 330 \text{ (N)}$	Allow ecf from (b) 2.25 (m) seen in working $(F \times 2.25) + (200 \times 0.75)$ Allow 333 N	4

Total 8 marks

Pearson Education Limited. Registered company number 872828
with its registered office at Edinburgh Gate, Harlow, Essex CM20 2JE