GradeMax

Cambridge International AS & A Level

Cambridge International Examinations
Cambridge International Advanced Subsidiary and Advanced Level

CANDIDATE NAME

CENTRE NUMBER

CANDIDATE NUMBER

* 9 0 8 0 4 1 9 4 3 *

CHEMISTRY **9701/35**

Paper 3 Advanced Practical Skills 1 **October/November 2015**

2 hours

Candidates answer on the Question Paper.

Additional Materials: As listed in the Confidential Instructions

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in.
Give details of the practical session and laboratory where appropriate, in the boxes provided.
Write in dark blue or black pen.
You may use an HB pencil for any diagrams or graphs.
Do not use staples, paper clips, glue or correction fluid.
DO **NOT** WRITE IN ANY BARCODES.

Answer **all** questions.
Electronic calculators may be used.
You may lose marks if you do not show your working or if you do not use appropriate units.
Use of a Data Booklet is unnecessary.
A copy of the Periodic Table is printed on page 12.

Qualitative Analysis Notes are printed on pages 10 and 11.

At the end of the examination, fasten all your work securely together.
The number of marks is given in brackets [ ] at the end of each question or part question.

Session
Laboratory

For Examiner's Use	
1	
2	
3	
Total	

This document consists of **12** printed pages and **1** insert.

© UCLES 2015

CAMBRIDGE International Examinations

GradeMax

1 In this experiment you will determine the ionic equation for the reaction of acidified potassium manganate(VII) with potassium iodide. Excess potassium iodide is used and the reaction produces iodine. The amount of iodine produced is measured by titration with sodium thiosulfate.

FA 1 is 0.0180 mol dm^{-3} potassium manganate(VII), $KMnO_4$.
FA 2 is 1.00 mol dm^{-3} sulfuric acid, H_2SO_4.
FA 3 is 0.500 mol dm^{-3} potassium iodide, KI.
FA 4 is 0.100 mol dm^{-3} sodium thiosulfate, $Na_2S_2O_3$.
starch indicator

(a) Method

- Pipette 25.0 cm^3 of **FA 1** into a conical flask.
- Use the measuring cylinder to add 25 cm^3 of **FA 2** to the conical flask.
- Use the measuring cylinder to add 20 cm^3 of **FA 3** to the conical flask.
- Fill the burette with **FA 4**.
- Carry out a rough titration. When the colour of the mixture becomes yellow/orange, add a few drops of starch indicator. Then titrate until the mixture goes colourless.
- Record all your burette readings in the space below.

The rough titre is ..... cm^3.

- Carry out as many accurate titrations as you think necessary to obtain consistent results.
- Make sure any recorded results show the precision of your practical work.
- Record in a suitable form below all of your burette readings and the volume of **FA 4** added in each accurate titration.

Keep FA 1 and FA 2 for use in Question 3 and FA 4 for use in Question 2.

I	
II	
III	
IV	
V	
VI	
VII	

[7]

(b) From your accurate titration results, obtain a suitable value for the volume of **FA 4** to be used in your calculations.
Show clearly how you have obtained this value.

Volume of **FA 4** required is ..... cm^3. [1]

© UCLES 2015

(c) Calculations

Show your working and appropriate significant figures in the final answer to **each** step of your calculations.

(i) Calculate the number of moles of sodium thiosulfate in the volume of **FA 4** calculated in **(b)**.

moles of $Na_2S_2O_3$ = ... mol

(ii) Use the equation below to calculate the number of moles of iodine that reacted with the sodium thiosulfate in the titration.

$$I_2 + 2Na_2S_2O_3 \rightarrow Na_2S_4O_6 + 2NaI$$

moles of I_2 = ... mol

(iii) Use information on page 2 to calculate the number of moles of potassium manganate(VII) in **FA 1** used in the titration.

moles of $KMnO_4$ = ... mol

(iv) From your answers to **(ii)** and **(iii)**, calculate the number of moles of iodine produced by the reaction of **2.00** moles of potassium manganate(VII) with excess potassium iodide.

moles I_2 = ... mol

(v) Using your answer to **(iv)**, put a tick next to the ionic equation that represents the reaction between **FA 1** and **FA 3**.

$2MnO_4^- + 2I^- + 16H^+ \rightarrow I_2 + 2Mn^{6+} + 8H_2O$ .....

$2MnO_4^- + 4I^- + 16H^+ \rightarrow 2I_2 + 2Mn^{5+} + 8H_2O$ .....

$2MnO_4^- + 6I^- + 16H^+ \rightarrow 3I_2 + 2Mn^{4+} + 8H_2O$ .....

$2MnO_4^- + 8I^- + 16H^+ \rightarrow 4I_2 + 2Mn^{3+} + 8H_2O$ .....

$2MnO_4^- + 10I^- + 16H^+ \rightarrow 5I_2 + 2Mn^{2+} + 8H_2O$ .....

$2MnO_4^- + 12I^- + 16H^+ \rightarrow 6I_2 + 2Mn^{+} + 8H_2O$ .....

© UCLES 2015

GradeMax

(vi) Prove that the iodide ion has been oxidised in the equation that you selected in **(v)**.

.....

.....

.....

[5]

(d) (i) The error in calibration of the pipette you used is $\pm 0.06\,cm^3$.
Calculate the percentage error when measuring **FA 1**, using the pipette.

percentage error = ... %

(ii) A student suggested that the experiment would be more accurate if a pipette was used to measure solution **FA 3**.
State and explain whether you agree with the student.

.....

.....

.....

[2]

[Total: 15]

© UCLES 2015

GradeMax

2 In this experiment you will investigate how the rate of reaction between sodium thiosulfate and hydrochloric acid is affected by the concentration of the acid.

When aqueous thiosulfate ions react with hydrogen ions, H^+, in any acid, a pale yellow precipitate of sulfur is formed. The ionic equation for this reaction is given below.

$$S_2O_3^{2-}(aq) + 2H^+(aq) \rightarrow S(s) + SO_2(aq) + H_2O(l)$$

The rate of the reaction can be determined by measuring the time taken to produce a fixed quantity of sulfur.

FA 4 is 0.10 mol dm^{-3} sodium thiosulfate, $Na_2S_2O_3$.
FA 5 is 0.20 mol dm^{-3} hydrochloric acid, HC*l*.

(a) Method

Record **all** your measurements, in an appropriate form, in the space below.

Experiment 1

- Use the larger measuring cylinder to transfer 40 cm^3 of **FA 4** into the 100 cm^3 beaker.
- Rinse the larger measuring cylinder thoroughly with water, then add 30 cm^3 of **FA 5** to the beaker and start timing **immediately**.
- Stir the mixture once and place the beaker on top of the printed insert page provided.
- Look down through the solution in the beaker at the print on the insert.
- Stop timing as soon as the precipitate of sulfur makes the print on the insert invisible.
- Record the reaction time to the **nearest second**.
- Empty and rinse the 100 cm^3 beaker.
- Dry the outside of the beaker ready for Experiment 2.

Experiment 2

- Rinse the larger measuring cylinder, then use it to transfer 40 cm^3 of **FA 4** into the 100 cm^3 beaker.
- Use the smaller measuring cylinder to add 10 cm^3 of distilled water to the beaker.
- Use the same measuring cylinder to add 20 cm^3 of **FA 5** to the mixture in the beaker and start timing **immediately**.
- Stir the mixture once and place the beaker on top of the printed insert page provided.
- Stop timing as soon as the print on the insert becomes invisible.
- Record the reaction time to the **nearest second**.
- Empty and rinse the 100 cm^3 beaker.
- Dry the outside of the beaker ready for Experiment 3.

Experiment 3

- Carry out the reaction using a mixture of 40 cm^3 of **FA 4**, 20 cm^3 of distilled water and 10 cm^3 of **FA 5**.
- Measure and record the reaction time to the **nearest second**.

I	
II	
III	
IV	

[4]

© UCLES 2015

[Turn over

(b) (i) The 'rate of reaction' can be represented by the formula below.

$$\text{'rate of reaction'} = \frac{1000}{\text{reaction time}}$$

Use this formula to calculate the 'rate of reaction' for Experiments 1 and 3.
Give the unit.

'rate of reaction' for Experiment 1 ..... unit .....

'rate of reaction' for Experiment 3 ..... unit .....

(ii) Calculate the initial concentrations of hydrochloric acid in the reaction mixtures in Experiments 1 and 3.

initial concentration of HC*l* in Experiment 1 = ... mol dm^{-3}

initial concentration of HC*l* in Experiment 3 = ... mol dm^{-3}

(iii) How is the 'rate of reaction' affected by the concentration of hydrochloric acid in the mixture?

.....

.....

(iv) Predict how the reaction time measured in Experiment 1 would have been affected if the experiment had been carried out using 0.20 mol dm^{-3} sulfuric acid instead of 0.20 mol dm^{-3} hydrochloric acid.
Explain your answer.

.....

.....

.....

(v) Predict how the reaction time measured in Experiment 3 would have been affected if the experiment had been carried out in a 250 cm^3 beaker instead of a 100 cm^3 beaker.
Explain your answer.

.....

.....

[5]

[Total: 9]

3 Qualitative Analysis

At each stage of any test you are to record details of the following.

- colour changes seen
- the formation of any precipitate
- the solubility of such precipitates in an excess of the reagent added

Where gases are released they should be identified by a test, **described in the appropriate place in your observations**.

You should indicate clearly at what stage in a test a change occurs.
No additional tests for ions present should be attempted.

If any solution is warmed, a boiling tube MUST be used.

Rinse and reuse test-tubes and boiling tubes where possible.

Where reagents are selected for use in a test, the name or correct formula of the element or compound must be given.

(a) **FA 6** is a sodium compound containing one anion listed on page 11.

Dissolve the **FA 6** provided in about 15 cm^3 of distilled water in a boiling tube.
Carry out the following tests and record your observations in the table below.

test		*observations*
(i)	To a 1cm depth of the solution of **FA 6** in a test-tube, add a few drops of aqueous barium chloride or aqueous barium nitrate, then	
	add dilute hydrochloric acid.	
(ii)	To a 1cm depth of the solution of **FA 6** in a test-tube, add an equal volume of aqueous hydrogen peroxide, then	
	add a few drops of aqueous barium chloride or aqueous barium nitrate, then	
	add dilute hydrochloric acid.	

GradeMax

	test	*observations*
(iii)	To a 1 cm depth of the solution of **FA 6** in a boiling tube, add an equal volume of **FA 2**, sulfuric acid, then	
	heat the mixture **gently and cautiously**.	
(iv)	To a 1 cm depth of the solution of **FA 6** in a test-tube, add an equal volume of aqueous sodium hydroxide, then	
	add a few drops of **FA 1**, aqueous potassium manganate(VII), then	
	add **FA 2**, sulfuric acid.	

(v) Identify the anion in **FA 6**, and state **one** piece of evidence for your identification.

anion .....

evidence .....

.....

.....

(vi) Give the chemical equation for the reaction between **FA 6** and hydrogen peroxide, H_2O_2, in test **(ii)**. State symbols are **not** required.

.....

[7]

© UCLES 2015

(b) **FA 7**, **FA 8**, **FA 9** and **FA 10** each contain one cation from the list on page 10.
You will attempt to identify the cations by testing with aqueous sodium hydroxide and aqueous ammonia.
In each case, use a 1 cm depth of the solution in a test-tube.

(i) Complete the table below.

test	*observations*			
	FA 7	**FA 8**	**FA 9**	**FA 10**
add sodium hydroxide				
add aqueous ammonia				

(ii) Use your observations to identify, as far as possible, the cation present in each solution. If alternative identities are possible, state this clearly.

FA 7 cation .....

FA 8 cation .....

FA 9 cation .....

FA 10 cation .....

(iii) Give the ionic equation for the reaction of **one** of your cations with a few drops of sodium hydroxide. State symbols are **not** required.

.....

(iv) The precipitates obtained when alkalis are added to solutions of certain cations are sometimes difficult to see. Suggest how, using no additional apparatus, the experiment could be repeated in a way that would make these precipitates more visible.

.....

.....

[9]

[Total: 16]

© UCLES 2015

Qualitative Analysis Notes

Key: [ppt. = precipitate]

1 Reactions of aqueous cations

ion	*reaction with*	
	NaOH(aq)	NH_3(aq)
aluminium, Al^{3+}(aq)	white ppt. soluble in excess	white ppt. insoluble in excess
ammonium, NH_4^+(aq)	no ppt. ammonia produced on heating	–
barium, Ba^{2+}(aq)	no ppt. (if reagents are pure)	no ppt.
calcium, Ca^{2+}(aq)	white ppt. with high [Ca^{2+}(aq)]	no ppt.
chromium(III), Cr^{3+}(aq)	grey-green ppt. soluble in excess giving dark green solution	grey-green ppt. insoluble in excess
copper(II), Cu^{2+}(aq)	pale blue ppt. insoluble in excess	blue ppt. soluble in excess giving dark blue solution
iron(II), Fe^{2+}(aq)	green ppt. turning brown on contact with air insoluble in excess	green ppt. turning brown on contact with air insoluble in excess
iron(III), Fe^{3+}(aq)	red-brown ppt. insoluble in excess	red-brown ppt. insoluble in excess
magnesium, Mg^{2+}(aq)	white ppt. insoluble in excess	white ppt. insoluble in excess
manganese(II), Mn^{2+}(aq)	off-white ppt. rapidly turning brown on contact with air insoluble in excess	off-white ppt. rapidly turning brown on contact with air insoluble in excess
zinc, Zn^{2+}(aq)	white ppt. soluble in excess	white ppt. soluble in excess

© UCLES 2015

2 Reactions of anions

ion	*reaction*
carbonate, CO_3^{2-}	CO_2 liberated by dilute acids
chloride, Cl^-(aq)	gives white ppt. with Ag^+(aq) (soluble in NH_3(aq))
bromide, Br^-(aq)	gives cream ppt. with Ag^+(aq) (partially soluble in NH_3(aq))
iodide, I^-(aq)	gives yellow ppt. with Ag^+(aq) (insoluble in NH_3(aq))
nitrate, NO_3^-(aq)	NH_3 liberated on heating with OH^-(aq) and Al foil
nitrite, NO_2^-(aq)	NH_3 liberated on heating with OH^-(aq) and Al foil; NO liberated by dilute acids (colourless NO → (pale) brown NO_2 in air)
sulfate, SO_4^{2-}(aq)	gives white ppt. with Ba^{2+}(aq) (insoluble in excess dilute strong acids)
sulfite, SO_3^{2-}(aq)	SO_2 liberated with dilute acids; gives white ppt. with Ba^{2+}(aq) (soluble in excess dilute strong acids)

3 Tests for gases

gas	*test and test result*
ammonia, NH_3	turns damp red litmus paper blue
carbon dioxide, CO_2	gives a white ppt. with limewater (ppt. dissolves with excess CO_2)
chlorine, Cl_2	bleaches damp litmus paper
hydrogen, H_2	“pops” with a lighted splint
oxygen, O_2	relights a glowing splint
sulfur dioxide, SO_2	turns acidified aqueous potassium manganate(VII) from purple to colourless

GradeMax

The Periodic Table of the Elements

	Group																	
	I	II									III	IV	V	VI	VII	0
																4.0 **He** Helium 2
						1.0 **H** Hydrogen 1										
											10.8 **B** Boron 5	12.0 **C** Carbon 6	14.0 **N** Nitrogen 7	16.0 **O** Oxygen 8	19.0 **F** Fluorine 9	20.2 **Ne** Neon 10
	6.9 **Li** Lithium 3	9.0 **Be** Beryllium 4									27.0 **Al** Aluminium 13	28.1 **Si** Silicon 14	31.0 **P** Phosphorus 15	32.1 **S** Sulfur 16	35.5 **Cl** Chlorine 17	39.9 **Ar** Argon 18
	23.0 **Na** Sodium 11	24.3 **Mg** Magnesium 12									69.7 **Ga** Gallium 31	72.6 **Ge** Germanium 32	74.9 **As** Arsenic 33	79.0 **Se** Selenium 34	79.9 **Br** Bromine 35	83.8 **Kr** Krypton 36
	39.1 **K** Potassium 19	40.1 **Ca** Calcium 20	45.0 **Sc** Scandium 21	47.9 **Ti** Titanium 22	50.9 **V** Vanadium 23	52.0 **Cr** Chromium 24	54.9 **Mn** Manganese 25	55.8 **Fe** Iron 26	58.9 **Co** Cobalt 27	58.7 **Ni** Nickel 28	63.5 **Cu** Copper 29	65.4 **Zn** Zinc 30	112 **Cd** Cadmium 48	115 **In** Indium 49	119 **Sn** Tin 50	122 **Sb** Antimony 51	128 **Te** Tellurium 52	127 **I** Iodine 53	131 **Xe** Xenon 54
	85.5 **Rb** Rubidium 37	87.6 **Sr** Strontium 38	88.9 **Y** Yttrium 39	91.2 **Zr** Zirconium 40	92.9 **Nb** Niobium 41	95.9 **Mo** Molybdenum 42	**Tc** Technetium 43	101 **Ru** Ruthenium 44	103 **Rh** Rhodium 45	106 **Pd** Palladium 46	108 **Ag** Silver 47	201 **Hg** Mercury 80	204 **Tl** Thallium 81	207 **Pb** Lead 82	209 **Bi** Bismuth 83	**Po** Polonium 84	**At** Astatine 85	**Rn** Radon 86
	133 **Cs** Caesium 55	137 **Ba** Barium 56	139 **La** Lanthanum 57 *	178 **Hf** Hafnium 72	181 **Ta** Tantalum 73	184 **W** Tungsten 74	186 **Re** Rhenium 75	190 **Os** Osmium 76	192 **Ir** Iridium 77	195 **Pt** Platinum 78	197 **Au** Gold 79	201 **Hg** Mercury 80	204 **Tl** Thallium 81	207 **Pb** Lead 82	209 **Bi** Bismuth 83	**Po** Polonium 84	**At** Astatine 85	**Rn** Radon 86
	Fr Francium 87	**Ra** Radium 88	**Ac** Actinium 89 †	**Rf** Rutherfordium 104	**Db** Dubnium 105	**Sg** Seaborgium 106	**Bh** Bohrium 107	**Hs** Hassium 108	**Mt** Meitnerium 109	**Uun** Ununnilium 110	**Uuu** Unununium 111	**Uub** Ununbium 112		**Uuq** Ununquadium 114		**Uuh** Ununhexium 116		**Uuo** Ununoctium 118

*	140 **Ce** Cerium 58	141 **Pr** Praseodymium 59	144 **Nd** Neodymium 60	**Pm** Promethium 61	150 **Sm** Samarium 62	152 **Eu** Europium 63	157 **Gd** Gadolinium 64	159 **Tb** Terbium 65	163 **Dy** Dysprosium 66	165 **Ho** Holmium 67	167 **Er** Erbium 68	169 **Tm** Thulium 69	173 **Yb** Ytterbium 70	175 **Lu** Lutetium 71
†	**Th** Thorium 90	**Pa** Protactinium 91	**U** Uranium 92	**Np** Neptunium 93	**Pu** Plutonium 94	**Am** Americium 95	**Cm** Curium 96	**Bk** Berkelium 97	**Cf** Californium 98	**Es** Einsteinium 99	**Fm** Fermium 100	**Md** Mendelevium 101	**No** Nobelium 102	**Lr** Lawrencium 103

*58-71 Lanthanides
†90-103 Actinides

Key

a	a = relative atomic mass
X	X = atomic symbol
b	b = proton (atomic) number

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination series.

© UCLES 2015