GradeMax

Cambridge International AS & A Level

Cambridge International Examinations
Cambridge International Advanced Subsidiary and Advanced Level

CHEMISTRY **9701/52**

Paper 5 Planning, Analysis and Evaluation **May/June 2016**

MARK SCHEME

Maximum Mark: 30

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the May/June 2016 series for most Cambridge IGCSE®, Cambridge International A and AS Level components and some Cambridge O Level components.

® IGCSE is the registered trademark of Cambridge International Examinations.

© UCLES 2016

Question	Expected answer	Mark
1 (a)	(As the $E^{\ominus}_{cell}$ value increases) ΔH_r decreases or ΔH_r becomes more negative or ΔH_r becomes more exothermic. AND The more reactive the metal then the greater the energy release will be. OR Energy output of both reactions is dependent upon the difference in reactivity (of metals).	[1]
(b)	Independent variable: The (type of) metal Dependent variable: temperature change or rise or increase OR enthalpy change	[1] [1]
(c) (i)	Diagram should indicate a labelled insulated container AND a labelled thermometer in the liquid.	[1]
(ii)	Mass of metal before and after Initial temperature (before metal added) AND Highest temperature (after metal added)	[1] [1]
(iii)	Wear gloves	[1]
(iv)	Moles $CuSO_4$ = 0.025 mol, therefore moles of magnesium = 0.025 mol (minimum) mass Mg > (0.025 × 24.3 =) 0.6075 g AND mass required value is greater than 0.6075 g	[1] [1]
(v)	Larger surface area AND causes increased rate of reaction	[1]
(vi)	Ensure uniformity of heating (of solution)	[1]

© Cambridge International Examinations 2016

Question	Expected answer	Mark
(d)	$50.0 \times 4.18 \times 58.5 = 12\,226.5$ (J) $\Delta H_r = 12\,226.5 / 0.025 = \frac{489\,000}{1000} = -489$	[1] [1]
(e)	Complete circuit involving labelled voltmeter; labelled salt bridge; two separate solutions; (Solutions are) magnesium sulfate or $MgSO_4$ with magnesium or Mg rod and copper(II) sulfate $CuSO_4$ with copper or Cu rod Concentration of solution(s) is 1 mol dm^{-3} or 1 M	[1] [1] [1]
(f)	So that values can be compared	[1]
(g)	Both ΔH_r (Zn) and ΔH_r (Fe) values which are consistent with the prediction in **(a)**.	[1]
		[18]

© Cambridge International Examinations 2016

Question	Expected answer	Mark
2 (a)		

Mass of liquid Y used / g	Volume of vapour Y / cm^3
0.15	48
0.10	35
0.21	72
0.17	58
0.24	83
0.09	31
0.20	70
0.23	79
0.12	41
0.22	73

All mass values.	[1]
All volume values.	[1]
(b) Candidate's points plotted correctly from table in 2(a).	[1]
Line of best fit drawn.	[1]

© Cambridge International Examinations 2016

Question	Expected answer	Mark
(c) (i)	Y evaporates from the (hypodermic) syringe OR Y evaporates before injection OR Y evaporates before weighing / after injection	[1]
(ii)	(Stop evaporation by) Keeping the syringe as cool as possible OR Closing off the needle end to stop evaporation OR Minimising length of time between each weighing	[1]
(d) (i)	correct co-ordinates. correct calculation of the gradient must be **three** significant figures	[1] [1]
(ii)	Calculation of $M_r = 3.07 \times 10^4$ / gradient in 2(d)(i) Answer	[1] [1]
(e)	M_r (from mass spectrum) = 84 OR empirical formula = CH_2 OR ratio of C and H seen as 1:2 Y is C_6H_{12}	[1] [1]
		[12]

© Cambridge International Examinations 2016