

Cambridge  
International  
AS & A Level

**Cambridge Assessment International Education**  
Cambridge International Advanced Subsidiary and Advanced Level

CANDIDATE  
NAME

|  |
|--|
|  |
|--|

CENTRE  
NUMBER

|  |  |  |  |  |
|--|--|--|--|--|
|  |  |  |  |  |
|--|--|--|--|--|

CANDIDATE  
NUMBER

|  |  |  |  |
|--|--|--|--|
|  |  |  |  |
|--|--|--|--|

\* 4 6 2 9 2 7 9 9 2 0 \*



**CHEMISTRY**

**9701/32**

Paper 3 Advanced Practical Skills 2

**May/June 2019**

**2 hours**

Candidates answer on the Question Paper.

Additional Materials: As listed in the Confidential Instructions

**READ THESE INSTRUCTIONS FIRST**

Write your centre number, candidate number and name on all the work you hand in.  
Give details of the practical session and laboratory where appropriate, in the boxes provided.  
Write in dark blue or black pen.  
You may use an HB pencil for any diagrams or graphs.  
Do not use staples, paper clips, glue or correction fluid.  
**DO NOT WRITE IN ANY BARCODES.**

Answer **all** questions.  
Electronic calculators may be used.  
You may lose marks if you do not show your working or if you do not use appropriate units.  
Use of a Data Booklet is unnecessary.

Qualitative Analysis Notes are printed on pages 10 and 11.  
A copy of the Periodic Table is printed on page 12.

At the end of the examination, fasten all your work securely together.  
The number of marks is given in brackets [ ] at the end of each question or part question.

| Session    |
|------------|
|            |
| Laboratory |
|            |

| For Examiner's Use |  |
|--------------------|--|
| 1                  |  |
| 2                  |  |
| 3                  |  |
| <b>Total</b>       |  |

This document consists of **12** printed pages.

## Quantitative Analysis

Read through the whole method before starting any practical work. Where appropriate, prepare a table for your results in the space provided.

Show your working and appropriate significant figures in the final answer to **each** step of your calculations.

- 1 Amidosulfonic acid is a monoprotic acid with the formula  $\text{HSO}_x\text{NH}_2$ . In this experiment you will determine the value of  $x$  in the formula by titration with an alkali.

**FB 1** is amidosulfonic acid,  $\text{HSO}_x\text{NH}_2$ . You are supplied with approximately 3.0 g.

**FB 2** is  $0.110 \text{ mol dm}^{-3}$  sodium hydroxide, NaOH.

thymolphthalein indicator

### (a) Method

#### Preparing a solution of amidosulfonic acid

- Weigh the empty beaker. Record the mass.
- Add 2.50–2.70 g of **FB 1** to the beaker. Weigh the beaker and its contents. Record the mass.
- Calculate and record the mass of **FB 1** used.
- Add approximately  $100 \text{ cm}^3$  of distilled water to the beaker and stir to dissolve **FB 1**.
- Transfer the solution to the  $250 \text{ cm}^3$  volumetric flask.
- Rinse the beaker twice with approximately  $20 \text{ cm}^3$  of distilled water each time and add to the volumetric flask.
- Add distilled water to the volumetric flask to make  $250 \text{ cm}^3$  of solution.
- Shake the volumetric flask to mix the solution thoroughly and label it **FB 3**.

#### Titration of FB 3

- Fill the burette with **FB 2**.
- Pipette  $25.0 \text{ cm}^3$  of **FB 3** into a conical flask.
- Add a few drops of thymolphthalein to the conical flask.
- Carry out a **rough** titration.
- Record your burette readings and the rough titre in the space below.

The rough titre is .....  $\text{cm}^3$ .

3

- Carry out as many accurate titrations as you think necessary to obtain consistent results.
- Make sure any recorded results show the precision of your practical work.
- Record all of your burette readings and the volume of **FB 2** added in each accurate titration.

|      |  |
|------|--|
| I    |  |
| II   |  |
| III  |  |
| IV   |  |
| V    |  |
| VI   |  |
| VII  |  |
| VIII |  |

[8]

- (b) From your accurate titration results, obtain a suitable value for the volume of **FB 2** to be used in your calculations.  
Show clearly how you obtained this value.

25.0 cm<sup>3</sup> of **FB 3** required ..... cm<sup>3</sup> of **FB 2**. [1]

(c) **Calculations**

- (i) Give your answers to (ii), (iii), (iv) and (v) to the appropriate number of significant figures. [1]

- (ii) Calculate the number of moles of sodium hydroxide, NaOH, in the volume of **FB 2** calculated in (b).

moles of NaOH = ..... mol [1]

- (iii) One mole of sodium hydroxide neutralises one mole of amidosulfonic acid.

Complete the equation for the reaction of amidosulfonic acid with sodium hydroxide.  
Include state symbols.



State the number of moles of amidosulfonic acid that reacted with the number of moles of NaOH calculated in (ii).

moles of HSO<sub>x</sub>NH<sub>2</sub> = ..... mol [1]

4

- (iv) Use your results on page 2 and your answer to (iii) to calculate the relative formula mass,  $M_r$ , of amidosulfonic acid.

$M_r$  of amidosulfonic acid = ..... [1]

- (v) Calculate the value of  $x$  in the formula of amidosulfonic acid,  $\text{HSO}_x\text{NH}_2$ .

$x$  = ..... [1]

- (d) The salts produced from amidosulfonic acid are called amidosulfonates.

You are to carry out an experiment using **FB 1** to find out if barium amidosulfonate is soluble in water.

- Describe your experiment.
- Record your observation(s) **and** conclusion.

.....

.....

.....

.....

..... [2]

[Total: 16]

**Question 2 starts on the next page.**

## 6

- 2 When hydrated copper hydroxycarbonate,  $\text{CuCO}_3 \cdot \text{Cu}(\text{OH})_2 \cdot y\text{H}_2\text{O}$  is heated, it decomposes as shown.



In this experiment, you will heat hydrated copper hydroxycarbonate to decompose it. You will use your results to investigate the value of  $y$ .

**FB 4** is hydrated copper hydroxycarbonate,  $\text{CuCO}_3 \cdot \text{Cu}(\text{OH})_2 \cdot y\text{H}_2\text{O}$ .

**(a) Method**

- Weigh the crucible with its lid and record the mass.
- Add all the **FB 4** from the container into the crucible.
- Weigh the crucible and lid with **FB 4** and record the mass.
- Place the crucible and contents on the pipe-clay triangle.
- Heat the crucible and contents gently for approximately two minutes with the lid on.
- Use tongs to remove the lid and heat strongly for approximately three minutes.
- Replace the lid and leave the crucible and residue to cool for several minutes.

**While the crucible is cooling, you may wish to begin work on Question 3.**

- When the crucible is cool, reweigh it with its lid and contents. Record the mass.
- Calculate and record the mass of **FB 4** and the mass of residue obtained.

**Results**

|     |  |
|-----|--|
| I   |  |
| II  |  |
| III |  |
| IV  |  |

State the observations made while the decomposition of **FB 4** was taking place.

.....

.....

[4]

**(b) Calculations**

- (i) Calculate the number of moles of copper oxide, CuO, obtained as residue.

moles of CuO obtained = ..... mol [1]

- (ii) Use your results to calculate the relative formula mass,  $M_r$ , of hydrated copper hydroxycarbonate,  $\text{CuCO}_3 \cdot \text{Cu}(\text{OH})_2 \cdot y\text{H}_2\text{O}$ .

$M_r$  of  $\text{CuCO}_3 \cdot \text{Cu}(\text{OH})_2 \cdot y\text{H}_2\text{O}$  = ..... [2]

- (iii) Use the Periodic Table to calculate the relative formula mass of  $\text{CuCO}_3 \cdot \text{Cu}(\text{OH})_2$ .

$M_r$  of  $\text{CuCO}_3 \cdot \text{Cu}(\text{OH})_2$  = .....

Use your answer to (ii) and this  $M_r$  to comment on the value of  $y$  in the formula of hydrated copper hydroxycarbonate.

.....  
..... [2]

- (c) State **one** way to improve the accuracy of the experiment, using the same mass of **FB 4**.

.....  
..... [1]

[Total: 10]

### Qualitative Analysis

Where reagents are selected for use in a test, the **name** or **correct formula** of the element or compound must be given.

At each stage of any test you are to record details of the following:

- colour changes seen;
- the formation of any precipitate and its solubility in an excess of the reagent added;
- the formation of any gas and its identification by a suitable test.

You should indicate clearly at what stage in a test a change occurs.

If any solution is warmed, a **boiling tube** must be used.

Rinse and reuse test-tubes and boiling tubes where possible.

**No additional tests for ions present should be attempted.**

**3 (a) FB 5** contains one cation and one anion from those listed in the Qualitative Analysis Notes.

- (i) Transfer a **small** spatula measure of **FB 5** into a hard-glass test-tube. Heat gently at first and then heat strongly, until no further change occurs.

Record **all** your observations.

.....

.....

.....

.....

.....

..... [3]

- (ii) To a 1 cm depth of dilute hydrochloric acid in a test-tube, add a **small** spatula measure of **FB 5**.

Record your observations.

.....

.....

.....

..... [2]

- (iii) Deduce the formula of **FB 5**.

..... [1]



(b) **FB 6** is a solution of a compound containing one cation from those listed in the Qualitative Analysis Notes.

(i) Carry out the following tests using a 1 cm depth of **FB 6** in a test-tube for each test.

| <i>test</i>                                            | <i>observations</i> |
|--------------------------------------------------------|---------------------|
| Add two drops of Universal Indicator solution.         |                     |
| Add an equal volume of aqueous manganese(II) chloride. |                     |
| Add an equal volume of dilute sulfuric acid.           |                     |
| Add a few drops of aqueous silver nitrate, then        |                     |
| add aqueous ammonia.                                   |                     |
| Add a few drops of aqueous copper(II) sulfate, then    |                     |
| add excess dilute hydrochloric acid.                   |                     |

[4]

(ii) What can be determined about **FB 6** from its reaction with manganese(II) chloride?

..... [1]

(iii) Give the ionic equation for the reaction between **FB 6** and sulfuric acid. Include state symbols.

..... [1]

(iv) When aqueous copper(II) sulfate was added to **FB 6**, **two** precipitates were formed.

Deduce the identity of the two precipitates.

..... and .....

Explain the observation you made when you added excess dilute hydrochloric acid in the final test of (b)(i).

.....

.....

.....

[2]

[Total: 14]

## Qualitative Analysis Notes

## 1 Reactions of aqueous cations

| <i>ion</i>                                     | <i>reaction with</i>                                                               |                                                                                    |
|------------------------------------------------|------------------------------------------------------------------------------------|------------------------------------------------------------------------------------|
|                                                | NaOH(aq)                                                                           | NH <sub>3</sub> (aq)                                                               |
| aluminium,<br>Al <sup>3+</sup> (aq)            | white ppt.<br>soluble in excess                                                    | white ppt.<br>insoluble in excess                                                  |
| ammonium,<br>NH <sub>4</sub> <sup>+</sup> (aq) | no ppt.<br>ammonia produced on heating                                             | —                                                                                  |
| barium,<br>Ba <sup>2+</sup> (aq)               | faint white ppt. is nearly always<br>observed unless reagents are pure             | no ppt.                                                                            |
| calcium,<br>Ca <sup>2+</sup> (aq)              | white ppt. with high [Ca <sup>2+</sup> (aq)]                                       | no ppt.                                                                            |
| chromium(III),<br>Cr <sup>3+</sup> (aq)        | grey-green ppt.<br>soluble in excess                                               | grey-green ppt.<br>insoluble in excess                                             |
| copper(II),<br>Cu <sup>2+</sup> (aq)           | pale blue ppt.<br>insoluble in excess                                              | blue ppt. soluble in excess<br>giving dark blue solution                           |
| iron(II),<br>Fe <sup>2+</sup> (aq)             | green ppt. turning brown on contact<br>with air<br>insoluble in excess             | green ppt. turning brown on contact<br>with air<br>insoluble in excess             |
| iron(III),<br>Fe <sup>3+</sup> (aq)            | red-brown ppt.<br>insoluble in excess                                              | red-brown ppt.<br>insoluble in excess                                              |
| magnesium,<br>Mg <sup>2+</sup> (aq)            | white ppt.<br>insoluble in excess                                                  | white ppt.<br>insoluble in excess                                                  |
| manganese(II),<br>Mn <sup>2+</sup> (aq)        | off-white ppt. rapidly turning brown<br>on contact with air<br>insoluble in excess | off-white ppt. rapidly turning brown<br>on contact with air<br>insoluble in excess |
| zinc,<br>Zn <sup>2+</sup> (aq)                 | white ppt.<br>soluble in excess                                                    | white ppt.<br>soluble in excess                                                    |

## 2 Reactions of anions

| <i>ion</i>                                | <i>reaction</i>                                                                                 |
|-------------------------------------------|-------------------------------------------------------------------------------------------------|
| carbonate,<br>$\text{CO}_3^{2-}$          | $\text{CO}_2$ liberated by dilute acids                                                         |
| chloride,<br>$\text{Cl}^-(\text{aq})$     | gives white ppt. with $\text{Ag}^+(\text{aq})$ (soluble in $\text{NH}_3(\text{aq})$ )           |
| bromide,<br>$\text{Br}^-(\text{aq})$      | gives cream ppt. with $\text{Ag}^+(\text{aq})$ (partially soluble in $\text{NH}_3(\text{aq})$ ) |
| iodide,<br>$\text{I}^-(\text{aq})$        | gives yellow ppt. with $\text{Ag}^+(\text{aq})$ (insoluble in $\text{NH}_3(\text{aq})$ )        |
| nitrate,<br>$\text{NO}_3^-(\text{aq})$    | $\text{NH}_3$ liberated on heating with $\text{OH}^-(\text{aq})$ and $\text{Al}$ foil           |
| nitrite,<br>$\text{NO}_2^-(\text{aq})$    | $\text{NH}_3$ liberated on heating with $\text{OH}^-(\text{aq})$ and $\text{Al}$ foil           |
| sulfate,<br>$\text{SO}_4^{2-}(\text{aq})$ | gives white ppt. with $\text{Ba}^{2+}(\text{aq})$ (insoluble in excess dilute strong acids)     |
| sulfite,<br>$\text{SO}_3^{2-}(\text{aq})$ | gives white ppt. with $\text{Ba}^{2+}(\text{aq})$ (soluble in excess dilute strong acids)       |

## 3 Tests for gases

| <i>gas</i>                    | <i>test and test result</i>                                                   |
|-------------------------------|-------------------------------------------------------------------------------|
| ammonia, $\text{NH}_3$        | turns damp red litmus paper blue                                              |
| carbon dioxide, $\text{CO}_2$ | gives a white ppt. with limewater (ppt. dissolves with excess $\text{CO}_2$ ) |
| chlorine, $\text{Cl}_2$       | bleaches damp litmus paper                                                    |
| hydrogen, $\text{H}_2$        | 'pops' with a lighted splint                                                  |
| oxygen, $\text{O}_2$          | relights a glowing splint                                                     |

## The Periodic Table of Elements

| Group                               |                                      |                                     |                                      |                                                                              |                                       |                                      |                                       |                                     |                                       |                                       |                                      |                                      |                                      |                                      |                                       |                                    |                                      |                                      |    |    |  |  |
|-------------------------------------|--------------------------------------|-------------------------------------|--------------------------------------|------------------------------------------------------------------------------|---------------------------------------|--------------------------------------|---------------------------------------|-------------------------------------|---------------------------------------|---------------------------------------|--------------------------------------|--------------------------------------|--------------------------------------|--------------------------------------|---------------------------------------|------------------------------------|--------------------------------------|--------------------------------------|----|----|--|--|
| 1                                   | 2                                    |                                     | Group                                |                                                                              |                                       |                                      |                                       |                                     |                                       |                                       |                                      |                                      |                                      |                                      | 13                                    | 14                                 | 15                                   | 16                                   | 17 | 18 |  |  |
| 3<br><b>Li</b><br>lithium<br>6.9    | 4<br><b>Be</b><br>beryllium<br>9.0   | 11<br><b>Na</b><br>sodium<br>23.0   | 12<br><b>Mg</b><br>magnesium<br>24.3 | <b>Key</b><br>atomic number<br>atomic symbol<br>name<br>relative atomic mass |                                       |                                      |                                       |                                     |                                       |                                       |                                      |                                      |                                      |                                      |                                       | 1<br><b>H</b><br>hydrogen<br>1.0   |                                      |                                      |    |    |  |  |
|                                     |                                      |                                     |                                      |                                                                              |                                       |                                      |                                       |                                     |                                       |                                       |                                      |                                      |                                      |                                      |                                       |                                    |                                      |                                      |    |    |  |  |
|                                     |                                      |                                     |                                      |                                                                              |                                       |                                      |                                       |                                     |                                       |                                       |                                      |                                      |                                      |                                      |                                       |                                    |                                      |                                      |    |    |  |  |
| 19<br><b>K</b><br>potassium<br>39.1 | 20<br><b>Ca</b><br>calcium<br>40.1   | 21<br><b>Sc</b><br>scandium<br>45.0 | 22<br><b>Ti</b><br>titanium<br>47.9  | 23<br><b>V</b><br>vanadium<br>50.9                                           | 24<br><b>Cr</b><br>chromium<br>52.0   | 25<br><b>Mn</b><br>manganese<br>54.9 | 26<br><b>Fe</b><br>iron<br>55.8       | 27<br><b>Co</b><br>cobalt<br>58.9   | 28<br><b>Ni</b><br>nickel<br>58.7     | 29<br><b>Cu</b><br>copper<br>63.5     | 30<br><b>Zn</b><br>zinc<br>65.4      | 31<br><b>Ga</b><br>gallium<br>69.7   | 32<br><b>Ge</b><br>germanium<br>72.6 | 33<br><b>As</b><br>arsenic<br>74.9   | 34<br><b>Se</b><br>selenium<br>79.0   | 35<br><b>Br</b><br>bromine<br>79.9 | 36<br><b>Kr</b><br>krypton<br>83.8   |                                      |    |    |  |  |
| 37<br><b>Rb</b><br>rubidium<br>85.5 | 38<br><b>Sr</b><br>strontium<br>87.6 | 39<br><b>Y</b><br>yttrium<br>88.9   | 40<br><b>Zr</b><br>zirconium<br>91.2 | 41<br><b>Nb</b><br>niobium<br>92.9                                           | 42<br><b>Mo</b><br>molybdenum<br>95.9 | 43<br><b>Tc</b><br>technetium<br>—   | 44<br><b>Ru</b><br>ruthenium<br>101.1 | 45<br><b>Rh</b><br>rhodium<br>102.9 | 46<br><b>Pd</b><br>palladium<br>106.4 | 47<br><b>Ag</b><br>silver<br>107.9    | 48<br><b>Cd</b><br>cadmium<br>112.4  | 49<br><b>In</b><br>indium<br>114.8   | 50<br><b>Sn</b><br>tin<br>118.7      | 51<br><b>Sb</b><br>antimony<br>121.8 | 52<br><b>Te</b><br>tellurium<br>127.6 | 53<br><b>I</b><br>iodine<br>126.9  | 54<br><b>Xe</b><br>xenon<br>131.3    |                                      |    |    |  |  |
| 55<br><b>Cs</b><br>caesium<br>132.9 | 56<br><b>Ba</b><br>barium<br>137.3   | 57–71<br>lanthanoids                |                                      | 72<br><b>Hf</b><br>hafnium<br>178.5                                          | 73<br><b>Ta</b><br>tantalum<br>180.9  | 74<br><b>W</b><br>tungsten<br>183.8  | 75<br><b>Re</b><br>rhenium<br>186.2   | 76<br><b>Os</b><br>osmium<br>190.2  | 77<br><b>Ir</b><br>iridium<br>192.2   | 78<br><b>Pt</b><br>platinum<br>195.1  | 79<br><b>Au</b><br>gold<br>197.0     | 80<br><b>Hg</b><br>mercury<br>200.6  | 81<br><b>Tl</b><br>thallium<br>204.4 | 82<br><b>Pb</b><br>lead<br>207.2     | 83<br><b>Bi</b><br>bismuth<br>209.0   | 84<br><b>Po</b><br>polonium<br>—   | 85<br><b>At</b><br>astatine<br>—     | 86<br><b>Rn</b><br>radon<br>—        |    |    |  |  |
| 87<br><b>Fr</b><br>francium<br>—    | 88<br><b>Ra</b><br>radium<br>—       | 89–103<br>actinoids                 |                                      | 104<br><b>Rf</b><br>rutherfordium<br>—                                       | 105<br><b>Db</b><br>dubnium<br>—      | 106<br><b>Sg</b><br>seaborgium<br>—  | 107<br><b>Bh</b><br>bohrium<br>—      | 108<br><b>Hs</b><br>hassium<br>—    | 109<br><b>Mt</b><br>meitnerium<br>—   | 110<br><b>Ds</b><br>darmstadtium<br>— | 111<br><b>Rg</b><br>roentgenium<br>— | 112<br><b>Cn</b><br>copernicium<br>— | 114<br><b>Fl</b><br>flerovium<br>—   | 116<br><b>Lv</b><br>livermorium<br>— | 117<br><b>Ts</b><br>tennessine<br>—   | 118<br><b>Og</b><br>oganeson<br>—  | 119<br><b>Uue</b><br>ununennium<br>— | 120<br><b>Uuh</b><br>ununhexium<br>— |    |    |  |  |

|    |    |           |       |    |    |         |       |    |    |              |       |    |    |           |       |    |    |            |       |    |    |           |       |    |    |           |       |    |    |            |       |    |    |           |       |    |    |             |       |    |    |             |       |     |    |         |       |     |    |              |       |     |    |           |       |     |    |            |       |
|----|----|-----------|-------|----|----|---------|-------|----|----|--------------|-------|----|----|-----------|-------|----|----|------------|-------|----|----|-----------|-------|----|----|-----------|-------|----|----|------------|-------|----|----|-----------|-------|----|----|-------------|-------|----|----|-------------|-------|-----|----|---------|-------|-----|----|--------------|-------|-----|----|-----------|-------|-----|----|------------|-------|
| 57 | La | lanthanum | 138.9 | 58 | Ce | cerium  | 140.1 | 59 | Pr | praseodymium | 140.9 | 60 | Nd | neodymium | 144.4 | 61 | Pm | promethium | —     | 62 | Sm | samarium  | 150.4 | 63 | Eu | euroium   | 152.0 | 64 | Gd | gadolinium | 157.3 | 65 | Tb | terbium   | 158.9 | 66 | Dy | dysprosium  | 162.5 | 67 | Ho | holmium     | 164.9 | 68  | Er | erbium  | 167.3 | 69  | Tm | thulium      | 168.9 | 70  | Yb | ytterbium | 173.1 | 71  | Lu | lutetium   | 175.0 |
| 89 | Ac | actinium  | 227.0 | 90 | Th | thorium | 232.0 | 91 | Pa | protactinium | 231.0 | 92 | U  | uranium   | 238.0 | 93 | Np | neptunium  | 237.0 | 94 | Pu | plutonium | 244.0 | 95 | Am | americium | 243.0 | 96 | Cm | curium     | 247.0 | 97 | Bk | berkelium | 247.0 | 98 | Cf | californium | 251.0 | 99 | Es | einsteinium | 252.0 | 100 | Fm | fermium | 257.0 | 101 | Md | mendeleevium | 258.0 | 102 | No | nobelium  | 259.0 | 103 | Lr | lawrencium | 262.0 |

© UCLES 2019