



# Cambridge International AS & A Level

CANDIDATE  
NAME
CENTRE  
NUMBER

|  |  |  |  |  |
|--|--|--|--|--|
|  |  |  |  |  |
|--|--|--|--|--|

CANDIDATE  
NUMBER

|  |  |  |  |
|--|--|--|--|
|  |  |  |  |
|--|--|--|--|

## CHEMISTRY

9701/53

Paper 5 Planning, Analysis and Evaluation

May/June 2022

1 hour 15 minutes

You must answer on the question paper.

No additional materials are needed.

## INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- You may use a calculator.
- You should show all your working and use appropriate units.

## INFORMATION

- The total mark for this paper is 30.
- The number of marks for each question or part question is shown in brackets [ ].
- The Periodic Table is printed in the question paper.
- Important values, constants and standards are printed in the question paper.

This document has **16** pages. Any blank pages are indicated.



## 2

- 1 A student plans an investigation to find the molar ratio of the reaction between sodium chloride,  $\text{NaCl}$ , and a lead compound.

The student is provided with solid  $\text{NaCl}$  and  $0.200 \text{ mol dm}^{-3}$  aqueous lead compound.

The reaction between  $\text{NaCl}(\text{aq})$  and the aqueous lead compound produces an insoluble compound as a precipitate.

- (a) The student prepares  $0.200 \text{ mol dm}^{-3}$   $\text{NaCl}(\text{aq})$ .

Calculate the mass of  $\text{NaCl}(\text{s})$  needed to make  $250.0 \text{ cm}^3$  of  $0.200 \text{ mol dm}^{-3}$   $\text{NaCl}(\text{aq})$ .

mass of  $\text{NaCl} = \dots\dots\dots \text{ g}$  [1]

- (b) The student weighs the mass of  $\text{NaCl}(\text{s})$  calculated in (a) in a weighing boat. The solid mass is then transferred into a small beaker.

Describe how the student should accurately weigh by difference so the exact mass of  $\text{NaCl}$  transferred into the small beaker is known.

.....  
.....  
.....  
..... [1]

- (c) The student is given a small beaker containing the mass of  $\text{NaCl}$  calculated in (a).

Describe how the student should prepare  $250.0 \text{ cm}^3$  of  $0.200 \text{ mol dm}^{-3}$   $\text{NaCl}(\text{aq})$ .

Include the names and capacities of each piece of apparatus used in the preparation of the solution.

.....  
.....  
.....  
.....  
..... [2]

3

- (d) The student plans the following method using the  $0.200 \text{ mol dm}^{-3}$  aqueous lead compound and the  $0.200 \text{ mol dm}^{-3}$   $\text{NaCl(aq)}$  prepared in (c).

**Step 1** Mix the  $\text{NaCl(aq)}$  and the aqueous lead compound in eight separate beakers in the proportions by volume shown in Table 1.1.

Table 1.1

| beaker | volume of $0.200 \text{ mol dm}^{-3}$ $\text{NaCl(aq)}$ / $\text{cm}^3$ | volume of $0.200 \text{ mol dm}^{-3}$ aqueous lead compound / $\text{cm}^3$ |
|--------|-------------------------------------------------------------------------|-----------------------------------------------------------------------------|
| 1      | 10.00                                                                   | 40.00                                                                       |
| 2      | 15.00                                                                   | 35.00                                                                       |
| 3      | 20.00                                                                   | 30.00                                                                       |
| 4      | 25.00                                                                   | 25.00                                                                       |
| 5      | 30.00                                                                   | 20.00                                                                       |
| 6      | 35.00                                                                   | 15.00                                                                       |
| 7      | 40.00                                                                   | 10.00                                                                       |
| 8      | 45.00                                                                   | 5.00                                                                        |

**Step 2** Filter the contents of each beaker to collect the precipitate.

**Step 3** Dry the precipitate for 3 minutes in an oven and allow to cool.

**Step 4** Weigh and record the mass of precipitate produced in each beaker.

- (i) State **one** extra step that would improve this method. Explain why this step is necessary.

extra step:

.....

.....

explanation:

.....

.....

.....

[2]

4

- (ii) The volumes of solutions are measured using a burette.

Calculate the percentage error when measuring  $10.00\text{ cm}^3$  of solution.  
Show your working.

percentage error = ..... [1]

- (iii) Explain how you would ensure that the results of the investigation are reliable.

.....  
.....  
..... [1]

- (e) The results of the investigation are shown on the graph in Fig. 1.1.

- (i) Draw **two** straight lines of best fit through the points. Extrapolate both lines so they intersect. [1]
- (ii) Using Fig. 1.1 and Table 1.1, state the volumes of  $0.200\text{ mol dm}^{-3}$   $\text{NaCl(aq)}$  and  $0.200\text{ mol dm}^{-3}$  aqueous lead compound which produce the maximum mass of precipitate.

Calculate the molar ratio in which the  $\text{NaCl}$  and the lead compound react.

volume of  $0.200\text{ mol dm}^{-3}$   $\text{NaCl(aq)}$  = .....  $\text{cm}^3$

volume of  $0.200\text{ mol dm}^{-3}$  aqueous lead compound = .....  $\text{cm}^3$

molar ratio of  $\text{NaCl}$ : lead compound = ..... : ..... [2]

- (f) Use the molar ratio in (e)(ii) to deduce the formula of the precipitate.

formula = ..... [1]

5




Fig. 1.1

6

(g) A student suggests that a simpler method can be used to find the molar ratio.

Different volumes of  $0.200 \text{ mol dm}^{-3} \text{ NaCl(aq)}$  and  $0.200 \text{ mol dm}^{-3}$  aqueous lead compound are mixed in test-tubes. The resulting precipitates are allowed to settle. The height of each precipitate is then measured.


A further two investigations are carried out. The volumes used and the results of the two investigations are shown.

### Investigation 1

Precipitate heights are measured after 1 minute.

**Table 1.2**

|                                                                         |   |   |   |   |   |   |   |   |   |
|-------------------------------------------------------------------------|---|---|---|---|---|---|---|---|---|
| volume $0.200 \text{ mol dm}^{-3} \text{ NaCl(aq)}/\text{cm}^3$         | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 |
| volume $0.200 \text{ mol dm}^{-3}$ aqueous lead compound/ $\text{cm}^3$ | 9 | 8 | 7 | 6 | 5 | 4 | 3 | 2 | 1 |




**Fig. 1.2**

**Investigation 2**

Precipitate heights are measured after 5 minutes.

**Table 1.3**

|                                                                            |   |   |   |   |   |   |   |   |   |
|----------------------------------------------------------------------------|---|---|---|---|---|---|---|---|---|
| volume $0.200 \text{ mol dm}^{-3}$<br>$\text{NaCl(aq)}/\text{cm}^3$        | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 |
| volume $0.200 \text{ mol dm}^{-3}$<br>aqueous lead compound/ $\text{cm}^3$ | 5 | 5 | 5 | 5 | 5 | 5 | 5 | 5 | 5 |

**Fig. 1.3**

Neither investigation produced the expected results. Both investigations, 1 and 2, contain weaknesses in the experimental procedure.

State how you would modify the experimental procedure in each case so that the expected results are obtained.

modification for investigation 1:

.....

.....

modification for investigation 2:


.....

.....

[2]

[Total: 14]

- 2 A student plans to study the rate of hydrolysis of 2-bromo-2-methylpropane.



As the concentration of 2-bromo-2-methylpropane decreases during the reaction, the concentration of hydrogen ions increases.

The student plans the following method.

- Step 1** Place 100 cm<sup>3</sup> of a mixture of propanone and water into a conical flask.
- Step 2** Heat the mixture to 35 °C and maintain this temperature.
- Step 3** Add 1.00 cm<sup>3</sup> of 2-bromo-2-methylpropane to the mixture and start timing.
- Step 4** After 1 minute, transfer a 10.00 cm<sup>3</sup> sample of the reaction mixture into a conical flask containing ice and 4 drops of methyl orange indicator.
- Step 5** Immediately titrate the 10.00 cm<sup>3</sup> of the reaction mixture with 0.0200 mol dm<sup>-3</sup> sodium hydroxide.
- Step 6** Repeat sampling and titrating at regular time intervals over a total time of 45 minutes.
- Step 7** Heat the reaction mixture to 50 °C, remove the final sample, and titrate this.
- (a) (i) State the apparatus you would use to maintain the temperature of the reaction mixture.

..... [1]

- (ii) Suggest why the experiment is carried out away from naked flames.

.....  
 ..... [1]

- (b) State the pieces of equipment and their capacities that you would use to:

- (i) measure 1.00 cm<sup>3</sup> of 2-bromo-2-methylpropane in step 3

..... [1]

- (ii) transfer a 10.00 cm<sup>3</sup> sample of the mixture in step 4.

..... [1]

**Question 2 continues on the next page.**

10

(c) Explain why the reaction mixture is transferred into a conical flask containing ice.

.....  
 ..... [1]

(d) State the measured dependent variable for this experiment.

..... [1]

(e) (i) The student recorded the results.  $V_{\text{final}}$  is the final titre volume,  $47.25 \text{ cm}^3$  in step 7.

Complete Table 2.1 by calculating the value of  $V_{\text{final}} - V_{\text{t}}$ .

Record the values to **2 decimal places**.

**Table 2.1**

| time/s | titre, $V_{\text{t}}/\text{cm}^3$ | $V_{\text{final}} - V_{\text{t}}/\text{cm}^3$ |
|--------|-----------------------------------|-----------------------------------------------|
| 60     | 1.25                              |                                               |
| 300    | 7.75                              |                                               |
| 600    | 17.75                             |                                               |
| 900    | 20.00                             |                                               |
| 1200   | 24.25                             |                                               |
| 1500   | 28.40                             |                                               |
| 1800   | 31.15                             |                                               |
| 2700   | 38.00                             |                                               |
| final  | 47.25                             |                                               |

[1]

(ii) The titre,  $V_{\text{t}}$ , is proportional to the concentration of the hydrogen ions.


State what  $V_{\text{final}} - V_{\text{t}}$  is proportional to.

..... [1]

(iii) Plot a graph on the grid to show the relationship between  $V_{\text{final}} - V_{\text{t}}$  and time.

Use a cross (×) to plot each data point. Draw a curved line of best fit. [2]

11



(iv) Circle the point on the graph you consider to be most anomalous.

Suggest **one** reason why this anomaly may have occurred during this experimental procedure.

.....  
..... [2]

12

- (v) Use the graph to find two half-lives,  $t_{\frac{1}{2}}$ , for this reaction.

State the coordinates of both points you used in your calculations.

first  $t_{\frac{1}{2}}$ : coordinates ..... and .....

half-life = ..... s

second  $t_{\frac{1}{2}}$ : coordinates ..... and .....

half-life = ..... s

[3]

- (vi) Use your answer to (e)(v) to state the order of the reaction with respect to 2-bromo-2-methylpropane. Explain your answer.

(If you were unable to obtain an answer to (e)(v) you may use the values 1050 s and 1045 s for the half-lives. These are **not** the correct values.)

order = .....

explanation .....

[1]

[Total: 16]





**Important values, constants and standards**

|                                 |                                                                                                                                           |
|---------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------|
| molar gas constant              | $R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$                                                                                              |
| Faraday constant                | $F = 9.65 \times 10^4 \text{ C mol}^{-1}$                                                                                                 |
| Avogadro constant               | $L = 6.022 \times 10^{23} \text{ mol}^{-1}$                                                                                               |
| electronic charge               | $e = -1.60 \times 10^{-19} \text{ C}$                                                                                                     |
| molar volume of gas             | $V_m = 22.4 \text{ dm}^3 \text{ mol}^{-1}$ at s.t.p. (101 kPa and 273 K)<br>$V_m = 24.0 \text{ dm}^3 \text{ mol}^{-1}$ at room conditions |
| ionic product of water          | $K_w = 1.00 \times 10^{-14} \text{ mol}^2 \text{ dm}^{-6}$ (at 298 K (25 °C))                                                             |
| specific heat capacity of water | $c = 4.18 \text{ kJ kg}^{-1} \text{ K}^{-1}$ (4.18 J g <sup>-1</sup> K <sup>-1</sup> )                                                    |

| Group                    |                           |                                                                |                             |                           |                            |                           |                            |                          |                            |                           |                           |  |  |  |  |  |                     |                      |                        |                      |                           |                           |                           |                            |                          |                         |
|--------------------------|---------------------------|----------------------------------------------------------------|-----------------------------|---------------------------|----------------------------|---------------------------|----------------------------|--------------------------|----------------------------|---------------------------|---------------------------|--|--|--|--|--|---------------------|----------------------|------------------------|----------------------|---------------------------|---------------------------|---------------------------|----------------------------|--------------------------|-------------------------|
| 1                        | 2                         | <div>Key</div>                                                 |                             |                           |                            |                           |                            | <div>1Hhydrogen1.0</div> |                            |                           |                           |  |  |  |  |  | 13                  | 14                   | 15                     | 16                   | 17                        | 18                        |                           |                            |                          |                         |
| 3Li<br>lithium<br>6.9    | 4Be<br>beryllium<br>9.0   | atomic number<br>atomic symbol<br>name<br>relative atomic mass |                             |                           |                            |                           |                            |                          |                            |                           |                           |  |  |  |  |  | 5B<br>boron<br>10.8 | 6C<br>carbon<br>12.0 | 7N<br>nitrogen<br>14.0 | 8O<br>oxygen<br>16.0 | 9F<br>fluorine<br>19.0    | 10Ne<br>neon<br>20.2      |                           |                            |                          |                         |
| 11Na<br>sodium<br>23.0   | 12Mg<br>magnesium<br>24.3 | 3                                                              | 4                           | 5                         | 6                          | 7                         | 8                          | 9                        | 10                         | 11                        | 12                        |  |  |  |  |  |                     |                      |                        |                      | 13Al<br>aluminium<br>27.0 | 14Si<br>silicon<br>28.1   | 15P<br>phosphorus<br>31.0 | 16S<br>sulfur<br>32.1      | 17Cl<br>chlorine<br>35.5 | 18Ar<br>argon<br>39.9   |
| 19K<br>potassium<br>39.1 | 20Ca<br>calcium<br>40.1   | 21Sc<br>scandium<br>45.0                                       | 22Ti<br>titanium<br>47.9    | 23V<br>vanadium<br>50.9   | 24Cr<br>chromium<br>52.0   | 25Mn<br>manganese<br>54.9 | 26Fe<br>iron<br>55.8       | 27Co<br>cobalt<br>58.9   | 28Ni<br>nickel<br>58.7     | 29Cu<br>copper<br>63.5    | 30Zn<br>zinc<br>65.4      |  |  |  |  |  |                     |                      |                        |                      | 31Ga<br>gallium<br>69.7   | 32Ge<br>germanium<br>72.6 | 33As<br>arsenic<br>74.9   | 34Se<br>selenium<br>79.0   | 35Br<br>bromine<br>79.9  | 36Kr<br>krypton<br>83.8 |
| 37Rb<br>rubidium<br>85.5 | 38Sr<br>strontium<br>87.6 | 39Y<br>yttrium<br>88.9                                         | 40Zr<br>zirconium<br>91.2   | 41Nb<br>niobium<br>92.9   | 42Mo<br>molybdenum<br>95.9 | 43Tc<br>technetium<br>—   | 44Ru<br>ruthenium<br>101.1 | 45Rh<br>rhodium<br>102.9 | 46Pd<br>palladium<br>106.4 | 47Ag<br>silver<br>107.9   | 48Cd<br>cadmium<br>112.4  |  |  |  |  |  |                     |                      |                        |                      | 49In<br>indium<br>114.8   | 50Sn<br>tin<br>118.7      | 51Sb<br>antimony<br>121.8 | 52Te<br>tellurium<br>127.6 | 53I<br>iodine<br>126.9   | 54Xe<br>xenon<br>131.3  |
| 55Cs<br>caesium<br>132.9 | 56Ba<br>barium<br>137.3   | 57–71lanthanoids                                               | 72Hf<br>hafnium<br>178.5    | 73Ta<br>tantalum<br>180.9 | 74W<br>tungsten<br>183.8   | 75Re<br>rhenium<br>186.2  | 76Os<br>osmium<br>190.2    | 77Ir<br>iridium<br>192.2 | 78Pt<br>platinum<br>195.1  | 79Au<br>gold<br>197.0     | 80Hg<br>mercury<br>200.6  |  |  |  |  |  |                     |                      |                        |                      | 81Tl<br>thallium<br>204.4 | 82Pb<br>lead<br>207.2     | 83Bi<br>bismuth<br>209.0  | 84Po<br>polonium<br>—      | 85At<br>astatine<br>—    | 86Rn<br>radon<br>—      |
| 87Fr<br>francium<br>—    | 88Ra<br>radium<br>—       | 89–103actinoids                                                | 104Rf<br>rutherfordium<br>— | 105Db<br>dubnium<br>—     | 106Sg<br>seaborgium<br>—   | 107Bh<br>bohrium<br>—     | 108Hs<br>hassium<br>—      | 109Mt<br>meitnerium<br>— | 110Ds<br>darmstadtium<br>— | 111Rg<br>roentgenium<br>— | 112Cn<br>copernicium<br>— |  |  |  |  |  |                     |                      |                        |                      | 113Nh<br>nihonium<br>—    | 114Fl<br>flerovium<br>—   | 115Mc<br>moscovium<br>—   | 116Lv<br>livermorium<br>—  | 117Ts<br>tennessine<br>— | 118Og<br>oganesson<br>— |

|    |    |              |       |
|----|----|--------------|-------|
| 57 | La | lanthanum    | 138.9 |
| 58 | Ce | cerium       | 140.1 |
| 59 | Pr | praseodymium | 140.9 |
| 60 | Nd | neodymium    | 144.4 |
| 61 | Pm | promethium   | —     |
| 62 | Sm | samarium     | 150.4 |
| 63 | Eu | europtium    | 152.0 |
| 64 | Gd | gadolinium   | 157.3 |
| 65 | Tb | terbium      | 158.9 |
| 66 | Dy | dysprosium   | 162.5 |
| 67 | Ho | holmium      | 164.9 |
| 68 | Er | erbium       | 167.3 |
| 69 | Tm | thulium      | 168.9 |
| 70 | Yb | yterbium     | 173.1 |
| 71 | Lu | lutetium     | 175.0 |

|    |    |              |       |
|----|----|--------------|-------|
| 57 | La | lanthanum    | 138.9 |
| 58 | Ce | cerium       | 140.1 |
| 59 | Pr | praseodymium | 140.9 |
| 60 | Nd | neodymium    | 144.4 |
| 61 | Pm | promethium   | —     |
| 62 | Sm | samarium     | 150.4 |
| 63 | Eu | europtium    | 152.0 |
| 64 | Gd | gadolinium   | 157.3 |
| 65 | Tb | terbium      | 158.9 |
| 66 | Dy | dysprosium   | 162.5 |
| 67 | Ho | holmium      | 164.9 |
| 68 | Er | erbium       | 167.3 |
| 69 | Tm | thulium      | 168.9 |
| 70 | Yb | yterbium     | 173.1 |
| 71 | Lu | lutetium     | 175.0 |