

Cambridge Assessment
International Education


Cambridge International AS & A Level

CANDIDATE
NAMECENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--

**CHEMISTRY****9701/22**

Paper 2 AS Level Structured Questions

May/June 2024**1 hour 15 minutes**

You must answer on the question paper.

No additional materials are needed.

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- You may use a calculator.
- You should show all your working and use appropriate units.

INFORMATION

- The total mark for this paper is 60.
- The number of marks for each question or part question is shown in brackets [].
- The Periodic Table is printed in the question paper.
- Important values, constants and standards are printed in the question paper.

This document has **16** pages. Any blank pages are indicated.



2

- 1 (a) Complete Table 1.1 using relevant information from the Periodic Table.

Table 1.1

	nucleon number	proton number	number of electrons
Mg^{2+}	24		
Al^{3+}	27		

[2]

- (b) State and explain the difference in the ionic radius of Al^{3+} compared to Mg^{2+} .

.....

.....

.....

..... [3]

- (c) Draw a labelled diagram to show the structure and bonding in sodium.

[1]

- (d) Fig. 1.1 shows the variation in melting point of some Period 3 elements in their standard states at room temperature and pressure.




Fig. 1.1



* 0019655306503 *



3

- (i) Explain why Si has a high melting point.

.....

 [1]

- (ii) Complete Fig. 1.1 to show the variation in the melting points of the elements P, S and Cl. [2]

- (e) Two Period 3 elements react with an excess of oxygen at room pressure.

- (i) Complete Table 1.2.

Table 1.2

1	2	3
Period 3 element	state of oxide at room temperature and pressure	approximate pH of solution made when oxide is added to water
Na		
S		

[2]

- (ii) The solutions made in column 3 of Table 1.2 are mixed together. Name the type of reaction that occurs.

..... [1]

- (iii) Write an equation to describe the reaction between P_4O_{10} and an excess of water.

..... [1]

- (f) Aluminium hydroxide is amphoteric.

- (i) Explain what is meant by amphoteric.

.....
 [1]

- (ii) Write an equation to describe the reaction that occurs when aluminium hydroxide, $Al(OH)_3$, reacts with $NaOH(aq)$.

..... [1]

[Total: 15]





- 2 Separate samples of Na_2CO_3 and NaHCO_3 react with $\text{HCl}(\text{aq})$ to produce the same products, as shown in Table 2.1.

Table 2.1

reaction	equation	$\Delta H/\text{kJ mol}^{-1}$
1	$\text{Na}_2\text{CO}_3 + 2\text{HCl} \rightarrow 2\text{NaCl} + \text{H}_2\text{O} + \text{CO}_2$	ΔH_1
2	$\text{NaHCO}_3 + \text{HCl} \rightarrow \text{NaCl} + \text{H}_2\text{O} + \text{CO}_2$	$\Delta H_2 = +27.2$

- (a) Complete the reaction pathway diagram in Fig. 2.1 for reaction 2.

Label the diagram to show the enthalpy change, ΔH_2 , and the activation energy, E_A .




Fig. 2.1

[2]

- (b) The value for ΔH_1 is determined by experiment using the following method.

- 50.0 cm^3 of 2.00 mol dm^{-3} $\text{HCl}(\text{aq})$ is added to a polystyrene cup.
- The initial temperature of the acid is recorded as 19.6°C .
- 0.0400 mol of Na_2CO_3 is added and the mixture is stirred.
- All the solid Na_2CO_3 disappears and a colourless solution is produced.

The maximum temperature recorded during the reaction is 26.2°C .

- (i) Describe **one** other observation that shows the reaction is complete.

..... [1]





5


(ii) Calculate the value of ΔH_1 in kJ mol^{-1} .

Assume the specific heat capacity of the reaction mixture is the same as for water and no heat is lost to the surroundings.

Show your working.

$$\Delta H_1 = \dots\dots\dots \text{kJ mol}^{-1} \quad [3]$$

(iii) Thermal decomposition occurs when NaHCO_3 is heated.



Calculate the enthalpy change for reaction 3, ΔH_r , using the data in Table 2.1 and the value of ΔH_1 calculated in (b)(ii).

(If you were unable to calculate a value for ΔH_1 in (b)(ii), assume the enthalpy change is $-38.4 \text{ kJ mol}^{-1}$. This is **not** the correct value.)

$$\Delta H_r = \dots\dots\dots \text{kJ mol}^{-1} \quad [2]$$

(c) **Z** is a salt that contains a Period 4 element from Group 2. When **Z** is heated brown gas forms.

Identify the formula of **Z** and use it to write an equation for the reaction.

..... [2]

[Total: 10]






3 (a) Describe what is meant by dynamic equilibrium.

.....

.....

..... [2]

(b) Reaction 4 describes the reversible reaction between yellow $\text{Fe}^{3+}(\text{aq})$ and colourless $\text{SCN}^{-}(\text{aq})$ to produce red $\text{FeSCN}^{2+}(\text{aq})$.



An equilibrium mixture contains $\text{Fe}^{3+}(\text{aq})$, $\text{SCN}^{-}(\text{aq})$ and $\text{FeSCN}^{2+}(\text{aq})$. A few colourless crystals of soluble $\text{KSCN}(\text{s})$ are added. The mixture is then left until it reaches equilibrium again. The temperature of both equilibrium mixtures is the same.

(i) Deduce the changes that occur, if any, in the equilibrium mixture after $\text{KSCN}(\text{s})$ is added compared to the original equilibrium mixture.

- change in appearance

.....

- change in relative concentration of $\text{Fe}^{3+}(\text{aq})$

.....

- change in value of the equilibrium constant, K_c

.....

[3]





(ii) The expression for the equilibrium constant, K_c , for reaction 4 is shown.

$$K_c = \frac{[\text{FeSCN}^{2+}(\text{aq})]}{[\text{Fe}^{3+}(\text{aq})] \times [\text{SCN}^{-}(\text{aq})]}$$

$5.00 \times 10^{-5} \text{ mol}$ of $\text{Fe}^{3+}(\text{aq})$ and $5.00 \times 10^{-5} \text{ mol}$ of $\text{SCN}^{-}(\text{aq})$ are added together and allowed to reach equilibrium. The total volume of the mixture is 25.0 cm^3 .

At equilibrium the concentration of $\text{FeSCN}^{2+}(\text{aq})$ is $4.23 \times 10^{-4} \text{ mol dm}^{-3}$.

Calculate the equilibrium constant, K_c , for reaction 4.

Include the units in your answer.

$K_c = \dots\dots\dots$

units $\dots\dots\dots$

[4]





(c) Determine the full electronic configuration of Fe^{3+} .

..... [1]

(d) $\text{SCN}^{-}(\text{aq})$ is colourless.

Complete the dot-and-cross diagram in Fig. 3.1 to show the arrangement of outer electrons in an SCN^{-} ion.




Fig. 3.1

[2]

[Total: 12]



**BLANK PAGE**



4 $\text{CH}_3(\text{CH}_2)_5\text{CHBrCH}_3$ exists as a pair of stereoisomers.

- (a) Draw the three-dimensional structures of the **two** stereoisomers of $\text{CH}_3(\text{CH}_2)_5\text{CHBrCH}_3$. R can be used to represent $\text{CH}_3(\text{CH}_2)_5$.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
84

[2]

- (b) A sample of $\text{CH}_3(\text{CH}_2)_5\text{CHBrCH}_3$ reacts with NaOH to make $\text{CH}_3(\text{CH}_2)_5\text{CH}(\text{OH})\text{CH}_3$ in an $\text{S}_{\text{N}}1$ mechanism.

Complete Fig. 4.1 to show the mechanism for the reaction of $\text{CH}_3(\text{CH}_2)_5\text{CHBrCH}_3$ and NaOH .

Include charges, dipoles, lone pairs of electrons and curly arrows, as appropriate.



Fig. 4.1

[3]

- (c) Separate samples of $\text{CH}_3(\text{CH}_2)_5\text{CHBrCH}_3$, $\text{CH}_3(\text{CH}_2)_5\text{CH}(\text{OH})\text{CH}_3$ and $\text{CH}_3(\text{CH}_2)_5\text{CHCH}_2$ are tested with different reagents.

Complete Table 4.1. If no reaction occurs, write x in the relevant box.

Table 4.1

reagent added	observation with $\text{CH}_3(\text{CH}_2)_5\text{CHBrCH}_3$	observation with $\text{CH}_3(\text{CH}_2)_5\text{CH(OH)CH}_3$	observation with $\text{CH}_3(\text{CH}_2)_5\text{CHCH}_2$
$\text{Br}_2(\text{l})$ in the dark			
$\text{PCl}_5(\text{s})$			
$\text{AgNO}_3(\text{aq})$			

[3]

* 0019655306611 *



11

(d) $\text{CH}_3(\text{CH}_2)_5\text{CHBrCH}_3$ is heated with **D** to produce three different molecules, **E**, **F** and **G**.




Fig. 4.2

(i) Name the type of reaction.

..... [1]

(ii) Identify **D** and the conditions used.

..... [1]

(e) (i) Both σ and π bonds are present in a molecule of **E** as a result of different types of hybridisation in the carbon atoms.

Complete Table 4.2 to show the number of carbon atoms with each type of hybridisation in a molecule of **E**.

Table 4.2

	number of carbon atoms		
	sp hybridised	sp^2 hybridised	sp^3 hybridised
E			

[2]

(ii) Describe the essential feature of an unbranched hydrocarbon that causes its molecules to show stereoisomerism. Explain how this feature leads to stereoisomerism.

.....

.....

..... [3]

[Total: 15]





5 Compound **W** has molecular formula $C_4H_{10}O$. It contains only **one** functional group.

(a) Table 5.1 shows the two peaks with the greatest m/e values in the mass spectrum of **W**.

Table 5.1

m/e	relative abundance
74	50
75	x

(i) Calculate the relative abundance, x, of the peak at $m/e = 75$ using the information from Table 5.1.

x = [1]

(ii) The mass spectrum of **W** also shows peaks at $m/e = 29$ and $m/e = 59$.

Suggest the molecular formulae of these fragments.

$m/e = 29$

$m/e = 59$

[2]

(b) A sample of **W**, $C_4H_{10}O$, is heated under reflux with an excess of acidified $K_2Cr_2O_7$ until there is no further reaction. Only **one** organic product, **X**, is present in the mixture at the end of the reaction.

Fig. 5.1 shows the infrared spectrum of **W**.




Fig. 5.1





13


Fig. 5.2 shows the infrared spectrum of **X**.

Fig. 5.2

Table 5.2

bond	functional groups containing the bond	characteristic infrared absorption range (in wavenumbers) / cm^{-1}
C–O	hydroxy, ester	1040–1300
C=C	aromatic compound, alkene	1500–1680
C=O	amide carbonyl, carboxyl ester	1640–1690 1670–1740 1710–1750
C≡N	nitrile	2200–2250
C–H	alkane	2850–2950
N–H	amine, amide	3300–3500
O–H	carboxyl hydroxy	2500–3000 3200–3650

- (i) Absorption **A** is shown in Fig. 5.1.
Absorption **B** is shown in Fig. 5.2.

Complete Table 5.3 using the information given in Fig. 5.1, Fig. 5.2 and Table 5.2.

Table 5.3

absorption	bond	functional group containing the bond
A		
B		

[1]





14

(ii) Use the information in (a) and (b)(i) to draw the structure of **X** in the box in Fig. 5.3.




Fig. 5.3

[1]

(c) **Y** is a structural isomer of **W**.

Both **W** and **Y** produce colourless bubbles when sodium is added to them.

Y does **not** react when heated with acidified $\text{K}_2\text{Cr}_2\text{O}_7$.

Y does **not** react when warmed with alkaline $\text{I}_2(\text{aq})$.

(i) Name the functional group present in **Y**.

..... [1]

(ii) Complete the equation to describe the reaction of **W** or **Y** with sodium.

..... $\text{C}_4\text{H}_{10}\text{O}$ + Na → [1]

(iii) Draw the structure of **Y**.

[1]

[Total: 8]



* 0019655306615 *



15

**Important values, constants and standards**

molar gas constant	$R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$
Faraday constant	$F = 9.65 \times 10^4 \text{ C mol}^{-1}$
Avogadro constant	$L = 6.022 \times 10^{23} \text{ mol}^{-1}$
electronic charge	$e = -1.60 \times 10^{-19} \text{ C}$
molar volume of gas	$V_m = 22.4 \text{ dm}^3 \text{ mol}^{-1}$ at s.t.p. (101 kPa and 273 K) $V_m = 24.0 \text{ dm}^3 \text{ mol}^{-1}$ at room conditions
ionic product of water	$K_w = 1.00 \times 10^{-14} \text{ mol}^2 \text{ dm}^{-6}$ (at 298 K (25 °C))
specific heat capacity of water	$c = 4.18 \text{ kJ kg}^{-1} \text{ K}^{-1}$ (4.18 J g ⁻¹ K ⁻¹)





Group																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																				
1	2	Key															18																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																			
		atomic number atomic symbol name relative atomic mass																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																		
3	4																2																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																			
Li lithium 6.9	Be beryllium 9.0	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57–71 lanthanoids	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89–103 actinoids	104	105	106	107	108	109	110	111	112	113	114	115	116	117	118	119	120	121	122	123	124	125	126	127	128	129	130	131	132	133	134	135	136	137	138	139	140	141	142	143	144	145	146	147	148	149	150	151	152	153	154	155	156	157	158	159	160	161	162	163	164	165	166	167	168	169	170	171	172	173	174	175	176	177	178	179	180	181	182	183	184	185	186	187	188	189	190	191	192	193	194	195	196	197	198	199	200	201	202	203	204	205	206	207	208	209	210	211	212	213	214	215	216	217	218	219	220	221	222	223	224	225	226	227	228	229	230	231	232	233	234	235	236	237	238	239	240	241	242	243	244	245	246	247	248	249	250	251	252	253	254	255	256	257	258	259	260	261	262	263	264	265	266	267	268	269	270	271	272	273	274	275	276	277	278	279	280	281	282	283	284	285	286	287	288	289	290	291	292	293	294	295	296	297	298	299	300	301	302	303	304	305	306	307	308	309	310	311	312	313	314	315	316	317	318	319	320	321	322	323	324	325	326	327	328	329	330	331	332	333	334	335	336	337	338	339	340	341	342	343	344	345	346	347	348	349	350	351	352	353	354	355	356	357	358	359	360	361	362	363	364	365	366	367	368	369	370	371	372	373	374	375	376	377	378	379	380	381	382	383	384	385	386	387	388	389	390	391	392	393	394	395	396	397	398	399	400	401	402	403	404	405	406	407	408	409	410	411	412	413	414	415	416	417	418	419	420	421	422	423	424	425	426	427	428	429	430	431	432	433	434	435	436	437	438	439	440	441	442	443	444	445	446	447	448	449	450	451	452	453	454	455	456	457	458	459	460	461	462	463	464	465	466	467	468	469	470	471	472	473	474	475	476	477	478	479	480	481	482	483	484	485	486	487	488	489	490	491	492	493	494	495	496	497	498	499	500	501	502	503	504	505	506	507	508	509	510	511	512	513	514	515	516	517	518	519	520	521	522	523	524	525	526	527	528	529	530	531	532	533	534	535	536	537	538

lanthanoids

actinoids

57	La	lanthanum	138.9
58	Ce	cerium	140.1
59	Pr	praseodymium	140.9
60	Nd	neodymium	144.2
61	Pm	promethium	—
62	Sm	samarium	150.4
63	Eu	euroium	152.0
64	Gd	gadolinium	157.3
65	Tb	terbium	158.9
66	Dy	dysprosium	162.5
67	Ho	holmium	164.9
68	Er	erbium	167.3
69	Tm	thulium	168.9
70	Yb	ytterbium	173.1
71	Lu	lutetium	175.0

© UCLES 2024

