

Cambridge
International
A Level

Cambridge Assessment International Education
Cambridge International Advanced Level

CANDIDATE
NAME

--

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--

FURTHER MATHEMATICS

9231/22

Paper 2

May/June 2019

3 hours

Candidates answer on the Question Paper.

Additional Materials: List of Formulae (MF10)

READ THESE INSTRUCTIONS FIRST

Write your centre number, candidate number and name in the spaces at the top of this page.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Answer **all** the questions in the space provided. If additional space is required, you should use the lined page at the end of this booklet. The question number(s) must be clearly shown.

Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.

Where a numerical value is necessary, take the acceleration due to gravity to be 10 m s^{-2} .

The use of a calculator is expected, where appropriate.

Results obtained solely from a graphic calculator, without supporting working or reasoning, will not receive credit.

You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.

This document consists of **25** printed pages and **3** blank pages.



BLANK PAGE

3

1 A particle P moves along an arc of a circle with centre O and radius 2 m. At time t seconds, the angle POA is θ , where $\theta = 1 - \cos 2t$, and A is a fixed point on the arc of the circle.

(i) Show that the magnitude of the radial component of the acceleration of P when $t = \frac{1}{6}\pi$ is 6 m s^{-2} . [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(ii) Find the magnitude of the transverse component of the acceleration of P when $t = \frac{1}{6}\pi$. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

2 A particle P moves on a straight line in simple harmonic motion. The centre of the motion is O . The points A and B are on the line on opposite sides of O such that $OA = 3.5\text{ m}$ and $OB = 1\text{ m}$. The speed of P when it is at B is twice its speed when it is at A . The maximum acceleration of P is 1 m s^{-2} .

(i) Find the speed of P when it is at O . [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

[4]

This image shows a full page of white paper with horizontal dotted lines. The lines are evenly spaced and run across the width of the page, providing a guide for handwriting practice. There are no margins, text, or other markings on the page.

3 Three uniform small spheres A , B and C have equal radii and masses $2m$, $4m$ and m respectively. The spheres are moving in a straight line on a smooth horizontal surface, with B between A and C . The coefficient of restitution between each pair of spheres is e . Spheres A and B are moving towards each other with speeds $2u$ and u respectively. The first collision is between A and B .

(i) Find the velocities of A and B after this collision. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

Sphere C is moving towards B with speed $\frac{4}{3}u$ and now collides with it. As a result of this collision, B is brought to rest.

(ii) Find the value of e . [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(iii) Find the total kinetic energy lost by the three spheres as a result of the two collisions. [3]

.....

.....

.....

.....

.....

.....

.....


.....

.....

.....

.....

4




A uniform rod AB of length $4a$ and weight W rests with the end A in contact with a rough vertical wall. A light inextensible string of length $\frac{5}{2}a$ has one end attached to the point C on the rod, where $AC = \frac{5}{2}a$. The other end of the string is attached to a point D on the wall, vertically above A . The vertical plane containing the rod AB is perpendicular to the wall. The angle between the rod and the wall is θ , where $\tan \theta = 2$ (see diagram). The end A of the rod is on the point of slipping down the wall and the coefficient of friction between the rod and the wall is μ .

Find, in either order, the tension in the string and the value of μ .

[10]

[illegible]

5



A thin uniform rod AB has mass kM and length $2a$. The end A of the rod is rigidly attached to the surface of a uniform hollow sphere with centre O , mass kM and radius $2a$. The end B of the rod is rigidly attached to the circumference of a uniform ring with centre C , mass M and radius a . The points C, B, A, O lie in a straight line. The horizontal axis L passes through the mid-point of the rod and is perpendicular to the rod and in the plane of the ring (see diagram). The object consisting of the rod, the ring and the hollow sphere can rotate freely about L .

- (i) Show that the moment of inertia of the object about L is $\frac{3}{2}(8k + 3)Ma^2$. [6]

[illegible]

The object performs small oscillations about L , with the ring above the sphere as shown in the diagram.

- (ii) Find the set of possible values of k and the period of these oscillations in terms of k . [6]

This image shows a single page of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

12

- 6** A fair six-sided die is thrown until a 3 or a 4 is obtained. The number of throws taken is denoted by the random variable X .

(i) State the mean value of X . [1]

.....

.....

.....

.....

(ii) Find the probability that obtaining a 3 or a 4 takes exactly 6 throws. [1]

.....

.....

.....

.....

.....

(iii) Find the probability that obtaining a 3 or a 4 takes more than 4 throws. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (iv) Find the greatest integer n such that the probability of obtaining a 3 or a 4 in fewer than n throws is less than 0.95. [3]

This image shows a full page of primary-ruled paper. It features multiple sets of horizontal dashed lines spaced evenly down the page, providing a guide for handwriting practice. The lines are thin and black, set against a plain white background. There are no margins, text, or other markings on the page.

7 The continuous random variable X has probability density function f given by

$$f(x) = \begin{cases} \frac{3}{4x^2} + \frac{1}{4} & 1 \leq x \leq 3, \\ 0 & \text{otherwise.} \end{cases}$$

(i) Find the distribution function of X . [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

15

(ii) Find the exact value of the interquartile range of X . [5]

This image shows a full page of a document template designed for handwriting practice. It consists of approximately 20 evenly spaced horizontal dotted lines across the entire width of the page, providing a guide for letter height and placement. The background is plain white, and there are no margins, text, or other markings present.

Runner	<i>A</i>	<i>B</i>	<i>C</i>	<i>D</i>	<i>E</i>	<i>F</i>
Time at beginning of camp	3.82	3.62	3.55	3.71	3.75	3.92
Time at end of camp	3.72	3.55	3.52	3.68	3.54	3.73

The organiser of the training camp claims that a runner's time will improve by more than 0.05 minutes between the beginning and end of the camp. Assuming that differences in time over the two runs are normally distributed, test at the 10% significance level whether the organiser's claim is justified. [8]

This image shows a full page of white paper with horizontal dashed lines, typical of primary-ruled notebook paper. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

This image shows a full page of white paper with horizontal dashed lines, typical of primary-ruled notebook paper. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

It is required to test the goodness of fit of the distribution with probability density function f given by

The relevant expected frequencies, correct to 2 decimal places, are given in the following table.

(i) Show how the expected frequency for $1.6 \leq x < 2.4$ is obtained. [3]

This image shows a full page of a worksheet designed for handwriting practice. It features 18 evenly spaced, horizontal dashed lines across the entire width of the page. The background is plain white, providing a clear guide for letter height and placement. There are no margins, text, or other markings present.

(ii) Carry out a goodness of fit test at the 5% significance level.

[7]

This image shows a full page of white paper with horizontal dashed lines, typical of primary school writing paper. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

x	3	4	4	6	8
y	5	7	q	6	7

(i) Given that q is an integer, find its value.

[5]

[illegible]

21**(ii)** Find the value of c .**[3]**

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(iii) Find the value of the product moment correlation coefficient.**[3]**

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

11 Answer only **one** of the following two alternatives.

EITHER

A particle P , of mass m , is able to move in a vertical circle on the smooth inner surface of a sphere with centre O and radius a . Points A and B are on the inner surface of the sphere and AOB is a horizontal diameter. Initially, P is projected vertically downwards with speed $\sqrt{\left(\frac{21}{2}ag\right)}$ from A and begins to move in a vertical circle. At the lowest point of its path, vertically below O , the particle P collides with a stationary particle Q , of mass $4m$, and rebounds. The speed acquired by Q , as a result of the collision, is just sufficient for it to reach the point B .

(i) Find the speed of P and the speed of Q immediately after their collision. [7]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

[5]

[illegible]

OR

A farmer grows two different types of cherries, Type *A* and Type *B*. He assumes that the masses of each type are normally distributed. He chooses a random sample of 8 cherries of Type *A*. He finds that the sample mean mass is 15.1 g and that a 95% confidence interval for the population mean mass, μ g, is $13.5 \leq \mu \leq 16.7$.

- (i) Find an unbiased estimate for the population variance of the masses of cherries of Type *A*. [3]

.....

.....

.....

.....

.....

.....

.....

.....

The farmer now chooses a random sample of 6 cherries of Type *B* and records their masses as follows.

12.2 13.3 16.4 14.0 13.9 15.4

- (ii) Test at the 5% significance level whether the mean mass of cherries of Type *B* is less than the mean mass of cherries of Type *A*. You should assume that the population variances for the two types of cherry are equal. [9]

.....

.....

.....

.....

.....

.....

.....

.....

This image shows a full page of white paper with horizontal dashed lines, typical of primary school writing paper. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

This image shows a full page of a worksheet designed for handwriting practice. It features 20 evenly spaced, horizontal dashed lines across the entire width of the page. The background is plain white, providing a clear guide for letter formation and alignment. There are no margins, text, or other markings present.

BLANK PAGE

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.