



Cambridge International AS & A Level

CANDIDATE
NAME

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--

FURTHER MATHEMATICS

9231/13

Paper 1 Further Pure Mathematics 1

May/June 2020

2 hours

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [].

This document has **16** pages. Blank pages are indicated.

1 The cubic equation $7x^3 + 3x^2 + 5x + 1 = 0$ has roots α, β, γ .

(a) Find a cubic equation whose roots are $\alpha^{-1}, \beta^{-1}, \gamma^{-1}$. [3]

.....

.....

.....

.....

.....

.....

.....

(b) Find the value of $\alpha^{-2} + \beta^{-2} + \gamma^{-2}$. [2]

.....

.....

.....

.....

.....

.....

.....

(c) Find the value of $\alpha^{-3} + \beta^{-3} + \gamma^{-3}$. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

3

2 The sequence $u_1, u_2, u_3, \dots$ is such that $u_1 = 1$ and $u_{n+1} = 2u_n + 1$ for $n \geq 1$.

(a) Prove by induction that $u_n = 2^n - 1$ for all positive integers n . [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Deduce that u_{2n} is divisible by u_n for $n \geq 1$. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

This image shows a full page of a handwriting practice worksheet. It consists of multiple sets of three horizontal dashed lines, providing a guide for letter height and placement. The lines are evenly spaced across the entire page, leaving ample room for practicing various letters and words. There is no text or other markings on the page.

(b) Express $\frac{n}{S_n}$ in partial fractions and find $\sum_{n=1}^N \frac{n}{S_n}$ in terms of N . [4]

[illegible]

(c) Deduce the value of $\sum_{n=1}^{\infty} \frac{n}{S_n}$. [1]

[illegible]

6**4** The matrix **A** is given by

$$\mathbf{A} = \begin{pmatrix} k & 0 & 2 \\ 0 & -1 & -1 \\ 1 & 1 & -k \end{pmatrix},$$

where k is a real constant.**(a)** Show that **A** is non-singular.

[3]

.....

.....

.....

.....

.....

.....

.....

.....

The matrices **B** and **C** are given by

$$\mathbf{B} = \begin{pmatrix} 0 & -3 \\ -1 & 3 \\ 0 & 0 \end{pmatrix} \text{ and } \mathbf{C} = \begin{pmatrix} -3 & -1 & 1 \\ 1 & 1 & 2 \end{pmatrix}.$$

It is given that $\mathbf{CAB} = \begin{pmatrix} -2 & -\frac{3}{2} \\ -1 & -\frac{3}{2} \end{pmatrix}.$

(b) Find the value of k .

[3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

[illegible]

8

5 The curve C has polar equation $r = a \tan \theta$, where a is a positive constant and $0 \leq \theta \leq \frac{1}{4}\pi$.

(a) Sketch C and state the greatest distance of a point on C from the pole. [2]

(b) Find the exact value of the area of the region bounded by C and the half-line $\theta = \frac{1}{4}\pi$. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

[illegible]

(d) Using your answer to part (b), deduce the exact value of $\int_0^{\frac{1}{2}a\sqrt{2}} \frac{x^2}{\sqrt{a^2 - x^2}} dx$. [2]

[illegible]

6 The curve C has equation $y = \frac{10 + x - 2x^2}{2x - 3}$.

(a) Find the equations of the asymptotes of C . [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Show that C has no turning points. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

12

(c) Sketch C , stating the coordinates of the intersections with the axes.

[3]

- (d) Sketch the curve with equation $y = \left| \frac{10+x-2x^2}{2x-3} \right|$ and find in exact form the set of values of x for which $\left| \frac{10+x-2x^2}{2x-3} \right| < 4$. [6]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

7 The lines l_1 and l_2 have equations $\mathbf{r} = -5\mathbf{j} + \lambda(5\mathbf{i} + 2\mathbf{k})$ and $\mathbf{r} = 4\mathbf{i} + 2\mathbf{j} - 2\mathbf{k} + \mu(\mathbf{j} + \mathbf{k})$ respectively. The plane Π contains l_1 and is parallel to l_2 .

(a) Find the equation of Π , giving your answer in the form $ax + by + cz = d$. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Find the distance between l_2 and Π . [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

-
- This image shows a full page of white paper with horizontal dotted lines. The lines are evenly spaced and run across the width of the page, providing a guide for handwriting practice. There are no margins, text, or other markings on the page.

[illegible]

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.