



Cambridge International AS & A Level

FURTHER MATHEMATICS**9231/33**

Paper 3 Further Mechanics

May/June 2022**MARK SCHEME**Maximum Mark: 50

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the May/June 2022 series for most Cambridge IGCSE, Cambridge International A and AS Level and Cambridge Pre-U components, and some Cambridge O Level components.

This document consists of **13** printed pages.

PUBLISHED**Generic Marking Principles**

These general marking principles must be applied by all examiners when marking candidate answers. They should be applied alongside the specific content of the mark scheme or generic level descriptors for a question. Each question paper and mark scheme will also comply with these marking principles.

GENERIC MARKING PRINCIPLE 1:

Marks must be awarded in line with:

- the specific content of the mark scheme or the generic level descriptors for the question
- the specific skills defined in the mark scheme or in the generic level descriptors for the question
- the standard of response required by a candidate as exemplified by the standardisation scripts.

GENERIC MARKING PRINCIPLE 2:

Marks awarded are always **whole marks** (not half marks, or other fractions).

GENERIC MARKING PRINCIPLE 3:

Marks must be awarded **positively**:

- marks are awarded for correct/valid answers, as defined in the mark scheme. However, credit is given for valid answers which go beyond the scope of the syllabus and mark scheme, referring to your Team Leader as appropriate
- marks are awarded when candidates clearly demonstrate what they know and can do
- marks are not deducted for errors
- marks are not deducted for omissions
- answers should only be judged on the quality of spelling, punctuation and grammar when these features are specifically assessed by the question as indicated by the mark scheme. The meaning, however, should be unambiguous.

GENERIC MARKING PRINCIPLE 4:

Rules must be applied consistently, e.g. in situations where candidates have not followed instructions or in the application of generic level descriptors.

GENERIC MARKING PRINCIPLE 5:

Marks should be awarded using the full range of marks defined in the mark scheme for the question (however; the use of the full mark range may be limited according to the quality of the candidate responses seen).

GENERIC MARKING PRINCIPLE 6:

Marks awarded are based solely on the requirements as defined in the mark scheme. Marks should not be awarded with grade thresholds or grade descriptors in mind.

Mathematics Specific Marking Principles	
1	Unless a particular method has been specified in the question, full marks may be awarded for any correct method. However, if a calculation is required then no marks will be awarded for a scale drawing.
2	Unless specified in the question, answers may be given as fractions, decimals or in standard form. Ignore superfluous zeros, provided that the degree of accuracy is not affected.
3	Allow alternative conventions for notation if used consistently throughout the paper, e.g. commas being used as decimal points.
4	Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored (isw).
5	Where a candidate has misread a number in the question and used that value consistently throughout, provided that number does not alter the difficulty or the method required, award all marks earned and deduct just 1 mark for the misread.
6	Recovery within working is allowed, e.g. a notation error in the working where the following line of working makes the candidate's intent clear.

PUBLISHED**Mark Scheme Notes**

The following notes are intended to aid interpretation of mark schemes in general, but individual mark schemes may include marks awarded for specific reasons outside the scope of these notes.

Types of mark

- M** Method mark, awarded for a valid method applied to the problem. Method marks are not lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. Correct application of a formula without the formula being quoted obviously earns the M mark and in some cases an M mark can be implied from a correct answer.
- A** Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated method mark is earned (or implied).
- B** Mark for a correct result or statement independent of method marks.
- DM or DB** When a part of a question has two or more ‘method’ steps, the M marks are generally independent unless the scheme specifically says otherwise; and similarly, when there are several B marks allocated. The notation DM or DB is used to indicate that a particular M or B mark is dependent on an earlier M or B (asterisked) mark in the scheme. When two or more steps are run together by the candidate, the earlier marks are implied and full credit is given.
- FT** Implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A or B marks are given for correct work only.
- A or B marks are given for correct work only (not for results obtained from incorrect working) unless follow through is allowed (see abbreviation FT above).
 - For a numerical answer, allow the A or B mark if the answer is correct to 3 significant figures or would be correct to 3 significant figures if rounded (1 decimal place for angles in degrees).
 - The total number of marks available for each question is shown at the bottom of the Marks column.
 - Wrong or missing units in an answer should not result in loss of marks unless the guidance indicates otherwise.
 - Square brackets [] around text or numbers show extra information not needed for the mark to be awarded.

Abbreviations

AEF/OE	Any Equivalent Form (of answer is equally acceptable) / Or Equivalent
AG	Answer Given on the question paper (so extra checking is needed to ensure that the detailed working leading to the result is valid)
CAO	Correct Answer Only (emphasising that no ‘follow through’ from a previous error is allowed)
CWO	Correct Working Only
ISW	Ignore Subsequent Working
SOI	Seen Or Implied
SC	Special Case (detailing the mark to be given for a specific wrong solution, or a case where some standard marking practice is to be varied in the light of a particular circumstance)
WWW	Without Wrong Working
AWRT	Answer Which Rounds To

PUBLISHED

Question	Answer			Marks	Guidance
1		Area	Distance from Oy	M1	Attempt at moments equation with all necessary terms. Other options possible for RHS of moments equation, for example: (1) OAC : 30×6 and ABC : 12×9 (2) OBC : 12×4 and OAB : 30×8 (3) Subtraction: $60 \times 7.5 - 6 \times 1 - 12 \times 13$
	Triangle $ OCD $	6	2		
	Rectangle $ DEBC $	24	6		
	Triangle $ BAE $	12	11		
	Trapezium $ OCBA $	42	$\bar{x}$		
	where D is point with coordinates (3, 0) and E is a point with coordinates (9, 0).				
	Parts that would give correct total area 42			B1	
	Moments about Oy $42\bar{x} = 6 \times 2 + 24 \times 6 + 12 \times 11$ (=288)			A1	Correct equation.
	$\bar{x} = \frac{288}{42} = 6.86$			A1	
				4	

PUBLISHED

Question	Answer	Marks	Guidance
2	$\frac{1}{2} \times \frac{\frac{4}{3}mg}{a} x^2$	B1	EPE term correct
	$\frac{1}{3}mgx$	B1	Work term correct
	Loss in KE = gain in EPE + work done against friction $\frac{1}{2}mv^2 = \frac{1}{2} \times \frac{\frac{4}{3}mg}{a} x^2 + \frac{1}{3}mgx$	M1	Energy equation with 3 terms, allow sign error.
	$\frac{1}{2} \times \frac{1}{4}ga = \frac{\frac{2}{3}g}{a}x^2 + \frac{1}{3}gx$ $16x^2 + 8ax - 3a^2 = 0$ $(4x - a)(4x + 3a) = 0$	M1	Obtain and attempt to solve a 3-term quadratic equation.
	$x = \frac{1}{4}a$	A1	
		5	

PUBLISHED

Question	Answer	Marks	Guidance
3(a)	Components of velocity : $\rightarrow 25 \cos \theta$ $\uparrow 25 \sin \theta - 2g$	B1	
	Speed = $\sqrt{(25 \cos \theta)^2 + (25 \sin \theta - 2g)^2}$	M1 A1	Expression for speed or square of speed.
	$(25 \cos \theta)^2 + (25 \sin \theta - 2g)^2 = 15^2$ $625 - 100g \sin \theta + 4g^2 = 225$	M1	Attempt to solve and find value for $\sin \theta$
	$\sin \theta = \frac{800}{1000} = \frac{4}{5}$	A1	
		5	
3(b)	Time of flight = $\left(\frac{2 \times 25 \sin \theta}{g} \right) = 4 \text{ (s)}$	B1	
	Range = $\frac{2 \times 25 \sin \theta}{g} \times 25 \cos \theta$	M1	Any equivalent method.
	Range = 60 (m)	A1	CWO
	Alternative method for question 3(b)		
	$y = \frac{4}{3}x - \frac{1}{45}x^2$	B1	Equation of trajectory..
	Substitute $y = 0$ and solve	M1	
	60 (m)	A1	
		3	

PUBLISHED

Question	Answer	Marks	Guidance
4(a)	$\frac{1}{2}mv^2 - \frac{1}{2}mu^2 = mga(\cos\theta + \cos\alpha)$	M1	Energy equation with all necessary terms, GPE terms must be resolved, allow sin/cos mix, allow sign error.
	$\frac{1}{2}m(2u)^2 - \frac{1}{2}mu^2 = mga(\cos\theta + \cos\alpha)$	A1	2u may be substituted later. Implied by $\frac{3}{2} \times \frac{2}{3} ag = ga(\cos\theta + \cos\alpha)$
	At A, $T + mg \cos\theta = \frac{m}{a}u^2$	B1	N2L
	Also, $10T - mg \cos\alpha = \frac{m}{a}4u^2$	B1	N2L and use of tension (10T).
	Use all three (two N2L and energy) equations to find T in terms of m and g only.	M1	Might see $9T - mg(\cos\theta + \cos\alpha) = \frac{3m}{a} \times \frac{2}{3} ga$ $(\cos\theta + \cos\alpha) = 1$ $(10\cos\theta + \cos\alpha) = 4$
	$T = \frac{1}{3}mg$	A1	
		6	
4(b)	Substitute back, $10 \times \frac{1}{3}mg - mg \cos\alpha = \frac{4m}{a} \times \frac{2}{3} ga$	M1	Any appropriate method to obtain $\cos\alpha$.
	$\cos\alpha = \frac{2}{3}$	A1	
		2	

PUBLISHED

Question	Answer	Marks	Guidance
5(a)	$4v \frac{dv}{dx} = -(4e^{-x} + 12)e^{-x}$	B1	
	$\frac{1}{2}v^2 = \frac{1}{2}e^{-2x} + 3e^{-x} (+A)$	M1	Expression of the correct form.
	$v = 4, \quad x = 0, \quad A = \frac{9}{2}$	A1	
	$v^2 = e^{-2x} + 6e^{-x} + 9 = (3 + e^{-x})^2$ $v = 3 + e^{-x} = \frac{1 + 3e^x}{e^x}$	A1	AG Must see the factorisation. Condone lack of justification for taking positive square root.
		4	
5(b)	$\frac{dx}{dt} = \frac{1 + 3e^x}{e^x}$ so $\int \frac{e^x}{3e^x + 1} dx = \int 1 dt$ $\frac{1}{3} \ln(3e^x + 1) = t (+B)$	M1* A1	Integration to obtain ln term Correct answer with constant of integration
	$t = 0, \quad x = 0, \quad B = \frac{1}{3} \ln 4$ $3t = \ln \frac{3e^x + 1}{4}$	DM1	Find the constant and substitute into their general solution.
	$x = \ln \left(\frac{4}{3} e^{3t} - \frac{1}{3} \right)$	A1	OE
		4	

PUBLISHED

Question	Answer	Marks	Guidance
6(a)	Let v be speed of rebound from 1 st collision: Energy loss: $\frac{1}{2}mv^2 = \frac{2}{5} \times \frac{1}{2}mu^2$, $v^2 = \frac{2}{5}u^2$	B1	Energy loss.
	$v \cos \alpha = u \cos \theta$ $v \sin \alpha = eu \sin \theta$	B1	Both.
	Combine to form equation in e only $\frac{2}{5} = \frac{1}{5} + e^2 \times \frac{4}{5}$	M1	$v^2 = (u \cos \theta)^2 + (eu \sin \theta)^2$
	$e = \frac{1}{2}$	A1	
		4	
6(b)	$\tan \alpha = e \tan \theta$, so $\tan \alpha = 1$, $\alpha = 45^\circ$	B1	
	For 2 nd collision $w \cos \beta = v \cos(180 - 60 - \alpha)$ $w \sin \beta = ev \sin(180 - 60 - \alpha)$	M1	Both. May be implied by the A1.
	$\tan \beta = e \tan(120 - \text{their } \alpha)$	M1	Divide to find β .
	$\beta = 61.8^\circ$	A1	
		4	

PUBLISHED

Question	Answer	Marks	Guidance
7(a)	Frictional force = $\mu \times$ normal reaction at D and E	B1	$F_{AB} = \mu R$, $F_{BC} = \mu N$
	Moments about B , $Na - Ra = Wa(\sin \theta - \cos \theta)$ Moments about centre, $F_{AB}a + F_{BC}a = Wa(\cos \theta - \sin \theta)$ Moments about D , $F_{BC}a + Na = Wa(\cos \theta + \sin \theta)$ Moments about E , $Ra - F_{AB}a = Wa(\cos \theta + \sin \theta)$	B1	One moments equation about any point involving all relevant forces, resolved if necessary (AEF).
	Parallel to AB , $N - F_{AB} = W \sin \theta + W \sin \theta$ Perpendicular to AB , $F_{BC} + R = W \cos \theta + W \cos \theta$	B1	Two resolutions: all relevant terms, different frictional forces [Vertical: $R \cos \theta + F_{BC} \cos \theta + N \sin \theta = F_{AB} \sin \theta + W + W$ Horizontal: $F_{BC} \sin \theta + F_{AB} \cos \theta + R \sin \theta = N \cos \theta$] Alternative approach using two moments equations can earn the B1B1
	$N - R = \frac{1}{2}((1 - \mu)N - (1 + \mu)R)$	M1	Combine appropriate equations.
	$N\left(1 - \frac{1}{2}(1 - \mu)\right) = R\left(1 - \frac{1}{2}(1 + \mu)\right)$ $N\left(\frac{1}{2} + \frac{1}{2}\mu\right) = R\left(\frac{1}{2} - \frac{1}{2}\mu\right)$	M1	Collect terms to obtain ratio/fraction in terms of μ only (CWO), any equivalent simplified form.
	$R : N = 1 + \mu : 1 - \mu$	A1	
		6	

Question	Answer	Marks	Guidance
7(b)	Divide resolution equations: $\tan \theta = \frac{N - \mu R}{\mu N + R}$	M1	Must include μ terms.
	Use $R = 2N$: $\tan \theta = \frac{\frac{1}{3}N}{\frac{7}{3}N}$	M1	FT their answer to part (a).
	$\tan \theta = \frac{1}{7}$	A1	
		3	