



**Cambridge Assessment
International Education**

Cambridge International AS & A Level

CANDIDATE
NAME

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--

FURTHER MATHEMATICS

9231/23

Paper 2 Further Pure Mathematics 2

October/November 2023

2 hours

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [].

This document has **16** pages. Any blank pages are indicated.



BLANK PAGE

$$\begin{array}{rcl} 14x - 4y + 6z & = & 5, \\ x + y + kz & = & 3, \\ -21x + 6y - 9z & = & 14. \end{array}$$

where k is a constant, does not have a unique solution and interpret this situation geometrically. [4]

This image shows a full page of white paper with horizontal dotted lines. The lines are evenly spaced and run across the width of the page, providing a guide for handwriting practice. There are no margins, text, or other markings on the page.

[illegible]

[illegible]

$$\frac{d^2y}{dx^2} + 2\frac{dy}{dx} + 3y = 27x^2,$$

given that, when $x = 0$, $y = 2$ and $\frac{dy}{dx} = -8$. [10]

This image shows a full page of white paper with horizontal dotted lines. The lines are evenly spaced and run across the width of the page, providing a guide for handwriting practice. There are no margins, text, or other markings on the page.

(a) Find the exact length of C . [5]

© UCLES 2023

This image shows a full page of white paper with horizontal dashed lines, typical of primary-ruled notebook paper. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

6 (a) Starting from the definitions of cosh and sinh in terms of exponentials, prove that

$\sinh 2x = 2 \sinh x \cosh x.$ [3]

.....

.....

.....

.....

.....

.....

.....

.....

(b) Using the substitution $u = \sinh x$, find $\int \sinh^2 2x \cosh x \, dx.$ [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(c) Find the particular solution of the differential equation

$$\frac{dy}{dx} + y \tanh x = \sinh^2 2x,$$

given that $y = 4$ when $x = 0$. Give your answer in the form $y = f(x)$. [7]

[illegible]

$$\mathbf{A} = \begin{pmatrix} -6 & 2 & 13 \\ 0 & -2 & 5 \\ 0 & 0 & 8 \end{pmatrix}.$$

- (a) Find a matrix $\mathbf{P}$ and a diagonal matrix $\mathbf{D}$ such that $\mathbf{A}^{-1} = \mathbf{P}\mathbf{D}\mathbf{P}^{-1}$. [7]

[illegible]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Use the characteristic equation of **A** to find $\mathbf{A}^{-1}$. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

.....

.....

$$1 + \cos \theta + \cos 2\theta + \dots + \cos(n-1)\theta = \frac{1}{2} \left(1 - \cos n\theta + \frac{\sin n\theta \sin \theta}{1 - \cos \theta} \right). \quad [7]$$



The diagram shows the curve with equation $y = \cos x$ for $0 \leq x \leq 1$, together with a set of n rectangles of width $\frac{1}{n}$.

(c) By considering the sum of the areas of these rectangles, show that

$$\int_0^1 \cos x \, dx < \frac{1}{2n} \left(1 - \cos 1 + \frac{\sin 1 \sin \frac{1}{n}}{1 - \cos \frac{1}{n}} \right).$$

[4]

(d) Use a similar method to find, in terms of n , a lower bound for $\int_0^1 \cos x \, dx$. [3]

[illegible]

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.