GradeMax

Cambridge International AS & A Level

Cambridge Assessment International Education
Cambridge International Advanced Subsidiary and Advanced Level

CANDIDATE NAME

CENTRE NUMBER | | | | | |

CANDIDATE NUMBER | | | | |

6611123410

MATHEMATICS **9709/12**

Paper 1 Pure Mathematics 1 **(P1)** **May/June 2019**

1 hour 45 minutes

Candidates answer on the Question Paper.

Additional Materials: List of Formulae (MF9)

READ THESE INSTRUCTIONS FIRST

Write your centre number, candidate number and name in the spaces at the top of this page.
Write in dark blue or black pen.
You may use an HB pencil for any diagrams or graphs.
Do not use staples, paper clips, glue or correction fluid.
DO **NOT** WRITE IN ANY BARCODES.

Answer **all** the questions in the space provided. If additional space is required, you should use the lined page at the end of this booklet. The question number(s) must be clearly shown.
Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.
The use of an electronic calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.
The number of marks is given in brackets [ ] at the end of each question or part question.
The total number of marks for this paper is 75.

This document consists of **18** printed pages and **2** blank pages.

© UCLES 2019

GradeMax

1 Find the coefficient of x in the expansion of $\left(\frac{2}{x} - 3x\right)^5$. [3]

© UCLES 2019

GradeMax

2 Two points A and B have coordinates $(1, 3)$ and $(9, -1)$ respectively. The perpendicular bisector of AB intersects the y-axis at the point C. Find the coordinates of C. [5]

© UCLES 2019

[Turn over

GradeMax

3 A curve is such that $\frac{dy}{dx} = x^3 - \frac{4}{x^2}$. The point $P(2, 9)$ lies on the curve.

(i) A point moves on the curve in such a way that the x-coordinate is decreasing at a constant rate of 0.05 units per second. Find the rate of change of the y-coordinate when the point is at P. [2]

.....

.....

.....

.....

.....

.....

.....

.....

(ii) Find the equation of the curve. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2019

GradeMax

4 Angle x is such that $\sin x = a + b$ and $\cos x = a - b$, where a and b are constants.

(i) Show that $a^2 + b^2$ has a constant value for all values of x. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(ii) In the case where $\tan x = 2$, express a in terms of b. [2]

.....

.....

.....

.....

.....

.....


.....

.....

.....

.....

5



The diagram shows a semicircle with diameter AB, centre O and radius r. The point C lies on the circumference and angle $AOC = \theta$ radians. The perimeter of sector BOC is twice the perimeter of sector AOC. Find the value of θ correct to 2 significant figures. [5]

GradeMax

6 The equation of a curve is $y = 3\cos 2x$ and the equation of a line is $2y + \frac{3x}{\pi} = 5$.

(i) State the smallest and largest values of y for both the curve and the line for $0 \leqslant x \leqslant 2\pi$. [3]

.....

.....

.....

.....

.....

.....

.....

.....

(ii) Sketch, on the same diagram, the graphs of $y = 3\cos 2x$ and $2y + \frac{3x}{\pi} = 5$ for $0 \leqslant x \leqslant 2\pi$. [3]

(iii) State the number of solutions of the equation $6\cos 2x = 5 - \frac{3x}{\pi}$ for $0 \leqslant x \leqslant 2\pi$. [1]

.....

.....

.....

.....

© UCLES 2019

[Turn over

GradeMax

7 Functions f and g are defined by

$$\mathrm{f} : x \mapsto 3x - 2, \quad x \in \mathbb{R},$$
$$\mathrm{g} : x \mapsto \frac{2x + 3}{x - 1}, \quad x \in \mathbb{R},\ x \neq 1.$$

(i) Obtain expressions for $\mathrm{f}^{-1}(x)$ and $\mathrm{g}^{-1}(x)$, stating the value of x for which $\mathrm{g}^{-1}(x)$ is not defined. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2019

GradeMax

(ii) Solve the equation $\mathrm{fg}(x) = \frac{7}{3}$. [3]

GradeMax

8 The position vectors of points A and B, relative to an origin O, are given by

$$\overrightarrow{OA} = \begin{pmatrix} 6 \\ -2 \\ -6 \end{pmatrix} \quad \text{and} \quad \overrightarrow{OB} = \begin{pmatrix} 3 \\ k \\ -3 \end{pmatrix},$$

where k is a constant.

(i) Find the value of k for which angle AOB is 90°. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(ii) Find the values of k for which the lengths of OA and OB are equal. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2019

The point C is such that $\overrightarrow{AC} = 2\overrightarrow{CB}$.

(iii) In the case where $k = 4$, find the unit vector in the direction of $\overrightarrow{OC}$. [4]

GradeMax

9 The curve C_1 has equation $y = x^2 - 4x + 7$. The curve C_2 has equation $y^2 = 4x + k$, where k is a constant. The tangent to C_1 at the point where $x = 3$ is also the tangent to C_2 at the point P. Find the value of k and the coordinates of P. [8]

GradeMax

GradeMax

10 **(a)** In an arithmetic progression, the sum of the first ten terms is equal to the sum of the next five terms. The first term is a.

(i) Show that the common difference of the progression is $\frac{1}{3}a$. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(ii) Given that the tenth term is 36 more than the fourth term, find the value of a. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

© UCLES 2019

GradeMax

(b) The sum to infinity of a geometric progression is 9 times the sum of the first four terms. Given that the first term is 12, find the value of the fifth term. [4]

11



The diagram shows part of the curve $y = \sqrt{(4x + 1)} + \frac{9}{\sqrt{(4x + 1)}}$ and the minimum point M.

(i) Find expressions for $\frac{dy}{dx}$ and $\int y \, dx$. [6]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2019

GradeMax

(ii) Find the coordinates of M. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

The shaded region is bounded by the curve, the y-axis and the line through M parallel to the x-axis.

(iii) Find, showing all necessary working, the area of the shaded region. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2019

Additional Page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.

GradeMax

BLANK PAGE

© UCLES 2019

GradeMax

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.

© UCLES 2019