GradeMax

Cambridge International AS Level

Cambridge Assessment International Education
Cambridge International Advanced Subsidiary Level

CANDIDATE NAME

CENTRE NUMBER

CANDIDATE NUMBER

8475261242

MATHEMATICS **9709/22**

Paper 2 Pure Mathematics 2 **(P2)** **October/November 2019**

1 hour 15 minutes

Candidates answer on the Question Paper.

Additional Materials: List of Formulae (MF9)

READ THESE INSTRUCTIONS FIRST

Write your centre number, candidate number and name in the spaces at the top of this page.
Write in dark blue or black pen.
You may use an HB pencil for any diagrams or graphs.
Do not use staples, paper clips, glue or correction fluid.
DO **NOT** WRITE IN ANY BARCODES.

Answer **all** the questions in the space provided. If additional space is required, you should use the lined page at the end of this booklet. The question number(s) must be clearly shown.
Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.
The use of an electronic calculator is expected, where appropriate.
You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.
The number of marks is given in brackets [ ] at the end of each question or part question.
The total number of marks for this paper is 50.

This document consists of **12** printed pages.

© UCLES 2019

Cambridge Assessment International Education

GradeMax

1 The polynomial $f(x)$ is defined by

$$f(x) = x^4 - 3x^3 + 5x^2 - 6x + 11.$$

Find the quotient and remainder when $f(x)$ is divided by $(x^2 + 2)$. [3]

© UCLES 2019

GradeMax

2 **(i)** Solve the equation $|4x + 5| = |x - 7|$. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(ii) Hence, using logarithms, solve the equation $\left|2^{y+2} + 5\right| = \left|2^{y} - 7\right|$, giving the answer correct to 3 significant figures. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

© UCLES 2019

[Turn over

3



The variables x and y satisfy the equation $y = kx^a$, where k and a are constants. The graph of $\ln y$ against $\ln x$ is a straight line passing through the points (0.22, 3.96) and (1.32, 2.43), as shown in the diagram. Find the values of k and a correct to 3 significant figures. [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2019

GradeMax

4 The sequence $x_1, x_2, x_3, \ldots$ defined by

$$x_1 = 1, \quad x_{n+1} = \frac{x_n}{\ln(2x_n)}$$

converges to the value α.

(i) Use the iterative formula to find the value of α correct to 4 significant figures. Give the result of each iteration to 6 significant figures. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(ii) State an equation satisfied by α and hence determine the exact value of α. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2019

[Turn over

GradeMax

5 Find the exact coordinates of the stationary point of the curve with equation $y = e^{-\frac{1}{2}x}(2x + 5)$. [5]

6 **(a)** Show that $\int_2^{18} \frac{3}{2x}\,\mathrm{d}x = \ln 27$. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Find the exact value of $\int_0^{\frac{1}{6}\pi} 4\sin^2\left(\frac{3}{2}x\right)\mathrm{d}x$. Show all necessary working. [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

7 The parametric equations of a curve are

$$x = 3\sin 2\theta, \quad y = 1 + 2\tan 2\theta,$$

for $0 \leqslant \theta < \frac{1}{4}\pi$.

(i) Find the exact gradient of the curve at the point for which $\theta = \frac{1}{6}\pi$. [4]

GradeMax

(ii) Find the value of θ at the point where the gradient of the curve is 2, giving the value correct to 3 significant figures. [4]

GradeMax

8 **(i)** Express $0.5\cos\theta - 1.2\sin\theta$ in the form $R\cos(\theta + \alpha)$, where $R > 0$ and $0° < \alpha < 90°$, giving the value of α correct to 2 decimal places. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(ii) Hence solve the equation $0.5\cos\theta - 1.2\sin\theta = 0.8$ for $0° < \theta < 360°$. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2019

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(iii) Determine the greatest and least possible values of $(3 - \cos\theta + 2.4\sin\theta)^2$ as θ varies. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2019

GradeMax

Additional Page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.

© UCLES 2019