GradeMax

Cambridge Assessment International Education

Cambridge International AS & A Level

CANDIDATE NAME									
CENTRE NUMBER						CANDIDATE NUMBER				

6412245051

MATHEMATICS **9709/13**

Paper 1 Pure Mathematics 1 **May/June 2021**

1 hour 50 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [ ].

This document has **20** pages. Any blank pages are indicated.

© UCLES 2021

PapaCambridge

GradeMax

BLANK PAGE

© UCLES 2021

PapaCambridge

GradeMax

1 A curve with equation $y = f(x)$ is such that $f'(x) = 6x^2 - \frac{8}{x^2}$. It is given that the curve passes through the point (2, 7).

Find $f(x)$. [3]

PapaCambridge

GradeMax

2 The function f is defined by $f(x) = \frac{1}{3}(2x - 1)^{\frac{3}{2}} - 2x$ for $\frac{1}{2} < x < a$. It is given that f is a decreasing function.

Find the maximum possible value of the constant a. [4]

PapaCambridge

GradeMax

3 A line with equation $y = mx - 6$ is a tangent to the curve with equation $y = x^2 - 4x + 3$.

Find the possible values of the constant m, and the corresponding coordinates of the points at which the line touches the curve. [6]

PapaCambridge

GradeMax

4 **(a)** Show that the equation

$$\frac{\tan x + \sin x}{\tan x - \sin x} = k,$$

where k is a constant, may be expressed as

$$\frac{1 + \cos x}{1 - \cos x} = k.$$ [2]

.....

.....

.....

.....

.....

.....

.....

(b) Hence express $\cos x$ in terms of k. [2]

.....

.....

.....

.....

.....

.....

(c) Hence solve the equation $\frac{\tan x + \sin x}{\tan x - \sin x} = 4$ for $-\pi < x < \pi$. [2]

.....

.....

.....

.....

.....

.....

© UCLES 2021

PapaCambridge

5



The diagram shows a triangle ABC, in which angle $ABC = 90°$ and $AB = 4$ cm. The sector ABD is part of a circle with centre A. The area of the sector is $10\,\text{cm}^2$.

(a) Find angle BAD in radians. [2]

.....

.....

.....

.....

.....

(b) Find the perimeter of the shaded region. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2021

PapaCambridge

GradeMax

6 Functions f and g are both defined for $x \in \mathbb{R}$ and are given by

$$f(x) = x^2 - 2x + 5,$$
$$g(x) = x^2 + 4x + 13.$$

(a) By first expressing each of $f(x)$ and $g(x)$ in completed square form, express $g(x)$ in the form $f(x + p) + q$, where p and q are constants. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Describe fully the transformation which transforms the graph of $y = f(x)$ to the graph of $y = g(x)$. [2]

.....

.....

.....

.....

.....

.....

.....

© UCLES 2021

PapaCambridge

GradeMax

7 **(a)** Write down the first four terms of the expansion, in ascending powers of x, of $(a - x)^6$. [2]

.....

.....

.....

.....

.....

.....

(b) Given that the coefficient of x^2 in the expansion of $\left(1 + \frac{2}{ax}\right)(a - x)^6$ is -20, find in exact form the possible values of the constant a. [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

PapaCambridge

GradeMax

8 Functions f and g are defined as follows:

$$\text{f} : x \mapsto x^2 - 1 \text{ for } x < 0,$$
$$\text{g} : x \mapsto \frac{1}{2x + 1} \text{ for } x < -\tfrac{1}{2}.$$

(a) Solve the equation $\text{fg}(x) = 3$. [4]

PapaCambridge

GradeMax

(b) Find an expression for $(\mathrm{fg})^{-1}(x)$. [3]

PapaCambridge

GradeMax

9 **(a)** A geometric progression is such that the second term is equal to 24% of the sum to infinity.

Find the possible values of the common ratio. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2021

PapaCambridge

GradeMax

(b) An arithmetic progression P has first term a and common difference d. An arithmetic progression Q has first term $2(a + 1)$ and common difference $(d + 1)$. It is given that

$$\frac{\text{5th term of } P}{\text{12th term of } Q} = \frac{1}{3} \quad \text{and} \quad \frac{\text{Sum of first 5 terms of } P}{\text{Sum of first 5 terms of } Q} = \frac{2}{3}.$$

Find the value of a and the value of d. [6]

PapaCambridge

GradeMax

10 Points $A\,(-2,\ 3)$, $B\,(3,\ 0)$ and $C\,(6,\ 5)$ lie on the circumference of a circle with centre D.

(a) Show that angle $ABC = 90°$. [2]

.....

.....

.....

.....

.....

.....

(b) Hence state the coordinates of D. [1]

.....

.....

.....

(c) Find an equation of the circle. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2021

PapaCambridge

The point E lies on the circumference of the circle such that BE is a diameter.

(d) Find an equation of the tangent to the circle at E. [5]

PapaCambridge

GradeMax

11



The diagram shows part of the curve with equation $y = x^{\frac{1}{2}} + k^2x^{-\frac{1}{2}}$, where k is a positive constant.

(a) Find the coordinates of the minimum point of the curve, giving your answer in terms of k. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2021

PapaCambridge

The tangent at the point on the curve where $x = 4k^2$ intersects the y-axis at P.

(b) Find the y-coordinate of P in terms of k. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

The shaded region is bounded by the curve, the x-axis and the lines $x = \frac{9}{4}k^2$ and $x = 4k^2$.

(c) Find the area of the shaded region in terms of k. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

© UCLES 2021

PapaCambridge

GradeMax

Additional Page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.

PapaCambridge

GradeMax

BLANK PAGE

© UCLES 2021

PapaCambridge

GradeMax

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.

PapaCambridge