



Cambridge International AS & A Level

CANDIDATE
NAME
CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--

MATHEMATICS**9709/42**

Paper 4 Mechanics

February/March 2023**1 hour 15 minutes**

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.
- Where a numerical value for the acceleration due to gravity (g) is needed, use 10 m s^{-2} .

INFORMATION

- The total mark for this paper is 50.
- The number of marks for each question or part question is shown in brackets [].

This document has **16** pages. Any blank pages are indicated.



1 A crate of mass 200 kg is being pulled at constant speed along horizontal ground by a horizontal rope attached to a winch. The winch is working at a constant rate of 4.5 kW and there is a constant resistance to the motion of the crate of magnitude 600 N.

(a) Find the time that it takes for the crate to move a distance of 15 m. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

The rope breaks after the crate has moved 15 m.

(b) Find the time taken, after the rope breaks, for the crate to come to rest. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

3

2 A particle P is projected vertically upwards from horizontal ground with speed 15 m s^{-1} .

(a) Find the speed of P when it is 10 m above the ground. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

At the same instant that P is projected, a second particle Q is dropped from a height of 18 m above the ground in the same vertical line as P .

(b) Find the height above the ground at which the two particles collide. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

3 A particle moves in a straight line starting from rest from a point O . The acceleration of the particle at time t s after leaving O is $a \text{ m s}^{-2}$, where $a = 4t^{\frac{1}{2}}$.

(a) Find the speed of the particle when $t = 9$. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Find the time after leaving O at which the speed (in metres per second) and the distance travelled (in metres) are numerically equal. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

4 A toy railway locomotive of mass 0.8 kg is towing a truck of mass 0.4 kg on a straight horizontal track at a constant speed of 2 m s^{-1} . There is a constant resistance force of magnitude 0.2 N on the locomotive, but no resistance force on the truck. There is a light rigid horizontal coupling connecting the locomotive and the truck.

(a) State the tension in the coupling. [1]

.....

.....

(b) Find the power produced by the locomotive’s engine. [1]

.....

.....

.....

The power produced by the locomotive’s engine is now changed to 1.2 W.

(c) Find the magnitude of the tension in the coupling at the instant that the locomotive begins to accelerate. [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

.....

.....

5




(a) Find the magnitude of the force in each of the struts AD and BD . [3]

[illegible]

-
- This image shows a full page of white paper with horizontal dotted lines. The lines are evenly spaced and run across the width of the page, providing a guide for handwriting practice. There are no margins, text, or other markings on the page.

8

6



A block B , of mass 2 kg , lies on a rough inclined plane sloping at 30° to the horizontal. A light rope, inclined at an angle of 20° above a line of greatest slope, is attached to B . The tension in the rope is $T\text{ N}$. There is a friction force of $F\text{ N}$ acting on B (see diagram). The coefficient of friction between B and the plane is μ .

(a) It is given that $F = 5$ and that the acceleration of B up the plane is 1.2 m s^{-2} .

(i) Find the value of T .

[3]

.....

.....

.....

.....

.....

.....

.....

(ii) Find the value of μ .

[3]

.....

.....

.....

.....

.....

.....

.....

(b) It is given instead that $\mu = 0.8$ and $T = 15$.

Determine whether B will move up the plane. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

.....

.....

7



The diagram shows a smooth track which lies in a vertical plane. The section AB is a quarter circle of radius 1.8 m with centre O . The section BC is a horizontal straight line of length 7.0 m and OB is perpendicular to BC . The section CFE is a straight line inclined at an angle of θ° above the horizontal.

A particle P of mass 0.5 kg is released from rest at A . Particle P collides with a particle Q of mass 0.1 kg which is at rest at B . Immediately after the collision, the speed of P is 4 ms^{-1} in the direction BC . You should assume that P is moving horizontally when it collides with Q .

- (a) Show that the speed of Q immediately after the collision is 10 ms^{-1} . [4]

This image shows a full page of a handwriting practice worksheet. It consists of multiple sets of three horizontal dotted lines, providing a guide for letter height and placement. The lines are evenly spaced across the entire page, which is otherwise blank.

11

When Q reaches C , it collides with a particle R of mass 0.4 kg which is at rest at C . The two particles coalesce. The combined particle comes instantaneously to rest at F . You should assume that there is no instantaneous change in speed as the combined particle leaves C , nor when it passes through C again as it returns down the slope.

- (b) Given that the distance CF is 0.4 m , find the value of θ . [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

[Question 7 continues on the next page.]

[5]

This image shows a full page of white paper with horizontal dashed lines, typical of primary school writing paper. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

Additional Page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.

[illegible]

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.