GradeMax

CAMBRIDGE
International Education

Cambridge International A Level

MATHEMATICS **9709/62**

Paper 6 Probability & Statistics 2 **May/June 2026**

MARK SCHEME

Maximum Mark: 50

Published

This mark scheme contains the instructions and marking guidance that our examiners used to mark candidate responses for this component.

Due to a series-specific issue during the live exam series, all candidates were awarded full marks for question 2. The mark scheme for this question was not used by examiners.

Before they start marking, examiners complete a standardisation process. They review a range of candidate responses and discuss the mark scheme in detail to agree which answers can and cannot be accepted. Examiners award marks where it is clear that candidate responses have met the requirements of the mark scheme. These requirements may vary between different components or different versions of the same component, depending on the exact requirements of the question and the candidate responses seen during the standardisation process.

Sometimes, our mark schemes may allow marks to be awarded to responses which contain elements that are factually incorrect or for content not covered by the syllabus. We do this in response to the demands of a specific question to support candidates and make sure that our assessments are fair. However, these allowances should not be used as a guide for teaching and learning.

We cannot discuss the mark scheme with schools, and we recommend that schools read the mark scheme together with the assessment material and the Principal Examiner Report for Teachers.

This document consists of **18** printed pages.

© Cambridge University Press & Assessment 2026

[Turn over

GradeMax

General marking principles

All examiners must apply these marking principles when marking candidate responses.

1	Examiners must award marks for each response in line with: • the specific **content** defined in the mark scheme • the specific **skills** defined in the mark scheme or in the level descriptions • the **standard of response** required as exemplified by the standardisation scripts.
2	Examiners must award **whole marks** (not half marks, or other fractions).
3	Examiners must award marks **positively**. This means that: • marks are not deducted for errors or omissions • marks are awarded for correct/valid answers as defined in the mark scheme and also for **other valid answers** which go beyond the scope of the syllabus and mark scheme referring to your team leader as appropriate • the quality of spelling, punctuation and grammar are judged **only** when the mark scheme indicates that these features are **specifically assessed** in the question. However, the meaning of all responses must be unambiguous.
4	Examiners must **consistently** apply the requirements of the mark scheme and any rules for what to do when candidates have not followed instructions correctly.
5	Examiners must use the **full range of marks** defined in the mark scheme, unless limited by the quality of the candidate responses seen.
6	Examiners must award marks according to the mark scheme and **must not** award marks with grade thresholds or grade descriptions in mind.

© Cambridge University Press & Assessment 2026

GradeMax

Mathematics-Specific Marking Principles

1 Unless a particular method has been specified in the question, full marks may be awarded for any correct method. However, if a calculation is required then no marks will be awarded for a scale drawing.

2 Unless specified in the question, non-integer answers may be given as fractions, decimals or in standard form. Ignore superfluous zeros, provided that the degree of accuracy is not affected.

3 Allow alternative conventions for notation if used consistently throughout the paper, e.g. commas being used as decimal points.

4 Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored (isw).

5 Where a candidate has misread a number or sign in the question and used that value consistently throughout, provided that number does not alter the difficulty or the method required, award all marks earned and deduct just 1 A or B mark for the misread.

6 Recovery within working is allowed, e.g. a notation error in the working where the following line of working makes the candidate's intent clear.

© Cambridge University Press & Assessment 2026

GradeMax

Annotations guidance for centres

Examiners use a system of annotations as a shorthand for communicating their marking decisions to one another. Examiners are trained during the standardisation process on how and when to use annotations. The purpose of annotations is to inform the standardisation and monitoring processes and guide the supervising examiners when they are checking the work of examiners within their team. The meaning of annotations and how they are used is specific to each component and is understood by all examiners who mark the component.

We publish annotations in our mark schemes to help centres understand the annotations they may see on copies of scripts. Note that there may not be a direct correlation between the number of annotations on a script and the mark awarded. Similarly, the use of an annotation may not be an indication of the quality of the response.

The annotations listed below were available to examiners marking this component in this series.

Annotations

Annotation	Meaning
	More information required
	Accuracy mark awarded zero
	Accuracy mark awarded one
	Independent accuracy mark awarded zero
	Independent accuracy mark awarded one
	Independent accuracy mark awarded two
	Benefit of the doubt
	Blank Page
	Incorrect
Dep	Used to indicate DM0 or DM1

© Cambridge University Press & Assessment 2026

GradeMax

Annotation	Meaning
DM1	Dependent on the previous M1 mark(s)
FT	Follow through
~~	Indicate working that is right or wrong
Highlighter	Highlight a key point in the working
ISW	Ignore subsequent work
J	Judgement
JU	Judgement
M0	Method mark awarded zero
M1	Method mark awarded one
M2	Method mark awarded two
MR	Misread
O	Omission or Other solution
Off-page comment	Allows comments to be entered at the bottom of the RM marking window and then displayed when the associated question item is navigated to.
On-page comment	Allows comments to be entered in speech bubbles on the candidate response.
PE	Judgment made by the PE
Pre	Premature approximation
SC	Special case
SEEN	Indicates that work/page has been seen

© Cambridge University Press & Assessment 2026

9709/62 Mark Scheme June 2026

GradeMax

Annotation	Meaning
SF	Error in number of significant figures
✓	Correct
TE	Transcription error
XP	Correct answer from incorrect working

© Cambridge University Press & Assessment 2026

GradeMax

PUBLISHED

Mark Scheme Notes

The following notes are intended to aid interpretation of mark schemes in general, but individual mark schemes may include marks awarded for specific reasons outside the scope of these notes.

Types of mark

M Method mark, awarded for a valid method applied to the problem. Method marks are not lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. Correct application of a formula without the formula being quoted obviously earns the M mark and in some cases an M mark can be implied from a correct answer.

A Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated method mark is earned (or implied).

B Mark for a correct result or statement independent of method marks.

DM or **DB** When a part of a question has two or more 'method' steps, the M marks are generally independent unless the scheme specifically says otherwise; and similarly, when there are several B marks allocated. The notation DM or DB is used to indicate that a particular M or B mark is dependent on an earlier M or B (asterisked) mark in the scheme. When two or more steps are run together by the candidate, the earlier marks are implied and full credit is given.

FT Implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A or B marks are given for correct work only.

- A or B marks are given for correct work only (not for results obtained from incorrect working) unless follow through is allowed (see abbreviation FT above).
- For a numerical answer, allow the A or B mark if the answer is correct to 3 significant figures or would be correct to 3 significant figures if rounded (1 decimal place for angles in degrees).
- The total number of marks available for each question is shown at the bottom of the Marks column.
- Wrong or missing units in an answer should not result in loss of marks unless the guidance indicates otherwise.
- Square brackets [ ] around text or numbers show extra information not needed for the mark to be awarded.

© Cambridge University Press & Assessment 2026

GradeMax

Abbreviations

AEF/OE	Any Equivalent Form (of answer is equally acceptable) / Or Equivalent
AG	Answer Given on the question paper (so extra checking is needed to ensure that the detailed working leading to the result is valid)
CAO	Correct Answer Only (emphasising that no 'follow through' from a previous error is allowed)
CWO	Correct Working Only
ISW	Ignore Subsequent Working
SOI	Seen Or Implied
SC	Special Case (detailing the mark to be given for a specific wrong solution, or a case where some standard marking practice is to be varied in the light of a particular circumstance)
WWW	Without Wrong Working
AWRT	Answer Which Rounds To

© Cambridge University Press & Assessment 2026

Question	Answer	Marks	Guidance
1	Est(μ) = $[\frac{88.9}{10}] = \frac{889}{100} = 8.89$	**B1**	
	Est(σ^2) = $\frac{10}{9}\left(\frac{794.47}{10} - '8.89'^2\right)$ or $\frac{1}{9}(794.47 - 88.9^2/10)$	**M1**	Use of correct formula with their values for Σx^2 and Σx (may need justification for Σx^2 must not be 7903.21) attempted.
	= 0.461	**A1**	(Biased 0.415 **M0A0**).
		3	

Question	Answer	Marks	Guidance
2	$C - 2(A + B) \sim N(-3, \ldots)$	**B1**	SOI.
	Var($C - 2(A + B)$) = 24 + 4(15 + 8) [= 116]	**M1**	or N(−3, 116) SOI **B1M1**.
	$\frac{0-(-3)}{\sqrt{'116'}}$ [= 0.27854…]	**M1**	
	1 − Φ('0.27854') = 1 – (0.6096 to 0.6099)	**M1**	P($X > 0$) SOI.
	= 0.390 (3 sf)	**A1**	Allow 0.39.
		5	

Question	Answer	Marks	Guidance
3(a)	H_0: Population mean = 2.4 H_1: Population mean > 2.4	**B1**	Not just 'mean'=. Allow μ or λ.
		1	

© Cambridge University Press & Assessment 2026

Question	Answer	Marks	Guidance
3(b)	**EITHER:** P($X \geqslant a \mid X \sim$ Po(2.4)) attempted, where a is any integer 3 or more	**(M1**	Must see Poisson expression.
	P($X \geqslant 5$ or 6 $\mid X \sim$ Po(2.4)) attempted,	**M1**	Must see Poisson expression.
	$[P(X \geqslant 5)] = 1 - e^{-2.4}(1 + 2.4 + \frac{2.4^2}{2!} + \frac{2.4^3}{3!} + \frac{2.4^4}{4!}) = 0.0959$ (>0.05) $= 1 - e^{-2.4}(1 + 2.4 + 2.88 + 2.304 + 1.3824)$ $= 1 - (0.0907 + 0.2177 + 0.2613 + 0.2090 + 0.1254)$		Accept AWRT 0.096.
	$[P(X \geqslant 6)] = 0.0959 - e^{-2.4} \times \frac{2.4^5}{5!}$ (or long method) $= 0.0357$ (<0.05) $= 0.0959 - 0.060196$	**A1**	Accept AWRT 0.036. Both correct. **SC:** both answers correct with no working **M0M0B2** or only P($X \geqslant 6$) = 0.0357 OE seen **M0B1**.
	Smallest value is 6	**B1dep)**	Condone $x \geqslant 6$ Dep on at least P($X \geqslant 6$) = 0.0357 OE.
			Note: Correct answer from incorrect method, using point probabilities, no marks.
	OR: P($X \leqslant b \mid$ X ~ Po(2,4)) attempted, where b is any integer 2 or more	**(M1**	Must see Poisson expression.
	OR P($X \leqslant 4$ or 5 $\mid X \sim$ Po(2,4)) attempted	**M1**	Must see Poisson expression.
	OR $[P(X \leqslant 4] = e^{-2.4}(1 + 2.4 + \frac{2.4^2}{2!} + \frac{2.4^3}{3!} + \frac{2.4^4}{4!}) = 0.9041$ (<0.95)		Accept AWRT 0.90.
	OR $[P(X \leqslant 5] = 0.9041 + e^{-2.4} \times \frac{2.4^5}{5!} = 0.9643$ (>0.95)	**A1**	Accept AWRT 0.964. Both correct. **SC:** both answers correct with no working **M0M0B2** or only P($X \leqslant 5$) = 0.9643 OE seen **M0B1**.

© Cambridge University Press & Assessment 2026

GradeMax

Question	Answer	Marks	Guidance
3(b)	Smallest value is 6	**B1dep)**	Condone $x \geqslant 6$ Dep on at least $P(X \leqslant 5) = 0.9643$ OE seen.
			Note: Correct answer from incorrect method, using point probabilities, no marks.
		4	
3(c)	0.0357(3.s.f) Accept 0.036 (2.s.f)	**B1FT**	**FT** *their* answer to **3(b)** must be < 0.05 and no **FT** for point probabilities.
		1	

Question	Answer	Marks	Guidance
4(a)	$[D + E \sim \text{Po}](3.8)$	**B1**	SOI.
	$P(D + E < 4) = e^{-3.8}(1 + 3.8 + \frac{3.8^2}{2!} + \frac{3.8^3}{3!}) = e^{-3.8}(1 + 3.8 + 7.22 + 9.1453) =$ $0.02237 + 0.08501 + 0.1615 + 0.2046$	**M1**	Must see expression (accept fully correct Σ notation). Accept any λ allow one end error.
	[= 0.47348] = 0.473 (3 sf) accept 0.474	**A1**	**SC**: Unjustified correct answer scores **B1B1**.
		3	

© Cambridge University Press & Assessment 2026

Question	Answer	Marks	Guidance
4(b)	$P(D = 1) \times P(E = 0, 1, 2) = e^{-1.7} \times 1.7 \times e^{-2.1} (1 + 2.1 + \frac{2.1^2}{2})$ $= 0.31056 \times 0.64963$ or $P(1, 0) + P(1, 1) + P(1, 2)$ $= (e^{-1.7} \times 1.7 \times e^{-2.1}) + (e^{-1.7} \times 1.7 \times e^{-2.1} \times 2.1) + (e^{-1.7} \times 1.7 \times e^{-2.1} \times \frac{2.1^2}{2})$ $=$ $0.03803 + 0.079864 + 0.083856$ [=0.20175]	**M1**	Attempted (allow with one missing term or one extra term). Need to see Poisson expression.
	$\frac{P(D=1)\times P(E=0,1,2)}{P(D+E<4)} = \frac{e^{-1.7} \times 1.7 \times e^{-2.1}(1+2.1+\frac{2.1^2}{2})}{\textit{Their } 0.47348}$ $[=\frac{0.20175}{0.47348}]$	**M1**	For attempt at their numerator divided by their 0.47348. The numerator must involve $P(D=1) = e^{-1.7} \times 1.7$ and Po(2.1).
	= 0.426 (3 sf) accept 0.427	**A1**	**SC**: correct answer not justified scores **B2.**
		3	

Question	Answer	Marks	Guidance
4(c)	$T \sim \text{N}(170, 170)$	**B1**	SOI
	$\frac{190.5-170}{\sqrt{170}}$ [= 1.572]	**M1**	For standardising. Allow no or wrong cc.
	$1 - \Phi(\text{'1.572'})$	**M1**	For area consistent with their working.
	= 0.0580 Allow 0.058 Allow 0.0579	**A1**	(Note alternative method $\text{N}(1.7, \frac{1.7}{100})$ **B1**: Use of $\frac{1.9+\frac{1}{200}-1.7}{\sqrt{\frac{1.7}{100}}}$, Ans 0.0580 **M1M1A1**. No mixed methods. Use of $(1.9-1.7)/\sqrt{(1.7/100)}$ $\frac{1.9-1.7}{\sqrt{\frac{1.7}{100}}}$, Ans 0.0625 **M1M1A1**).
		4	

© Cambridge University Press & Assessment 2026

GradeMax

Question	Answer	Marks	Guidance
5(a)	$2 \times 1.96 \times \sqrt{\frac{x(100-x)}{100^3}} = 0.165$ or in terms of p (or $\frac{x}{100}$)	**M1**	Condone missing factor of 2, any z.
	1.96 seen	**B1**	
	$x^2 - 100x + 1771.72 = 0$ or $p^2 - p + 0.177172 = 0$	**M1**	OE. Rearrange to form 3-term quadratic equation in x or p.
	= 77 or 23	**A1**	Must be × (0.77 or 0.23 final answer scores **A0**). Must be whole number. Condone 77.0, 23.0.
		4	
5(b)	$2 \times y(1 - y) = 0.1128$	**M1**	Allow omission of '2'. Allow any letter, including 'p'
	$y^2 - y + 0.0564 = 0$ [$y = 0.94$ [or 0.06]]	**M1**	Rearrange equation of correct form to form 3-term quadratic.
	$\alpha = 94$	**A1**	Allow 94%.
		3	

© Cambridge University Press & Assessment 2026

9709/62 Mark Scheme June 2026

GradeMax

Question	Answer	Marks	Guidance
6(a)	H_0: $\mu = 12.4$ H_1: $\mu < 12.4$	**B1**	Or Population mean (not just 'mean').
	$\dfrac{11.9-12.4}{\sqrt{\frac{9.2}{75}}}$	**M1**	Must have $\sqrt{75}$.
	$= -1.428$ or -1.427 OR 0.0767 or 0.0768 or 0.0766 (3sf)	**A1**	Accept 3sf if nothing better seen.
	$-1.428 > -1.645$ or $1.428 < 1.645$ or $0.0767 > 0.05$	**M1**	Valid comparison.
	Insufficient evidence to suggest that the (mean) time (spent by customers) has decreased Insufficient evidence to suggest that $\mu < 12.4$ [Do not reject H_0]	**A1FT**	OE. In context, not definite, no contradictions. Need 'decreased' OE, not 'changed'. Condone 'prove' or similar. Note: Accept critical value method: Critical value = 11.82 **M1A1,** 11.82 < 11.9 **M1,** conclusion **A1**. Condone critical value =12.47 **M1A1,** 12.47 > 12.4 **M1** conclusion **A1**. Two tail test scores **B0M1A1M1** (comparison with 1.96 or 0.025 **A0** (max 3/5).
		5	
6(b)	No because population normal	**B1**	OE, need both ('no' and reason). Accept population/underlying distribution/parent distribution/T, not $\overline{T}$ and not 'time spent in shop' not data.
		1	

© Cambridge University Press & Assessment 2026

Question	Answer	Marks	Guidance
6(c)	$\dfrac{11.8-12.4}{\sqrt{\frac{9.2}{n}}} < -1.645$	**M1**	OE. Allow any z-value. Signs must be consistent. Accept any inequality sign or $=$.
	$\sqrt{n} > 1.645 \div 0.6 \times \sqrt{9.2}$ or $n > 1.645^2 \div 0.6^2 \times 9.2$	**M1**	OE. Accept any inequality sign or $=$. Attempt to rearrange to make $\sqrt{n}$ or n the subject. Dep z-value ± 1.645.
	$[n > 69.2]$ $\quad n \geqslant 70$	**A1**	CWO. Including correct inequality signs (n >70 **A0**).
		3	

© Cambridge University Press & Assessment 2026

GradeMax

Question	Answer	Marks	Guidance
7(a)	Area = $\frac{1}{2} \times a \times a$ or $\int_0^a (a-x)\,\mathrm{d}x$	**M1**	For expression for area or attempt to integrate f(x) with correct limits.
	'$\frac{1}{2}a^2$' = 1	**M1**	Equate *their* ∫f(x) to 1 and attempt to solve.
	$a = \sqrt{2}$	**A1**	
	$\mathrm{E}(X) = \int_0^{\sqrt{2}} (\sqrt{2}x - x^2)\,\mathrm{d}x$	**M1**	OE in terms of '*a*'/*their* '*a*'. Attempt to integrate xf(x), correct limits; accept in terms of *a*/ **FT** *their a.*
	$= \left[\sqrt{2}\frac{x^2}{2} - \frac{x^3}{3}\right]_0^{\sqrt{2}}$	**A1FT**	OE in terms of '*a*'/*their* '*a*'. Correct integration $[\frac{ax^2}{2} - \frac{x^3}{3}]$ limits 0 to *a*, **FT** *their a* [Note: E(X) = $\frac{a^3}{6}$ **M1A1**].
	$[= \sqrt{2} - \frac{2}{3}\sqrt{2}\,] = \frac{1}{3}\sqrt{2}$	**A1**	Correctly obtained. **SC**: correct answer not fully justified (integration not seen) scores **B1** for E(X) **B1** for $\frac{1}{3}$.
		6	

© Cambridge University Press & Assessment 2026

GradeMax

Question	Answer	Marks	Guidance
7(b)	$\frac{1}{2} \times (\sqrt{2} - m) \times (\sqrt{2} - m) = \frac{1}{2}$ or $\int_0^m (\sqrt{2} - x)\mathrm{d}x = \frac{1}{2}$ or $\int_m^{\sqrt{2}} (\sqrt{2} - x)\mathrm{d}x = \frac{1}{2}$ Or in terms of a $\int(a\text{-}x)\mathrm{d}x$ limits 0 and m	**M1**	Attempt area from 0 to m or from m to a. **FT** *their* $\sqrt{2}$ (accept in terms of a) and equate to $\frac{1}{2}$.
	$m^2 - 2\sqrt{2}\,m + 1 = 0 \quad (m^2 - 2am + 1 = 0)$ $[m = \frac{2\sqrt{2} \pm \sqrt{8-4}}{2} = \sqrt{2} \pm 1]$	**A1ft**	Correct 3-term quadratic equation, **FT** *their* $\sqrt{2}$. (accept in terms of a).
	$\sqrt{2} + 1$ is not in the domain of X	**B1**	OE $\sqrt{2} + 1 > \sqrt{2}$. Reject $m = \sqrt{2} + 1$ with reason. (accept in terms of a i.e. reject $a + \sqrt{a^2 - 1}$ as not in domain ($> a$)).
	$m = \sqrt{2} - 1$	**A1**	Must be exact. **SC**: If **B0A0** scored and final answer given as $m = \sqrt{2} \pm 1$ allow **B1**.
		4	

© Cambridge University Press & Assessment 2026