



Cambridge Assessment
International Education

Cambridge IGCSE™

CANDIDATE
NAME

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--

ADDITIONAL MATHEMATICS

0606/21

Paper 2

October/November 2023

2 hours

You must answer on the question paper.

No additional materials are needed.


INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 80.
- The number of marks for each question or part question is shown in brackets [].

This document has **16** pages. Any blank pages are indicated.



Mathematical Formulae**1. ALGEBRA***Quadratic Equation*

For the equation $ax^2 + bx + c = 0$,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Binomial Theorem

$$(a + b)^n = a^n + \binom{n}{1}a^{n-1}b + \binom{n}{2}a^{n-2}b^2 + \dots + \binom{n}{r}a^{n-r}b^r + \dots + b^n$$

where n is a positive integer and $\binom{n}{r} = \frac{n!}{(n-r)!r!}$

Arithmetic series $u_n = a + (n-1)d$

$$S_n = \frac{1}{2}n(a + l) = \frac{1}{2}n\{2a + (n-1)d\}$$

Geometric series $u_n = ar^{n-1}$

$$S_n = \frac{a(1-r^n)}{1-r} \quad (r \neq 1)$$

$$S_\infty = \frac{a}{1-r} \quad (|r| < 1)$$

2. TRIGONOMETRY*Identities*

$$\begin{aligned}\sin^2 A + \cos^2 A &= 1 \\ \sec^2 A &= 1 + \tan^2 A \\ \operatorname{cosec}^2 A &= 1 + \cot^2 A\end{aligned}$$

Formulae for $\triangle ABC$

$$\begin{aligned}\frac{a}{\sin A} &= \frac{b}{\sin B} = \frac{c}{\sin C} \\ a^2 &= b^2 + c^2 - 2bc \cos A \\ \Delta &= \frac{1}{2}bc \sin A\end{aligned}$$

3

1 (a) Write $19 - 12x - 3x^2$ in the form $a(x + b)^2 + c$ where a , b and c are integers. [4]

(b) Hence find the maximum value of $19 - 12x - 3x^2$ and the value of x at which this maximum occurs. [2]

(c) Use your answer to **part (a)** to solve the equation $19 - 12\sqrt{u} - 3u = 0$. [3]

4**2** Solve the following simultaneous equations.

$$5x - 3 \ln y = 2$$

$$x + \ln y = 1$$

[4]

5

3 (a) Find $\int \left(4x + 5 - \frac{1}{2x+3}\right) dx$. [3]

(b) Hence find the exact value of $\int_1^3 \left(4x + 5 - \frac{1}{2x+3}\right) dx$, simplifying your answer. [3]

6

4 In this question a and b are integers.

Three terms in the expansion of $(2 + ax)^5(1 + bx)$ are $32 + 112x - 240x^2$. Find the values of a and b .
[7]

5 In this question p and q are constants.

The normal to the curve $y = \frac{p}{x^2} + 5x - 2$, at the point where $x = 1$, has equation $y = -x + q$.

Find the values of p and q .

[6]

8

- 6** Find the value of the constant a for which the line $y = (2a + 1)x - 10$ is a tangent to the curve $y = ax^2 - 5x + 2$. [6]

9

- 7** A particle moves in a straight line. At time t seconds after passing through a fixed point O , its velocity, $v \text{ ms}^{-1}$, is given by $v = 10 \sin 2t - 6 \cos 2t$.

(a) Find an expression for the acceleration of the particle. [2]

(b) Find the acceleration when $t = \frac{\pi}{4}$. [1]

(c) Find the first time at which the acceleration is zero. [3]

(d) Find the displacement of the particle between $t = \frac{\pi}{4}$ and $t = \frac{\pi}{2}$. [4]

10**8 DO NOT USE A CALCULATOR IN THIS QUESTION.**

Solve the equation $(2 - \sqrt{10})x^2 + x + (2 + \sqrt{10}) = 0$, giving your answers in the form $a + b\sqrt{10}$,
where a and b are rational. [7]

11

- 9** The functions f and g are defined as follows, for all real values of x .

$$f(x) = 2x^2 - 1$$

$$g(x) = e^x + 1$$

- (a)** Solve the equation $fg(x) = 8$.


[3]

- (b)** For each of the functions f and g , either explain why the inverse function does not exist or find the inverse function, stating its domain.

[4]

12

10 In this question all lengths are in centimetres.



The diagram shows a circle centre O with radius 6 . The line AB is a tangent to the circle at the point B . The point C lies on the circle such that AOC is a straight line. $AB = 8$.

(a) Find the perimeter of the shaded region.

[6]

13**(b)** Find the area of the shaded region.**[3]**

14

11 (a) Show that $\frac{1}{\sec x - \operatorname{cosec} x} + \frac{1}{\sec x + \operatorname{cosec} x} = \frac{2 \cos x}{1 - \cot^2 x}$. [5]

15

- (b)** Solve the equation $3 \tan^2(y + \frac{\pi}{4}) = 1$ for $-2\pi < y < 0$. [4]

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.