



Cambridge IGCSE™

CANDIDATE
NAME
CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--

CHEMISTRY**0620/43**

Paper 4 Theory (Extended)

May/June 2023**1 hour 15 minutes**

You must answer on the question paper.

No additional materials are needed.

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- You may use a calculator.
- You should show all your working and use appropriate units.







INFORMATION

- The total mark for this paper is 80.
- The number of marks for each question or part question is shown in brackets [].
- The Periodic Table is printed in the question paper.

This document has **16** pages.

2

1 Some symbol equations and word equations, **A** to **J**, are shown.



Use the equations to answer the questions that follow.

Each equation may be used once, more than once, or not at all.

Give the letter, **A** to **J**, for the equation which represents:

(a) photosynthesis [1]

(b) an addition reaction [1]

(c) a precipitation reaction [1]

(d) incomplete combustion [1]

(e) a displacement reaction [1]

(f) a substitution reaction. [1]

[Total: 6]

3

Question 2 starts on the next page.

- 2 (a) The symbols of the elements in Period 3 of the Periodic Table are shown.

Na Mg Al Si P S Cl Ar

Use the symbols of the elements in Period 3 to answer the questions that follow.
Each symbol may be used once, more than once, or not at all.

Give the symbol of the element that:

- (i) is present in purified bauxite [1]
- (ii) contains atoms with a full outer shell of electrons [1]
- (iii) is used to kill microbes in water treatment [1]
- (iv) forms an amphoteric oxide [1]
- (v) forms an oxide which causes acid rain [1]
- (vi) has an oxidation number of -1 when it forms a compound with hydrogen.
..... [1]
- (b) The relative atomic masses of elements can be calculated from the relative masses of isotopes and their percentage abundances.
- (i) Identify the isotope to which all relative masses are compared.
..... [1]
- (ii) Table 2.1 shows the relative masses and the percentage abundances of the two isotopes in a sample of magnesium.

Table 2.1

relative mass of isotope	percentage abundance of isotope
24	85
26	15

Calculate the relative atomic mass of magnesium to **one** decimal place.

relative atomic mass = [2]

5

(c) An ion contains 10 electrons, 13 protons and 14 neutrons.

(i) State the nucleon number of the ion.

..... [1]

(ii) Identify the element that forms this ion.

..... [1]

[Total: 11]

3 Magnesium forms ionic compounds.

- (a)** Magnesium reacts with fluorine to form the ionic compound magnesium fluoride. The electronic configurations of an atom of magnesium and an atom of fluorine are shown in Fig. 3.1.




Fig. 3.1

- (i)** Ions are formed by the transfer of electrons from magnesium atoms to fluorine atoms.

Complete the dot-and-cross diagrams in Fig. 3.2 to show the electronic configurations of **one** magnesium ion and **one** fluoride ion. Show the charges on the ions.




Fig. 3.2

[3]

- (ii)** Deduce the formula of magnesium fluoride.

..... [1]

- (iii)** When solid magnesium fluoride is dissolved in water it forms a solution that conducts electricity.

State one other change that can be made to solid magnesium fluoride to allow it to conduct electricity.

..... [1]

7

- (b) Silicon tetrachloride, SiCl_4 , and silicon(IV) oxide, SiO_2 , are covalent compounds.

Complete the dot-and-cross diagram in Fig. 3.3 to show the electronic configuration in a molecule of silicon tetrachloride. Show outer shell electrons only.




Fig. 3.3

[2]

- (c) The melting points of silicon tetrachloride and silicon(IV) oxide are shown in Table 3.1.

Table 3.1

	melting point/ $^{\circ}\text{C}$
silicon tetrachloride	–69
silicon(IV) oxide	1710

- (i) Silicon tetrachloride has a low melting point because it has weak forces of attraction between particles.

Name the type of particles that are held together by these weak forces of attraction.

..... [1]


- (ii) Explain, in terms of structure and bonding, why silicon(IV) oxide has a high melting point.

.....

 [2]

[Total: 10]

- 4 Hydrogen is produced by the reaction between zinc and dilute sulfuric acid, H_2SO_4 .



- (a) A student carries out an experiment using excess zinc and dilute sulfuric acid.

The student measures the volume of hydrogen produced at regular time intervals using the apparatus shown in Fig. 4.1.

Lumps of zinc are used.




Fig. 4.1

The rate of reaction decreases as the reaction progresses. The rate eventually becomes zero.

- (i) Explain why the rate of reaction decreases as the reaction progresses.

.....
..... [1]

- (ii) Explain why the rate of reaction eventually becomes zero.

.....
..... [1]


- (b) The experiment is repeated using powdered zinc instead of lumps of zinc.
All other conditions remain the same.

Explain, in terms of collision theory, why the rate of reaction increases if powdered zinc is used.

.....
.....
..... [2]

9

(c) The equation for the reaction is shown.



25.0 cm³ of 2.00 mol/dm³ H₂SO₄(aq) is added to excess zinc.

Calculate the volume of H₂ formed at room temperature and pressure (r.t.p.). The volume of one mole of any gas is 24 dm³ at r.t.p.

Use the following steps.

- Calculate the number of moles of H₂SO₄ used.

..... mol

- Deduce the number of moles of H₂ produced.

..... mol

- Calculate the volume of H₂ formed at r.t.p.

..... dm³
[3]

(d) Hydrogen can also be produced by the reaction of zinc with dilute hydrochloric acid.

- (i) Write a symbol equation for this reaction.

..... [2]

- (ii) State the test for hydrogen gas.

test

positive result
[1]

[Total: 10]

5 This question is about electricity and chemical reactions.

(a) Aqueous copper(II) sulfate is an electrolyte.

The electrolysis of aqueous copper(II) sulfate using inert electrodes forms:

- copper at the cathode
- oxygen at the anode.

(i) State what is meant by the term electrolyte.

.....
..... [2]

(ii) State the term given to the Roman numeral, (II), in the name copper(II) sulfate.

..... [1]

(iii) State what happens to the colour of the aqueous copper(II) sulfate as this electrolysis progresses.

..... [1]

(iv) Write an ionic half-equation for the formation of copper at the cathode.

..... [2]

(v) Give the formula of the ion that forms oxygen at the anode.

..... [1]

(b) The electrolysis of aqueous copper(II) sulfate is repeated using **copper** electrodes.

State what happens to the anode.

..... [1]

(c) Spoons can be electroplated with silver.

(i) Name the substances used as:

the anode (positive electrode)

the cathode (negative electrode)

the electrolyte. [3]

(ii) State **two** reasons why spoons are electroplated.

1

2 [2]

11

- (d) Hydrogen–oxygen fuel cells can be used to produce electricity to power cars.
Petrol produces carbon dioxide and carbon monoxide when it powers cars.

- (i) State **one** adverse effect of carbon dioxide and carbon monoxide.

carbon dioxide

carbon monoxide

[2]


- (ii) State **one** disadvantage, other than cost, of using hydrogen–oxygen fuel cells to power cars compared to using petrol.

..... [1]

[Total: 16]

6 This question is about nitrogen and compounds of nitrogen.

- (a) Ammonia is manufactured by the reaction between nitrogen and hydrogen in the Haber process.
The equation is shown.



- (i) State the source of nitrogen for the Haber process.

..... [1]

- (ii) State the source of hydrogen for the Haber process.

..... [1]

- (iii) State the typical conditions used in the Haber process.

temperature °C

pressure atm

[2]

- (iv) Name the catalyst used in the Haber process.

..... [1]


- (v) State what is meant by the term catalyst.

.....

..... [2]

- (b) Ammonia is converted into nitric acid.

- (i) The first stage is the conversion of ammonia into nitrogen monoxide, NO.
The equation is shown.



The reaction is carried out at a temperature of 900 °C and a pressure of 7 atm.
The forward reaction is exothermic.

Using explanations that do **not** involve cost:

- explain why a temperature less than 900 °C is **not** used

.....

- explain why a pressure greater than 7 atm is **not** used.


.....

[2]

13

- (ii) In the second stage, nitrogen monoxide reacts with water and oxygen to produce nitric acid.

Balance the symbol equation for the reaction.



- (c) A student makes aqueous copper(II) nitrate by adding an excess of solid copper(II) carbonate to dilute nitric acid.

- (i) Write the symbol equation for this reaction.

..... [2]

- (ii) State **two** observations that indicate the copper(II) carbonate is in excess.

1

2 [2]

- (iii) Name **one** compound, other than copper(II) carbonate, that can be added to dilute nitric acid to produce aqueous copper(II) nitrate.


..... [1]

[Total: 15]

14

7 This question is about organic compounds.

(a) Propane and chlorine react at room temperature. An equation for the reaction is shown.



(i) State the condition required for this reaction.

..... [1]

(ii) Draw the displayed formulae of **two** structural isomers with the formula $\text{C}_3\text{H}_7\text{Cl}$.

[2]

(b) Alkenes are a homologous series of hydrocarbons.

(i) State **two** characteristics that all members of the same homologous series have in common.

1

2

[2]

(ii) Addition polymers are made from alkenes.

Complete Fig. 7.1 to show one repeat unit of the addition polymer formed from but-2-ene.




Fig. 7.1

[2]

15

- (c) A repeat unit of a condensation polymer is shown in Fig. 7.2.
The polymer is made from two monomers.






Fig. 7.2

- (i) Draw the structures of the monomers used to produce the polymer in Fig. 7.2.



[2]

- (ii) Name the **type** of condensation polymer in Fig. 7.2.

..... [1]

- (iii) Name the **two** homologous series to which the monomers in (i) belong.

1

2

[2]

[Total: 12]

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.

The Periodic Table of Elements

Group																		
I	II											III	IV	V	VI	VII	VIII	
		1Hhydrogen1																
		Key atomic number atomic symbol name relative atomic mass																
3Li lithium 7	4Be beryllium 9											5B boron 11	6C carbon 12	7N nitrogen 14	8O oxygen 16	9F fluorine 19	10Ne neon 20	
11Na sodium 23	12Mg magnesium 24											13Al aluminium 27	14Si silicon 28	15P phosphorus 31	16S sulfur 32	17Cl chlorine 35.5	18Ar argon 40	
19K potassium 39	20Ca calcium 40	21Sc scandium 45	22Ti titanium 48	23V vanadium 51	24Cr chromium 52	25Mn manganese 55	26Fe iron 56	27Co cobalt 59	28Ni nickel 59	29Cu copper 64	30Zn zinc 65	31Ga gallium 70	32Ge germanium 73	33As arsenic 75	34Se selenium 79	35Br bromine 80	36Kr krypton 84	
37Rb rubidium 85	38Sr strontium 88	39Y yttrium 89	40Zr zirconium 91	41Nb niobium 93	42Mo molybdenum 96	43Tc technetium —	44Ru ruthenium 101	45Rh rhodium 103	46Pd palladium 106	47Ag silver 108	48Cd cadmium 112	49In indium 115	50Sn tin 119	51Sb antimony 122	52Te tellurium 128	53I iodine 127	54Xe xenon 131	
55Cs caesium 133	56Ba barium 137	lanthanoids		73Ta tantalum 181	74W tungsten 184	75Re rhenium 186	76Os osmium 190	77Ir iridium 192	78Pt platinum 195	79Au gold 197	80Hg mercury 201	81Tl thallium 204	82Pb lead 207	83Bi bismuth 209	84Po polonium —	85At astatine —	86Rn radon —	
87Fr francium —	88Ra radium —	89–103 actinoids	104Rf rutherfordium —	105Db dubnium —	106Sg seaborgium —	107Bh bohrium —	108Hs hassium —	109Mt meitnerium —	110Ds darmstadtium —	111Rg roentgenium —	112Cn copernicium —	113Nh nihonium —	114Fl flerovium —	115Mc moscovium —	116Lv livermorium —	117Ts tennessine —	118Og oganeson —	

lanthanoids

actinoids

The volume of one mole of any gas is 24 dm³ at room temperature and pressure (r.t.p.).