

Please check the examination details below before entering your candidate information

Candidatesurname					Other names									
<b>Pearson Edexcel</b> <b>International</b> <b>Advanced Level</b>					Centre Number					Candidate Number				
					<input type="text"/>					<input type="text"/>				
Time 1 hour 30 minutes					Paper reference					<b>WME01/01</b>				
<b>Mathematics</b>														
<b>International Advanced Subsidiary/Advanced Level</b>														
<b>Mechanics M1</b>														
<b>You must have:</b> Mathematical Formulae and Statistical Tables (Yellow), calculator										Total Marks				

**Candidates may use any calculator permitted by Pearson regulations. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.**

### Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided  
– *there may be more space than you need.*
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Whenever a numerical value of  $g$  is required, take  $g = 9.8 \text{ m s}^{-2}$ , and give your answer to either 2 significant figures or 3 significant figures.

### Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- There are 8 questions in this question paper. The total mark for this paper is 75.
- The marks for each question are shown in brackets  
– *use this as a guide as to how much time to spend on each question.*

### Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.
- If you change your mind about an answer, cross it out and put your new answer and any working underneath.
- Good luck with your examination.

Turn over ►















Leave  
blank

3. [In this question  $\mathbf{i}$  and  $\mathbf{j}$  are perpendicular horizontal unit vectors.]

Three forces,  $\mathbf{F}_1$ ,  $\mathbf{F}_2$  and  $\mathbf{F}_3$ , are given by

$$\mathbf{F}_1 = (5\mathbf{i} + 2\mathbf{j})\text{N} \quad \mathbf{F}_2 = (-3\mathbf{i} + \mathbf{j})\text{N} \quad \mathbf{F}_3 = (a\mathbf{i} + b\mathbf{j})\text{N}$$

where  $a$  and  $b$  are constants.

The forces  $\mathbf{F}_1$ ,  $\mathbf{F}_2$  and  $\mathbf{F}_3$  act on a particle  $P$  of mass 4 kg.

Given that  $P$  rests in equilibrium on a smooth horizontal surface under the action of these three forces,

- (a) find the size of the angle between the direction of  $\mathbf{F}_3$  and the direction of  $-\mathbf{j}$ . (4)

The force  $\mathbf{F}_3$  is now removed and replaced by the force  $\mathbf{F}_4$  given by  $\mathbf{F}_4 = \lambda(\mathbf{i} + 3\mathbf{j})\text{N}$ , where  $\lambda$  is a positive constant.

When the three forces  $\mathbf{F}_1$ ,  $\mathbf{F}_2$  and  $\mathbf{F}_4$  act on  $P$ , the acceleration of  $P$  has magnitude  $3.25 \text{ m s}^{-2}$

- (b) Find the value of  $\lambda$ . (5)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA










Leave blank

4.



**Figure 1**

Figure 1 shows a large bucket used by a crane on a building site to move materials between the ground and the top of the building. The mass of the bucket is 15 kg.

The bucket is attached to a vertical cable with the bottom of the bucket horizontal. The cable is modelled as light and inextensible.

When the bucket is on the ground, a bag of cement of mass 25 kg is placed in the bucket.

The bucket with the bag of cement moves vertically upwards with constant acceleration  $0.2 \text{ m s}^{-2}$ . Air resistance is modelled as being negligible.

(a) Find the tension in the cable. (3)

At the top of the building, the bag of cement is removed. A box of tools of mass 12 kg is now placed in the bucket.

Later on the bucket with the box of tools is moving vertically downwards with constant deceleration  $0.1 \text{ m s}^{-2}$ . Air resistance is again modelled as being negligible.

(b) Find the magnitude of the normal reaction between the bucket and the box of tools. (3)

---

---

---

---

---

---

---

---

---

---

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA









Leave  
blank

5. [In this question  $\mathbf{i}$  and  $\mathbf{j}$  are perpendicular horizontal unit vectors.]

A particle  $P$  is moving with constant acceleration. At 2pm, the velocity of  $P$  is  $(3\mathbf{i} + 5\mathbf{j}) \text{ km h}^{-1}$  and at 2.30pm the velocity of  $P$  is  $(\mathbf{i} + 7\mathbf{j}) \text{ km h}^{-1}$

At time  $T$  hours after 2pm,  $P$  is moving in the direction of the vector  $(-\mathbf{i} + 2\mathbf{j})$

- (a) Find the value of  $T$ .

(6)

Another particle,  $Q$ , has velocity  $\mathbf{v}_Q \text{ km h}^{-1}$  at time  $t$  hours after 2pm, where

$$\mathbf{v}_Q = (-4 - 2t)\mathbf{i} + (\mu + 3t)\mathbf{j}$$

and  $\mu$  is a constant.

Given that there is an instant when the velocity of  $P$  is equal to the velocity of  $Q$ ,

- (b) find the value of  $\mu$ .

(3)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

















Leave blank

7.




Figure 2

A non-uniform beam  $AB$ , of mass  $60$  kg and length  $8a$  metres, rests in equilibrium in a horizontal position on two vertical supports. One support is at  $C$ , where  $AC = a$  metres and the other support is at  $D$ , where  $DB = 2a$  metres, as shown in Figure 2.

The magnitude of the normal reaction between the beam and the support at  $D$  is three times the magnitude of the normal reaction between the beam and the support at  $C$ .

By modelling the beam as a non-uniform rod whose centre of mass is at a distance  $x$  metres from  $A$ ,

- (a) find an expression for  $x$  in terms of  $a$ . (5)

A box of mass  $M$  kg is placed on the beam at  $E$ , where  $AE = 2a$  metres.

The beam remains in equilibrium in a horizontal position.

The magnitude of the normal reaction between the beam and the support at  $C$  is now equal to the magnitude of the normal reaction between the beam and the support at  $D$ .

By modelling the box as a particle,

- (b) find the value of  $M$ . (5)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA















