

Please check the examination details below before entering your candidate information

Candidate surname					Other names				
Centre Number					Candidate Number				

Pearson Edexcel International Advanced Level

Friday 11 October 2024

Afternoon (Time: 1 hour 30 minutes) **Paper reference** **WME01/01**

Mathematics

International Advanced Subsidiary/Advanced Level

Mechanics M1

You must have:
Mathematical Formulae and Statistical Tables (Yellow), calculator

Total Marks

Candidates may use any calculator permitted by Pearson regulations. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
– *there may be more space than you need.*
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Whenever a numerical value of g is required, take $g = 9.8 \text{ m s}^{-2}$, and give your answer to either 2 significant figures or 3 significant figures.

Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- There are 7 questions in this question paper. The total mark for this paper is 75.
- The marks for **each** question are shown in brackets
– *use this as a guide as to how much time to spend on each question.*

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.
- If you change your mind about an answer, cross it out and put your new answer and any working underneath.

Turn over ►

P76181A

©2024 Pearson Education Ltd.
V:1/1/1/1/1/




Pearson

1. Particle A has mass $4m$ and particle B has mass $3m$.

The particles are moving in opposite directions along the same straight line on a smooth horizontal surface when they collide directly.

Immediately **before** the collision, the speed of A is $2x$ and the speed of B is x .

Immediately **after** the collision, the speed of A is y and the speed of B is $5y$.

The direction of motion of each particle is reversed as a result of the collision.

(a) Show that $y = \frac{5}{11}x$. (3)

(b) Find, in terms of m and x , the magnitude of the impulse received by A in the collision. (3)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



2.




Figure 1

A non-uniform beam AB has length 6 m and mass 50 kg. The beam rests horizontally on two supports at C and D , where $AC = 0.9$ m and $DB = 1.8$ m.

A child of mass 25 kg stands on the beam at E , where $AE = EB = 3$ m, as shown in Figure 1.

The beam is in equilibrium.

The magnitude of the normal reaction between the beam and the support at C is R_C newtons.

The magnitude of the normal reaction between the beam and the support at D is R_D newtons.

The beam is modelled as a rod and the child is modelled as a particle.

The centre of mass of the beam is between C and D and is a distance x metres from D .

Given that $2R_D = 3R_C$

(a) show that $x = 1.38$

(6)

The child remains at E and a block of mass M kg is placed on the beam at B .

The block is modelled as a particle.

Given that the beam is on the point of tilting,

(b) find the value of M .

(3)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



3. [In this question \mathbf{i} and \mathbf{j} are horizontal unit vectors and position vectors are given relative to a fixed origin.]

A ship A is moving with constant velocity.

At 1 pm, the position vector of A is $(25\mathbf{i} + 10\mathbf{j})$ km.

At 3 pm, the position vector of A is $(55\mathbf{i} + 34\mathbf{j})$ km.

At time t hours after 1 pm, the position vector of A is \mathbf{r}_A km.

(a) Show that $\mathbf{r}_A = (25 + 15t)\mathbf{i} + (10 + 12t)\mathbf{j}$ (4)

The speed of A is $V \text{ ms}^{-1}$

(b) Find the value of V . (2)

A ship B is moving with constant velocity $(20\mathbf{i} - 6\mathbf{j}) \text{ km h}^{-1}$

At 1 pm, the position vector of B is $(35\mathbf{i} + 51\mathbf{j})$ km.

At 2:30 pm, B passes through the point P .

(c) Show that A also passes through P . (5)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



4. The points A and B lie on the same straight horizontal road.

Figure 2, on page 11, shows the speed-time graph of a cyclist P , for his journey from A to B .

At time $t = 0$, P starts from rest at A and accelerates uniformly for 9 seconds until his speed is $V \text{ m s}^{-1}$

He then travels at constant speed $V \text{ m s}^{-1}$

When $t = 42$, cyclist P passes B .

Given that the distance AB is 120 m,

- (a) show that $V = 3.2$ (3)
- (b) Find the acceleration of cyclist P between $t = 0$ and $t = 9$ (2)

Cyclist P continues to cycle along the road in the same direction at the same constant speed, $V \text{ m s}^{-1}$

When $t = 6$, a second cyclist Q sets off from A and travels in the same direction as P along the same road. She accelerates for T seconds until her speed is 3.6 m s^{-1}

She then travels at constant speed 3.6 m s^{-1}

Cyclist Q catches up with P when $t = 54$

- (c) On Figure 2, on page 11, sketch a speed-time graph showing the journeys of **both** cyclists, for the interval $0 \leq t \leq 54$ (3)
- (d) Find the value of T (5)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 4 continued




Figure 2

A copy of Figure 2 is on page 13 if you need to redraw your answer to part (c).

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA


DO NOT WRITE IN THIS AREA



P 7 6 1 8 1 A 0 1 1 2 4

Question 4 continued

Only use this copy of Figure 2 if you need to redraw your answer to part (c).



Copy of Figure 2

(Total for Question 4 is 13 marks)



P 7 6 1 8 1 A 0 1 3 2 4

5.




Figure 3

Two particles, P and Q , have masses 3 kg and 5 kg respectively. The particles are connected by a light inextensible string which passes over a small smooth fixed pulley.

The particles are released from rest with the string taut and the hanging parts of the string vertical, as shown in Figure 3.

Immediately after the particles are released from rest, P moves upwards with acceleration $a \text{ m s}^{-2}$ and the tension in the string is T newtons.

(a) Write down an equation of motion for P . (2)

(b) Find the value of T . (4)

The total force acting on the pulley due to the string has magnitude F newtons.

(c) Find the value of F . (2)

Initially, Q is 10 m above horizontal ground and P is more than 2 m below the pulley.

At the instant when Q has descended a distance of 2 m, the string breaks and Q falls to the ground.

(d) Find the speed of Q at the instant it hits the ground. (5)



6.




Figure 4

A particle P of mass 5 kg lies on the surface of a rough plane.

The plane is inclined at an angle α to the horizontal, where $\tan \alpha = \frac{3}{4}$

The particle is held in equilibrium by a horizontal force of magnitude H newtons, as shown in Figure 4.

The horizontal force acts in a vertical plane containing a line of greatest slope of the inclined plane.

The coefficient of friction between the particle and the plane is $\frac{1}{4}$

(a) Find the smallest possible value of H .

(6)

The horizontal force is now removed, and P starts to slide down the slope.

In the first T seconds after P is released from rest, P slides 1.5 m down the slope.

(b) Find the value of T .

(6)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



- 7 At time $t = 0$, a small ball A is projected vertically upwards with speed 8 m s^{-1} from a fixed point on horizontal ground.
The ball hits the ground again for the first time at time $t = T_1$ seconds.

Ball A is modelled as a particle moving freely under gravity.

- (a) Show that $T_1 = 1.63$ to 3 significant figures. (2)

After the first impact with the ground, A rebounds to a height of 2 m above the ground.

Given that the mass of A is 0.1 kg,

- (b) find the magnitude of the impulse received by A as a result of its first impact with the ground. (5)

At time $t = 1$ second, another small ball B is projected vertically upwards from another point on the ground with speed 5 m s^{-1}

Ball B is modelled as a particle moving freely under gravity.

At time $t = T_2$ seconds ($T_2 > 1$), A and B are at the same height above the ground for the first time.

- (c) Find the value of T_2 (4)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



