

Please check the examination details below before entering your candidate information

Candidate surname					Other names							
Pearson Edexcel					Centre Number				Candidate Number			
International Advanced Level					<input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>				<input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>			
Wednesday 8 January 2020												
Morning (Time: 1 hour 30 minutes)					Paper Reference WMA11/01							
Mathematics												
International Advanced Subsidiary/Advanced Level												
Pure Mathematics P1												
You must have: Mathematical Formulae and Statistical Tables (Lilac), calculator								Total Marks				

Candidates may use any calculator permitted by Pearson regulations. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
– *there may be more space than you need.*
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Inexact answers should be given to three significant figures unless otherwise stated.

Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- There are 11 questions in this question paper. The total mark for this paper is 75.
- The marks for **each** question are shown in brackets
– *use this as a guide as to how much time to spend on each question.*

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.
- If you change your mind about an answer, cross it out and put your new answer and any working underneath.

Turn over ►

P60796A

©2020 Pearson Education Ltd.

1/1/1/




Pearson

Leave
blank

1. Find, in simplest form,

$$\int \left(\frac{8x^3}{3} - \frac{1}{2\sqrt{x}} - 5 \right) dx$$

(4)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Leave
blank

2. Given $y = 3^x$, express each of the following in terms of y . Write each expression in its simplest form.

(a) 3^{3x}

(1)

(b) $\frac{1}{3^{x-2}}$

(2)

(c) $\frac{81}{9^{2-3x}}$

(2)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Leave
blank

Question 3 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Q3

(Total 6 marks)



Leave
blank

5. (a) Find, using algebra, all solutions of

$$20x^3 - 50x^2 - 30x = 0 \quad (3)$$

- (b) Hence find all real solutions of

$$20(y + 3)^{\frac{3}{2}} - 50(y + 3) - 30(y + 3)^{\frac{1}{2}} = 0 \quad (4)$$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Leave
blank

8. The straight line l has equation $y = k(2x - 1)$, where k is a constant.

The curve C has equation $y = x^2 + 2x + 11$

Find the set of values of k for which l does not cross or touch C .

(6)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Leave
blank

9. In this question you must show all stages of your working.

Solutions relying on calculator technology are not acceptable.

A curve has equation

$$y = \frac{4x^2 + 9}{2\sqrt{x}} \quad x > 0$$

Find the x coordinate of the point on the curve at which $\frac{dy}{dx} = 0$

(6)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Leave
blank

10. The curve C_1 has equation $y = f(x)$, where

$$f(x) = (4x - 3)(x - 5)^2$$

(a) Sketch C_1 showing the coordinates of any point where the curve touches or crosses the coordinate axes.

(3)

(b) Hence or otherwise

(i) find the values of x for which $f\left(\frac{1}{4}x\right) = 0$

(ii) find the value of the constant p such that the curve with equation $y = f(x) + p$ passes through the origin.

(2)

A second curve C_2 has equation $y = g(x)$, where $g(x) = f(x + 1)$

(c) (i) Find, in simplest form, $g(x)$. You may leave your answer in a factorised form.

(ii) Hence, or otherwise, find the y intercept of curve C_2

(3)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Leave
blank

11. A curve has equation $y = f(x)$, where

$$f''(x) = \frac{6}{\sqrt{x^3}} + x \quad x > 0$$

The point $P(4, -50)$ lies on the curve.

Given that $f'(x) = -4$ at P ,

(a) find the equation of the normal at P , writing your answer in the form $y = mx + c$, where m and c are constants,

(3)

(b) find $f(x)$.

(8)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



