



Mark Scheme (Results)

October 2024

Pearson Edexcel International Advanced Level
In Pure Mathematics (WMA13) Paper 01

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

October 2024

Question Paper Log Number P76188A

Publications Code WMA13_01_2410_MS

All the material in this publication is copyright

© Pearson Education Ltd 2024

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

General Instructions for Marking

1. The total number of marks for the paper is 75.
2. The Edexcel Mathematics mark schemes use the following types of marks:
 - **M marks: Method marks are awarded for 'knowing a method and attempting to apply it', unless otherwise indicated.**
 - A marks: Accuracy marks can only be awarded if the relevant method (M) marks have been earned.
 - B marks are unconditional accuracy marks (independent of M marks)

Marks should not be subdivided.

3. Abbreviations

These are some of the traditional marking abbreviations that will appear in the mark schemes and can be used if you are using the annotation facility on ePEN:

- bod – benefit of doubt
 - ft – follow through
 - the symbol \surd will be used for correct ft
 - cao – correct answer only
 - cso – correct solution only. There must be no errors in this part of the question to obtain this mark
 - isw – ignore subsequent working
 - awrt – answers which round to
 - SC – special case
 - oe – or equivalent (and appropriate)
 - **d... or dep** – dependent
 - indep – independent
 - dp – decimal places
 - sf – significant figures
 - * – The answer is printed on the paper or ag- answer given
 - \square or **d...** – The second mark is dependent on gaining the first mark
4. **All A marks are 'correct answer only' (cao), unless shown, for example, as A1** ft to indicate that previous wrong working is to be followed through. After a misread however, the subsequent A marks affected are treated as A ft, but manifestly absurd answers should never be awarded A marks.

5. For misreading which does not alter the character of a question or materially simplify it, deduct two from any A or B marks gained, in that part of the question affected. If you are using the annotation facility on ePEN, indicate **this action by 'MR' in the** body of the script.

6. If a candidate makes more than one attempt at any question:
 - a) If all but one attempt is crossed out, mark the attempt which is NOT crossed out.
 - b) If either all attempts are crossed out or none are crossed out, mark all the attempts and score the highest single attempt.

7. Ignore wrong working or incorrect statements following a correct answer.

General Principles for Pure Mathematics Marking
(NB specific mark schemes may sometimes override these general principles)

Method mark for solving 3 term quadratic:

1. Factorisation

$$(x^2 + bx + c) = (x + p)(x + q), \text{ where } |pq| = |c| \text{ leading to } x = \dots$$

$$(ax^2 + bx + c) = (mx + p)(nx + q), \text{ where } |pq| = |c| \text{ and } |mn| = |a| \text{ leading to } x = \dots$$

2. Formula

Attempt to use correct formula (with values for a, b and c)

3. Completing the square

$$\text{Solving } x^2 + bx + c = 0: (x \pm \frac{b}{2})^2 \pm q \pm c, \quad q \neq 0 \text{ leading to } x = \dots$$

Method marks for differentiation and integration:

1. Differentiation

Power of at least one term decreased by 1 ($x^n \rightarrow x^{n-1}$)

2. Integration

Power of at least one term increased by 1 ($x^n \rightarrow x^{n+1}$)

Use of a formula

Where a method involves using a formula that has been learnt, the advice given in recent examiners' reports is that the formula should be quoted first.

Normal marking procedure is as follows:

Method mark for quoting a correct formula and attempting to use it, even if there are small mistakes in the substitution of values.

Where the formula is not quoted, the method mark can be gained by implication from correct working with values, but may be lost if there is any mistake in the working.

Exact answers

Examiners' reports have emphasised that where, for example, an exact answer is asked for, or working with surds is clearly required, marks will normally be lost if the candidate resorts to using rounded decimals.

Answers without working

The rubric says that these may not gain full credit. Individual mark schemes will give details of what happens in particular cases. General policy is that if it could be done "in your head", detailed working would not be required. Most candidates do show working, but there are occasional awkward cases and if the mark scheme does not cover this, please contact your team leader for advice.

Question Number	Scheme	Marks
1	$3 \tan^2 \theta + 7 \sec \theta - 3 = 0 \Rightarrow 3(\sec^2 \theta - 1) + 7 \sec \theta - 3 = 0$	M1
	$3 \sec^2 \theta + 7 \sec \theta - 6 = 0$	A1
	$(3 \sec \theta - 2)(\sec \theta + 3) = 0 \Rightarrow \sec \theta = \dots \Rightarrow \cos \theta = \dots$	dM1
	$\theta = 109.5^\circ, 250.5^\circ$	A1, A1
		(5)
	Total 5	

Main method: Also allow if another variable is used e.g $x \leftrightarrow \theta$

M1: Attempts to set up a quadratic equation in $\sec \theta$ using $\tan^2 \theta = \pm 1 \pm \sec^2 \theta$

Condone bracketing slips, for example, where $3 \tan^2 \theta$ is replaced by $\pm 3 \sec^2 \theta \pm 1$

A1: Collects terms to form a correct 3TQ, $3 \sec^2 \theta + 7 \sec \theta - 6 = 0$ o.e

Terms do not need to be all on one side of the equation and the $= 0$ can be implied by subsequent work.

dM1: Dependent upon the previous M1, it is scored for the method of producing a solvable non zero solution for $\cos \theta$

Look for;

- A valid way of solving their quadratic equation in $\sec \theta$ using any allowable method including a calculator leading to $\sec \theta = \dots$
- Use of $\sec \theta = \frac{1}{\cos \theta}$ to produce a solution $\cos \theta = k$ where $-1 < k < 1, k \neq 0$

E.g. $3 \sec^2 \theta + 7 \sec \theta - 6 = 0 \Rightarrow \sec \theta = -3 \Rightarrow \cos \theta = -\frac{1}{3}$. You can ignore the 2nd solution

If the error had been $\tan^2 \theta = \sec^2 \theta + 1$ look for $3 \sec^2 \theta + 7 \sec \theta = 0 \Rightarrow \sec \theta = -\frac{7}{3} \Rightarrow \cos \theta = -\frac{3}{7}$

A1: One of $\theta = \text{awrt } 109.5^\circ, \text{awrt } 250.5^\circ$ following M1, A1, dM1

Allow either answer in radians here so score for awrt 1.91 or awrt 4.37

A1: CSO. Both $\theta = \text{awrt } 109.5^\circ$ and $\text{awrt } 250.5^\circ$ and no extras in the range

.....
Other methods do exist but the same marking principles will apply

Alt method:

M1: For an attempt to set up a quadratic equation in a single trigonometric term, e.g. $\cos \theta$

Replaces $\tan^2 \theta$ by $\frac{\sin^2 \theta}{\cos^2 \theta}$, $\sec \theta$ by $\frac{1}{\cos \theta}$ and then uses $\sin^2 \theta = \pm 1 \pm \cos^2 \theta \Rightarrow$ quadratic equation in

$\cos \theta$ E.g. $3 \tan^2 \theta + 7 \sec \theta - 3 = 0 \Rightarrow 3 \frac{\sin^2 \theta}{\cos^2 \theta} + 7 \times \frac{1}{\cos \theta} - 3 = 0 \Rightarrow 3 \sin^2 \theta + 7 \cos \theta - 3 \cos^2 \theta = 0$

and then uses $\sin^2 \theta = \pm 1 \pm \cos^2 \theta \Rightarrow 3(1 - \cos^2 \theta) + 7 \cos \theta - 3 \cos^2 \theta = 0$.

Condone slips

A1: For a correct 3TQ in $\cos \theta$. FYI $6 \cos^2 \theta - 7 \cos \theta - 3 = 0$. Terms do not have to be on one side

dM1: Solves the quadratic equation in $\cos \theta$ using an appropriate method including a calculator.

It must lead to a solution $\cos \theta = k$ where $-1 < k < 1, k \neq 0$

If a calculator is used the solution must be correct for their equation.

$$\text{E.g. } 6\cos^2 \theta - 7\cos \theta - 3 = 0 \Rightarrow \cos \theta = -\frac{1}{3}$$

A1: One of $\theta = \text{awrt } 109.5^\circ, \text{awrt } 250.5^\circ$ following M1, A1, dM1

Allow either answer in radians here so score for awrt 1.91 or awrt 4.37

A1: CSO. Both $\theta = \text{awrt } 109.5^\circ$ and $\text{awrt } 250.5^\circ$ and no extras in the range.

If both solutions to the quadratic were given then both must be correct

.....

Methods using calculator technology and not showing all stages of working as required by question:

Example 1:

$$3 \tan^2 \theta + 7 \sec \theta - 3 = 0 \Rightarrow \theta = 109.5^\circ, 250.5^\circ$$

No marks

Example 2:

$$3 \tan^2 \theta + 7 \sec \theta - 3 = 0 \Rightarrow 3(\sec^2 \theta - 1) + 7 \sec \theta - 3 = 0$$

$$\Rightarrow 3 \sec^2 \theta + 7 \sec \theta - 6 = 0 \Rightarrow \sec \theta = -3 \Rightarrow \theta = 109.5^\circ, 250.5^\circ$$

Scores M1, A1, dM0, A0, A0

Example 3:

$$3 \tan^2 \theta + 7 \sec \theta - 3 = 0 \Rightarrow 6 \cos^2 \theta - 7 \cos \theta - 3 = 0 \text{ via identities followed by } \Rightarrow \theta = 109.5^\circ, 250.5^\circ$$

Scores M1, A1, dM0, A0, A0

Methods using calculator technology and showing sufficient stages of working:

Example 4: $3 \tan^2 \theta + 7 \sec \theta - 3 = 0 \Rightarrow 3 \sec^2 \theta + 7 \sec \theta - 6 = 0$ via identity followed by

$$\sec \theta = -3 \Rightarrow \cos \theta = -\frac{1}{3} \Rightarrow \theta = \pm 109.5^\circ$$

Scores M1, A1, dM1, A1, A0 (only one of the two solutions given)

Example 5: $3 \tan^2 \theta + 7 \sec \theta - 3 = 0 \Rightarrow 6 \cos^2 \theta - 7 \cos \theta - 3 = 0$ via identities followed by

$$\cos \theta = -\frac{1}{3} \Rightarrow \theta = 109.5^\circ, 250.5^\circ \text{ M1, A1, dM1, A1, A1}$$

Question Number	Scheme	Marks
2 (a)	$x = 2y^2 + 5y - 6$	
	$\frac{dx}{dy} = 4y + 5 \Rightarrow \frac{dy}{dx} = \frac{1}{4y + 5}$	M1, A1
		(2)
(b)	Sets their $4y + 5 = 0 \Rightarrow y = -\frac{5}{4}$	M1
	Substitutes their $y = -\frac{5}{4}$ into $x = 2y^2 + 5y - 6$ to find x	dM1
	$\left(-\frac{73}{8}, -\frac{5}{4}\right)$	A1
		(3)
		Total 5

Mark as one complete question

Main method via differentiation in (a) and (b)

(a)

M1: Differentiates $2y^2 + 5y - 6$ to a linear term in y AND attempts to take the reciprocal.

Condoned for this mark attempts such as $\frac{dx}{dy} = 4y + 5 \Rightarrow \frac{dy}{dx} = \frac{1}{4y} + \frac{1}{5}$

We may see other methods including implicit differentiation from WMA14.

Look for $x = 2y^2 + 5y - 6 \Rightarrow 1 = ay \frac{dy}{dx} + b \frac{dy}{dx}$ followed by an attempt to make $\frac{dy}{dx}$ the subject

A1: $\frac{dy}{dx} = \frac{1}{4y + 5}$ ISW after a correct answer.

(b) **Marks in part (b) can only be awarded after sight of** $\frac{dx}{dy} = \frac{ay + b}{k}$ **or** $\frac{dy}{dx} = \frac{k}{ay + b}$

So full marks can only be awarded after sight of $\frac{dx}{dy} = 4y + 5$ **or** $\frac{dy}{dx} = \frac{1}{4y + 5}$

M1: Sets ' $ay + b = 0 \Rightarrow y = \dots$ ' following sight of $\frac{dx}{dy} = \frac{ay + b}{k}$ **or** $\frac{dy}{dx} = \frac{k}{ay + b}$

Ignore/condone incorrect statements such as $\frac{dy}{dx} = 0$ here

This can be implied. E.g $\frac{dy}{dx} = \frac{1}{4y + 5}$ followed by $y = -\frac{5}{4}$

dM1: Substitutes their solution of their $4y + 5 = 0$ into $x = 2y^2 + 5y - 6$ to find x .

This is dependent upon the previous M

A1: $\left(-\frac{73}{8}, -\frac{5}{4}\right)$ following $\frac{dx}{dy} = 4y + 5$ **or** $\frac{dy}{dx} = \frac{1}{4y + 5}$

The answer may be given separately, for example, as $x = -9.125, y = -1.25$

ISW after a correct answer.

Further guidance to highlighted sentence.

If (a) is incorrect or incomplete follow the following guidance.

Incorrect (a): If they incorrectly state $\frac{dy}{dx} = 4y + 5$ o.e. and go on to use $4y + 5 = 0 \Rightarrow y = \dots$ and write down a correct set of coordinates for P they will score NO MARKS in (b)

Incomplete (a): If they state $\frac{dx}{dy} = 4y + 5$ in (a) and go on to use $4y + 5 = 0 \Rightarrow y = \dots$ in (b) they will score no marks in (a) but could potentially score full marks in (b)

You can award all 3 marks for $\left(-\frac{73}{8}, -\frac{5}{4}\right)$ following either $\frac{dx}{dy} = 4y + 5$ or $\frac{dy}{dx} = \frac{1}{4y + 5}$

.....
 Alternative approach: Part (a) may well be blank or incorrect. Part(b) done via completion of square

It is possible to do part (b) by not doing any differentiating.



$$x = 2y^2 + 5y - 6 \Rightarrow x = 2\left(y + \frac{5}{4}\right)^2 - \frac{73}{8}$$

M1: An allowable attempt to complete the square followed by a valid attempt at either x or y from their attempt

A1: For a correct x or y

A1: For a correct x and y

.....

Question Number	Scheme	Marks
<p>3 (a)</p>	 <p style="text-align: right;">$y = 48$</p> <p style="text-align: center;">Graphical interpretation of $f(x) = 48$</p>	
	<p>One of 8, -8</p>	<p>B1</p>
	<p>Attempts to solve an appropriate equation E.g. $2x^2 - 10x = 48 \Rightarrow x^2 - 5x - 24 = 0 \Rightarrow (x \pm 8)(x \pm 3) = 0 \Rightarrow x = \dots$</p>	<p>M1</p>
	<p>$x = 8, -8$ with no additional values</p>	<p>A1</p>
	<p>(3)</p>	<p>(b)</p>  <p style="text-align: center;">Graphical interpretation of $f(x) \dots \frac{5}{2}x$</p> <p style="text-align: left;">$y = \frac{5}{2}x$</p>
<p>Attempts to solve $2x^2 - 10x = \frac{5}{2}x \Rightarrow 4x^2 - 25x = 0 \Rightarrow x = \frac{25}{4}$ OR attempts to solve (o.e.) $10x - 2x^2 = \frac{5}{2}x \Rightarrow 4x^2 - 15x = 0 \Rightarrow x = \frac{15}{4}$</p>	<p>M1</p>	
<p>Attempts to solve $2x^2 - 10x = \frac{5}{2}x \Rightarrow 4x^2 - 25x = 0 \Rightarrow x = \frac{25}{4}$ AND attempts to solve (o.e.) $10x - 2x^2 = \frac{5}{2}x \Rightarrow 4x^2 - 15x = 0 \Rightarrow x = \frac{15}{4}$</p>	<p>dM1</p>	
<p>Achieves both critical values $x = \frac{15}{4}, x = \frac{25}{4}$</p>	<p>A1</p>	
<p>Correct set of values $x, \frac{15}{4}$ or $x \dots \frac{25}{4}$</p>	<p>A1</p>	
<p>(4)</p>	<p>Total 7</p>	

This question demands that some working should be shown

(a) This part is now being marked B1 M1 A1 instead of M1 A1 A1

B1: Correctly obtains 8 or -8 as a solution (for x) . Condone solution appearing as $|x| = 8$

M1: Attempts to solve $f(|x|) = 48$ via an appropriate method including a calculator, factors or formula

So, look for a correct equation followed by an attempt to solve.

The equation must be given in the form of a correct 3TQ, but not necessarily collected on one side of the = sign. Solving is allowed by all methods including via a calculator.

All the following score this M mark

Example 1:

$$2x^2 - 10x = 48 \text{ o.e. } \Rightarrow x = 8, (-3) \dots \text{only the 8 is required}$$

Example 2:

$$2x^2 - 10x = 48 \text{ o.e. } \Rightarrow x^2 - 5x - 24 = 0 \Rightarrow (x-8)(x+3) = 0 \Rightarrow x = \dots$$

Example 3:

$$2|x|^2 - 10|x| - 48 = 0 \text{ or } |x|^2 - 5|x| - 24 = 0 \text{ leading to } |x| \text{ or } x = 8, (-3)$$

Example 4:

$$2x^2 + 10x - 48 = 0 \Rightarrow x = \frac{-10 \pm \sqrt{100 - 4(2)(-48)}}{2(2)} = \dots$$

Example 5:

$$2x^2 + 10x = 48 \text{ o.e. } \Rightarrow x = -8, (3) \dots \text{only the } -8 \text{ is required}$$

A1 $x = 8, -8$ with no additional values unless they have been rejected

(b)

M1: Attempts to solve a correct equation or inequality, leading to a non-zero value for x .

The method **must be shown** with the terms being collected as a minimum response

Look for $2x^2 - 10x = \frac{5}{2}x \Rightarrow ax^2 \pm bx = 0 \Rightarrow x = c, c \neq 0$. Allow with the = as any inequality including <

OR $-2x^2 + 10x = \frac{5}{2}x \Rightarrow ax^2 \pm bx = 0 \Rightarrow x = c, c \neq 0$. Allow with the = as any inequality including <

You may see as an alternative to the 2nd equation/inequality $2x^2 - 10x = -\frac{5}{2}x \Rightarrow ax^2 \pm bx = 0 \Rightarrow x = c, c \neq 0$

It is acceptable to divide through by x , so $2x^2 - 10x = \frac{5}{2}x \Rightarrow 2x - 10 = \frac{5}{2} \Rightarrow 2x = \frac{25}{2} \Rightarrow x = 6.25$

dM1: Attempts to solve BOTH correct equations or inequations leading to two non-zero values for x

Allow even if there are extra equations solved.

A1: Achieves both critical values $x = \frac{15}{4}$ and $x = \frac{25}{4}$ and no other values apart from 0 (and perhaps 5)

following both equations or inequalities. Allow with incorrect inequalities $x \dots \frac{15}{4}$ and $x \dots \frac{25}{4}$, you are

looking for the critical values only

A1: Correct range in allowable form. E.g. $x \dots \frac{15}{4}$ or $x \dots \frac{25}{4}, \left(-\infty, \frac{15}{4}\right] \cup \left[\frac{25}{4}, \infty\right)$ but condone $x \dots \frac{15}{4}, x \dots \frac{25}{4}$

Do NOT incorrect forms such as $x \dots \frac{15}{4}$ and $x \dots \frac{25}{4}, \left(-\infty, \frac{15}{4}\right] \cap \left[\frac{25}{4}, \infty\right), \frac{25}{4} \dots x \dots \frac{15}{4}$

The two aspects must be brought together on a single line. Mark their final answer

Alt Method: You may see attempts via squaring. It can be marked using the above guidelines

$$(2x^2 - 10x)^2 = \left(\frac{5}{2}x\right)^2 \Rightarrow x^2(16x^2 - 160x + 375) = 0 \Rightarrow x^2(4x - 15)(4x - 25) = 0$$

Question Number	Scheme	Marks
4 (a)	$\log_{10} N = 2 \Rightarrow N = 100$	B1
		(1)
(b)	$\log_{10} N = 0.35t + 2 \Rightarrow N = 10^{0.35t+2}$	M1
	$\Rightarrow N = 10^{0.35t} \times 10^2$	dM1
	$\Rightarrow N = 100 \times 2.24^t$	A1
		(3)
(c)	Rate of growth at 5 hours is $\frac{dN}{dt} = 100 \times \ln 2.24 \times 2.24^5$	M1
	Allow an answer in the range 4530 to 4550 (bacteria per hour)	A1
		(2)
		Total 6

(a)

B1: For 100 Do not accept just 10^2 and cannot be awarded from 100 in part (b)

(b) **The demand of the question is ‘show that’, not just find values of a and b**

Method 1: starting with $\log_{10} N = 0.35t + 2$

M1: Undoes the log. E.g. $\log_{10} N = 0.35t + 2 \Rightarrow N = 10^{0.35t+2}$

Give BOD for those who go from $\log_{10} N = 0.35t + 2$ straight to $N = 10^{0.35t} \times 10^2$

dM1: And follows with a correct application of the index law.

Look for $\log_{10} N = 0.35t + 2 \Rightarrow N = 10^{0.35t+2} = 10^{0.35t} \times 10^2$ o.e such as $10^{0.35t} \times 100$

But BOD for $\log_{10} N = 0.35t + 2 \Rightarrow N = 10^{0.35t} \times 10^2$ o.e. such as $10^{0.35t} \times 100$

And BOD for $\log_{10} N = 0.35t + 2 \Rightarrow N = 10^{0.35t+2} \Rightarrow N = 100 \times 2.24^t$

A1: CSO $N = 100 \times 2.24^t$ with no errors following M1, dM1. Allow awrt 2.24

Condone $a = 100, b = 2.24$ following the award of M1 dM1

You should expect to see at least one correct line of working and no errors

Method 2: starting with $N = ab^t$

M1: Takes logs of both sides and applies one correct rule. E.g. $N = ab^t \Rightarrow \log N = \log a + \log b^t$

Give BOD for those who write $N = ab^t$ followed by $\log N = \log a + t \log b$

dM1: And follows with $\log N = \log a + t \log b$ before setting

$\log a = 2$ and $\log b = 0.35$ and proceeding to values for a and b

A1: Fully shows and produces correct values following M1, dM1

CSO $N = 100 \times 2.24^t$ with no errors. Allow awrt 2.24

Condone $a = 100, b = 2.24$ following the award of M1 dM1

You should expect to see at least one correct line of working and no errors

Method 3: A simultaneous approach that finds values but does not show the result

M1: Sets up two simultaneous equations in a and b

E.g. Uses $t = 0, N = 100 \Rightarrow a = 100$ and $t = 1, N = 10^{2.35} \Rightarrow ab^1 = 223.9$

dM1: Solves to find values for a and b .

A0: Has not **shown** the result so cannot be awarded

Common error: Special Case

$$\log_{10} N = 0.35t + 2 \Rightarrow N = 10^{0.35t} + 10^2 \text{ so } a = 100 \text{ and } b = 2.24$$

Or even

$$\log_{10} N = 0.35t + 2 \Rightarrow N = 10^{0.35t} + 10^2 \Rightarrow N = 100 \times 2.24^t \text{ so } a = 100 \text{ and } b = 2.24$$

Such work resulting in the **correct** values for a and b scores SC 100

(c) Full marks in part (c) can be scored following the SC

M1: Attempts $\frac{dN}{dt} = K \times "2.24"^{t^5}$ following through on their 2.24.

This cannot be scored from substituting $t = 5$ into their ab^t

A1: Calculates $\frac{dN}{dt} = 100 \times \ln 2.24 \times 2.24^5$ and achieves an answer in the range 4530 to 4550 (bacteria per hour)

Question Number	Scheme	Marks
5(a)	$\sin 3x = \sin(2x + x) = \sin 2x \cos x + \cos 2x \sin x$	M1
	$= (2 \sin x \cos x) \cos x + (1 - 2 \sin^2 x) \sin x$	dM1
	$= 2 \sin x \cos^2 x + \sin x - 2 \sin^3 x$	A1
	$= 2 \sin x (1 - \sin^2 x) + \sin x - 2 \sin^3 x = 3 \sin x - 4 \sin^3 x$	A1
		(4)
(b)	$2 \sin 3\theta = 5 \sin 2\theta \Rightarrow 2(3 \sin \theta - 4 \sin^3 \theta) = 10 \sin \theta \cos \theta$	M1
	Divides or takes out $\sin \theta$ as a factor and uses $\sin^2 \theta = 1 - \cos^2 \theta$ to set up and solve a 3TQ in $\cos \theta$ E.g. $\Rightarrow 6 \sin \theta - 8 \sin^3 \theta = 10 \sin \theta \cos \theta \Rightarrow 6 - 8(1 - \cos^2 \theta) = 10 \cos \theta$ $\Rightarrow 4 \cos^2 \theta - 5 \cos \theta - 1 = 0$ $\Rightarrow \cos \theta = \frac{5 - \sqrt{41}}{8} = (-0.175\dots)$	dM1
	Any two of the following four answers $\sin \theta = 0 \Rightarrow \theta = 180^\circ, 360^\circ$ $\cos \theta = \frac{5 - \sqrt{41}}{8} \Rightarrow \theta = \text{awrt } 100^\circ \text{ or awrt } 260^\circ$	A1
	All of $180^\circ, 360^\circ, \text{awrt } 100.1^\circ, \text{awrt } 259.9^\circ$	A1
		(4)
		Total 8

(a)

M1: Attempts $\sin 3x = \sin(2x + x)$ and achieves $\pm \sin 2x \cos x \pm \cos 2x \sin x$. The $= \sin(2x + x)$ may be implied

dM1: Dependent upon the previous mark. It is for using $\sin 2x = 2 \sin x \cos x$ and $\cos 2x = \pm 1 \pm 2 \sin^2 x$, $\pm 1 \pm 2 \cos^2 x$ or $\pm(\cos^2 x - \sin^2 x)$ to form an expression in just $\sin x$ and $\cos x$.

A1: Fully correct AND expanded expression in just $\sin x$ and $\cos x$

Examples of correct expanded expressions are;

$$2 \sin x \cos^2 x + \sin x - 2 \sin^3 x$$

$$2 \sin x \cos^2 x + 2 \sin x \cos^2 x - \sin x$$

$$2 \sin x \cos^2 x + \sin x \cos^2 x - \sin^3 x$$

or $4 \sin x \cos^2 x - \sin x$

Condone a missing x on a $\sin x$, for example. Condone a slip in notation, e.g. writing $\sin x^3$ instead of $\sin^3 x$ as long as it is subsequently corrected.

A1: Proceeds correctly to $3 \sin x - 4 \sin^3 x$ using $\cos^2 x = 1 - \sin^2 x$

Penalise notational and bracketing errors on this mark only

E.g. Writing $\cos x^2$ for $\cos^2 x$, mixing variables, missing x on a $\sin x$ etc

(b) Allow if solved in terms of another variable, say x

The question states hence or otherwise solve $2\sin 3\theta = 5\sin 2\theta$

In almost all cases they will use their part (a) and proceed by the 'hence' route.

'Otherwise' routes could start with earlier versions of $2\sin 3\theta$ such as $2(2\sin x \cos^2 x + \sin x - 2\sin^3 x)$

In almost all cases the main mark scheme provides a framework of how the marks are awarded

- M1: Uses their part (a) and a double angle for $\sin 2\theta$ of the form $G\sin \theta \cos \theta$ to form an equation in $\sin \theta$ and $\cos \theta$. Look for $R\sin \theta + S\sin^3 \theta = T\sin \theta \cos \theta$ o.e. Condone slips such as forgetting to $\times 2$
- dM1: The mark is scored for dividing or taking out a factor of $\sin \theta$ and using $\sin^2 \theta = \pm 1 \pm \cos^2 \theta$ to set up and solve a 3TQ in $\cos \theta$. To solve you should expect to see a valid method used, leading to at least one value for $\cos \theta = k$, $-1 < k < 1, k \neq 0$
- If a calculator is used to solve the 3TQ in $\cos \theta$ the value(s) must be correct for their equation. Candidates cannot go straight from an equation e.g. $4\cos^2 \theta - 5\cos \theta - 1 = 0$ to a solution e.g. 100.1° without sight of the value for $\cos \theta$ in between.
- A1: Finds two angles of $180^\circ, 360^\circ$, awrt 100° , awrt 260° following at least one previous M1
- The 180° and/or 360° must be found via a correct method with $\sin \theta = 0$ being seen or implied and also produced from a factorised $P\sin \theta + Q\sin^3 \theta = R\sin \theta \cos \theta$
- Likewise, the 100° and 260° must have been achieved via solving a correct equation.
- Allow this mark if the answers are given in radians. So, for two angles of $\pi, 2\pi$, awrt 1.75, awrt 4.54
- A1: All of $180^\circ, 360^\circ$, awrt 100.1° , awrt 259.9° with no extra values within the range following M1, dM1

There are no marks for answers appearing without valid method and working

There are other (usually more complicated) ways of attempting part (b) so you must look carefully through each response. If you think a response deserves credit and you are unsure of how to mark, please send to review.

Question Number	Scheme	Marks
6 (a)	$gf(2) = g(3) = 3^2 + 5$	M1
	$= 14$	A1
		(2)
(b)	Correct attempt at inverse $y = 6 - \frac{21}{2x+3} \Rightarrow 2x+3 = \frac{21}{6-y} \Rightarrow x = \left(\frac{21}{6-y} - 3\right) \div 2$	M1
	$f^{-1}(x) = \frac{3x+3}{2(6-x)}$ or $f^{-1}(x) = \frac{21}{2(6-x)} - \frac{3}{2}$	A1
	$-1, x < 6$	B1
		(3)
(c)	$gg(x) = 126 \Rightarrow (x^2 + 5)^2 + 5 = 126 \Rightarrow x^2 + 5 = \sqrt{121}$	M1
	$\Rightarrow x^2 = "6" \Rightarrow x = (\pm)\sqrt{6}$	dM1
	$\Rightarrow x = \pm\sqrt{6}$	A1
		(3)
		Total 8

(a)

M1: Full attempt at $gf(2)$ applying the rules in the correct order

Look for one of

- An attempt at $6 - \frac{21}{2 \times 2 + 3} \rightarrow k$ followed by an attempt at $k^2 + 5$
- An attempt at substituting $x = 2$ into $\left(6 - \frac{21}{2x+3}\right)^2 + 5$

A1: 14

Allow just 14 with no incorrect working for both marks

(b)

M1: For an attempt at the method of finding the inverse. It is the process you are scoring here.

Look for a process that starts with y as a function of x and proceeds to x as a function of just y .

Starting with $y = 6 - \frac{21}{2x+3}$, or possibly a changed function of the form $y = \frac{ax+b}{2x+3}$, which proceeds to a

form $x =$ function of **just** y . Some candidates may swap x and y first. E.g. $x = 6 - \frac{21}{2y+3}$ and proceed to

function $y =$ function of just x which is fine.

A1: Correct inverse $f^{-1}(x) = \frac{3x+3}{2(6-x)}$ or $f^{-1}(x) = \frac{21}{2(6-x)} - \frac{3}{2}$ o.e. but condone y or f^{-1} for $f^{-1}(x)$

Allow other equivalents such as $y = \frac{-3x-3}{2(x-6)}$, $y = \frac{10.5}{(6-x)} - 1.5$ and $y = \left(\frac{21}{6-x} - 3\right) \div 2$

ISW following a correct answer

B1: Correct domain written in correct form. E.g. $-1, x < 6$, $[-1, 6)$

(c)

M1: For an attempt to "undo" g once in an attempt to find the value of $g(x)$, x^2 or $x^2 + 5$.

This may be scored for any of the following methods (condoning slips)

- $(x^2 + 5)^2 + 5 = 126 \Rightarrow x^2 + 5 = \sqrt{121}$
- $(x^2 + 5)^2 + 5 = 126 \Rightarrow x^4 + 10x^2 - 96 = 0 \Rightarrow (x^2 - 6)(x^2 + 16) = 0 \Rightarrow x^2 = \dots$
- Condone $(x^2 + 5)^2 + 5 = 126 \Rightarrow x^4 + 10x^2 - 96 = 0 \Rightarrow x = \text{awrt } 2.45 \text{ OR } x^2 = 6$
- $g(x) = \sqrt{126 - 5}$

Do not award this mark if the candidate believes that to find x you only need to undo g once.
So, answers like $x = 11$ or $x = 6$ will score 0,0,0 unless further work is seen

dM1: For a complete attempt to find the exact solution to the equation. Examples of this would include

- $(x^2 + 5)^2 + 5 = 126 \Rightarrow x^2 + 5 = \sqrt{121} \Rightarrow x = \sqrt{\sqrt{121} \pm 5}$
- $(x^2 + 5)^2 + 5 = 126 \Rightarrow x^4 + 10x^2 - 96 = 0 \Rightarrow (x^2 - 6)(x^2 + 16) = 0 \Rightarrow x^2 = 6 \Rightarrow x = \sqrt{6}$
- $(x^2 + 5)^2 + 5 = 126 \Rightarrow x = \sqrt{6}$ or $(x^2 + 5)^2 + 5 = 126 \Rightarrow x = -\sqrt{6}$
- $(x^2 + 5)^2 + 5 = 126 \Rightarrow x^4 + 10x^2 - 96 = 0 \Rightarrow x = \sqrt{6}$
- $x = \sqrt{11 - 5}$

A1: $x = \pm\sqrt{6}$ only. Ignore any reference to $\pm 4i$ if included.Sight of $\pm\sqrt{6}$ without incorrect working scores all 3 marks

Question Number	Scheme	Marks
7 (a)	$f(x) = x^3 \sqrt{4x+7} \Rightarrow f'(x) = 3x^2 \sqrt{4x+7} + 2x^3 (4x+7)^{-\frac{1}{2}}$	M1, A1
	$\Rightarrow f'(x) = 3x^2 \sqrt{4x+7} + \frac{2x^3}{\sqrt{4x+7}} = \frac{3x^2(4x+7) + 2x^3}{\sqrt{4x+7}}$	dM1
	$\Rightarrow f'(x) = \frac{7x^2(2x+3)}{\sqrt{4x+7}}$	A1
		(4)
(b)	Substitutes $x = -\frac{3}{2}$ into $x^3 \sqrt{4x+7} \Rightarrow y = \dots$	M1
	$\left(-\frac{3}{2}, -\frac{27}{8}\right)$	A1
		(2)
(c)	Attempts $-4 \times -\frac{27}{8}$	M1
	$y = \frac{27}{2}$	A1
		(2)
(d)	$(2, 7)$	M1, A1
		(2)
		Total 10

(a)

M1: Attempts the product rule to achieve $f'(x) = Px^2 \sqrt{4x+7} + Qx^3 (4x+7)^{-\frac{1}{2}}$ where P and Q are positive constants.

Other methods are possible so look at each one carefully.

For example: $f(x) = x^3 \sqrt{4x+7} \Rightarrow f(x) = \sqrt{4x^7 + 7x^6} \Rightarrow f'(x) = \frac{1}{2} (4x^7 + 7x^6)^{-\frac{1}{2}} \times (28x^6 + 42x^5)$

The same principles for the main mark scheme can be applied

A1: $f'(x) = 3x^2 \sqrt{4x+7} + 2x^3 (4x+7)^{-\frac{1}{2}}$ o.e. which may be left unsimplified

dM1: Dependent upon previous M1. It is for "correctly" producing a single fraction with a denominator of $\sqrt{4x+7}$.

Look for $f'(x) = Px^2 \sqrt{4x+7} + Qx^3 (4x+7)^{-\frac{1}{2}} \Rightarrow \frac{Px^2(4x+7) + Qx^3}{\sqrt{4x+7}}$ o.e

A1: $f'(x) = \frac{7x^2(2x+3)}{\sqrt{4x+7}}$

.....
It is possible to attempt this by WMA14 methods using $[f(x)]^2 = x^6(4x+7) = 4x^7 + 7x^6$

Score

M1, A1: $2f(x)f'(x) = 28x^6 + 42x^5$

dM1, A1: $f'(x) = \frac{14x^5(2x+5)}{2f(x)} = \frac{14x^5(2x+5)}{2x^3 \sqrt{4x+7}} = \frac{7x^2(2x+5)}{\sqrt{4x+7}}$

(b)

M1: Attempts to substitute $x = -\frac{3}{2}$ into $x^3\sqrt{4x+7}$ to find the y value of the minimum point .

This may be implied by the correct y value.

It can be attempted even if part (a) is missing or incorrect

If they ignore the given form of $f'(x)$ and use their own version, they must

- Set their $f'(x)$ which must be of the form $Px^2\sqrt{4x+7} + Qx^3(4x+7)^{-\frac{1}{2}} = 0$
- Solve $Px^2\sqrt{4x+7} + Qx^3(4x+7)^{-\frac{1}{2}} = 0$ by multiplying by $(4x+7)^{\frac{1}{2}}$ o.e. to achieve a non-zero value for x
- Substitute this non zero value for x in into $x^3\sqrt{4x+7}$ to find y

A1: $\left(-\frac{3}{2}, -\frac{27}{8}\right)$ which may be awarded separately as, for example, $x = -1.5, y = -3.375$

Ignore $(0, 0)$ if also given.

(c)

M1: Attempts $\pm 4 \times -\frac{27}{8}$ " which may be seen within an inequality.

A1: $y, \frac{27}{2}$ which must be correctly simplified.

Allow it to be written in other correct forms such as $g, \frac{27}{2}, g(x), \frac{27}{2}$ or $(-\infty, 13.5]$

The original function is f so $f(x), \frac{27}{2}$ would be M1 A0

(d)

M1: For one correct coordinate. Look for $(2, \dots)$ or $(\dots, 7)$. Allow for either $x = 2$ or $y = 7$

If the coordinates have been built up mark the final answer.

E.g. $\left(\frac{1}{2}, \frac{3}{8}\right) \rightarrow \left(2, \frac{3}{8}\right) \rightarrow (2, 15) \rightarrow (-6, 15)$ would score M0 A0

A1: $(2, 7)$. Allow $x = 2$ and $y = 7$ ISW after a correct answer

Question Number	Scheme	Marks
8 (a)	52 b.p.m.	B1
		(1)
(b)	32 b.p.m.	B1
		(1)
(c)	$\frac{dH}{dt} = -8e^{-0.2t} + 18e^{-0.9t}$	M1, A1
	Sets $-8e^{-0.2t} + 18e^{-0.9t} = 0 \Rightarrow 4e^{0.7t} = 9$	dM1
	$\Rightarrow 0.7t = \ln \frac{9}{4} \Rightarrow t = \dots$	M1
	$T = 1.158$ (minutes)	A1
		(5)
(d)	$37 = 32 + 40e^{-0.2t} - 20e^{-0.9t} \Rightarrow e^{-0.2t} = \frac{1 + 4e^{-0.9t}}{8}$	M1
	$\Rightarrow e^{0.2t} = \frac{8}{1 + 4e^{-0.9t}} \Rightarrow t = 5 \ln \left(\frac{8}{1 + 4e^{-0.9t}} \right)$	A1*
		(2)
(e)	$t_2 = 5 \ln \left(\frac{8}{1 + 4e^{-0.9 \times 10}} \right) = \dots$	M1
	$(t_2) = \text{awrt } 10.3947$	A1
	$(M) = 10.3955$	A1
		(3)
		Total 12

- (a) B1: 52 b.p.m. Units are not required. Check for answer given in the body of the question
- (b) B1: 32 b.p.m. . Units are not required. Check for answer given in the body of the question
- (c)

M1: Achieves $\frac{dH}{dt} = P e^{-0.2t} + Q e^{-0.9t}$ where P and Q are constants

A1: $\frac{dH}{dt} = -8 e^{-0.2t} + 18 e^{-0.9t}$ but allow un-simplified versions

dM1: Sets $P e^{-0.2t} + Q e^{-0.9t} = 0 \Rightarrow m e^{0.7t} = n$ or $m e^{-0.7t} = n$ where $m \times n > 0$

M1: Solves an equation of the form $m e^{\pm kt} = n$ where $m \times n > 0$ using correct \ln work.
It must proceed from a correct form of the derivative (so the first M1).

Look for $m e^{\pm kt} = n \Rightarrow e^{\pm kt} = \frac{n}{m} \Rightarrow \pm kt = \ln \left(\frac{n}{m} \right) \Rightarrow t = \dots$

A1: $T = \text{awrt } 1.158$ minutes following the award of all previous marks. You can ignore units

.....
An alternative for the last three marks in part (c) are;

dM1: $P e^{-0.9t} = Q e^{-0.2t} \Rightarrow \ln P - 0.9t = \ln Q - 0.2t$

M1: Collects terms in t and proceeds to $t = \dots$ Under this method it is dependent upon both previous M's

A1: $T = \text{awrt } 1.158$ minutes following the award of all previous marks. Units are not required

There may be answers produced from a calculator. In cases such as this we are going to apply the rule ‘answers without full working may not gain credit.’

$$\frac{dH}{dt} = -8 e^{-0.2t} + 18 e^{-0.9t} = 0 \Rightarrow t = 1.158 \text{ scores } 1,1,0,0,0$$

(d)

M1: Sets $H = 37$ and proceeds to make $e^{\pm 0.2t}$ the subject

A1*: Proceeds to the given answer showing all necessary steps.

Note that this is a given answer so the steps must be logical without huge jumps

Acceptable proofs:

Example 1

$$\begin{aligned} 37 &= 32 + 40 e^{-0.2t} - 20 e^{-0.9t} \Rightarrow e^{-0.2t} = \frac{1 + 4 e^{-0.9t}}{8} \Rightarrow -0.2t = \ln\left(\frac{1 + 4 e^{-0.9t}}{8}\right) \\ &\Rightarrow t = -5 \ln\left(\frac{1 + 4 e^{-0.9t}}{8}\right) \\ &\Rightarrow t = 5 \ln\left(\frac{8}{1 + 4 e^{-0.9t}}\right) \end{aligned}$$

Example 2

$$37 = 32 + 40 e^{-0.2t} - 20 e^{-0.9t} \Rightarrow e^{-0.2t} = \frac{1 + 4 e^{-0.9t}}{8} \Rightarrow e^{0.2t} = \frac{8}{1 + 4 e^{-0.9t}} \Rightarrow 0.2t = \ln\left(\frac{8}{1 + 4 e^{-0.9t}}\right) \Rightarrow t = 5 \ln\left(\frac{8}{1 + 4 e^{-0.9t}}\right)$$

Unacceptable proof:

$$\begin{aligned} 37 &= 32 + 40 e^{-0.2t} - 20 e^{-0.9t} \Rightarrow e^{-0.2t} = \frac{1 + 4 e^{-0.9t}}{8} \Rightarrow -0.2t = \ln\left(\frac{1 + 4 e^{-0.9t}}{8}\right) \\ &\Rightarrow t = 5 \ln\left(\frac{8}{1 + 4 e^{-0.9t}}\right) \end{aligned}$$

They need to deal with 0.2 and –sign in separate steps

Alt (d) lns can be taken earlier. The M1 is scored for making 0.2 t the subject. See below

$$\begin{aligned} 37 &= 32 + 40 e^{-0.2t} - 20 e^{-0.9t} \\ \Rightarrow 40 e^{-0.2t} &= 5 + 20 e^{-0.9t} \Rightarrow \ln 40 - 0.2t = \ln(5 + 20 e^{-0.9t}) \Rightarrow 0.2t = \ln 40 - \ln(5 + 20 e^{-0.9t}) \end{aligned}$$

We would need to see three further steps. E.g.

$$0.2t = \ln\left(\frac{40}{5 + 20 e^{-0.9t}}\right) \Rightarrow t = 5 \ln\left(\frac{40}{5 + 20 e^{-0.9t}}\right) \Rightarrow t = 5 \ln\left(\frac{8}{1 + 4 e^{-0.9t}}\right)$$

(e)

M1: Uses the iteration formula once. Allow for the embedded 10 leading to a value or sight of $t_2 = \text{awrt } 10.4$

A1: $(t_2) = \text{awrt } 10.3947$

A1: $(M) = 10.3955$ following some evidence of previous iterations, e.g. sight of

$t_2 = 5 \ln \left(\frac{8}{1 + 4e^{-0.9 \times 10}} \right) = \text{awrt } 10.4 \dots$ This is NOT awrt so just 10.3955 here

Question Number	Scheme	Marks
9 (a)	$f'(x) = \frac{(2x+1) \times (12x+4) - 2(6x^2 + 4x - 2)}{(2x+1)^2}$	M1 A1
	$= \frac{12x^2 + 12x + 8}{(2x+1)^2} \text{ o.e.}$	A1
		(3)
(b)	At $x=2 \Rightarrow f'(x) = \frac{12 \times 2^2 + 12 \times 2 + 8}{(2 \times 2 + 1)^2} = \left(\frac{16}{5} \right)$	M1
	Full method of normal $y - 6 = -\frac{5}{16}(x - 2)$	dM1
	$16y - 96 = -5x + 10 \Rightarrow 16y + 5x = 106 \quad *$	A1*
		(3)
(c)	Any correct value $A = 3, B = \frac{1}{2} \text{ or } D = -\frac{5}{2}$	B1
	$6x^2 + 4x - 2 = (Ax + B)(2x + 1) + D \Rightarrow$ Values of A, B and D	M1
	$3x + \frac{1}{2} + \frac{-5/2}{2x+1} \text{ o.e.}$	A1
		(3)
(d)	$\int 3x + \frac{1}{2} + \frac{-5/2}{2x+1} dx = \frac{3}{2}x^2 + \frac{1}{2}x - \frac{5}{4} \ln(2x+1)$	M1, A1 ft
	Area under curve = $\left[\frac{3}{2}x^2 + \frac{1}{2}x - \frac{5}{4} \ln(2x+1) \right]_{\frac{1}{3}}^2 = \left(\frac{3}{2} \times 2^2 + \frac{1}{2} \times 2 - \frac{5}{4} \ln(5) \right) - \left(\frac{3}{2} \times \left(\frac{1}{3} \right)^2 + \frac{1}{2} \times \left(\frac{1}{3} \right) - \frac{5}{4} \ln \left(\frac{5}{3} \right) \right)$	dM1
	Area of $R = \int_{\frac{1}{3}}^2 3x + \frac{1}{2} + \frac{-5/2}{2x+1} dx + \frac{1}{2} \times 6 \times \left(\frac{106}{5} - 2 \right)$	M1
	Area of $R = \frac{964}{15} - \frac{5}{4} \ln 3$	A1
		(5)
		Total 14

(a)

M1: Condoning slips, this is for an attempt using

- the quotient rule to obtain $f'(x) = \frac{(2x+1) \times (Ax + B) - C(6x^2 + 4x - 2)}{(2x+1)^2}$ Must be ‘-’ on the numerator
- the chain rule on $3x + A + \frac{B}{2x+1} \Rightarrow f'(x) = 3 \pm D(2x+1)^{-2}$
- the product rule on $(6x^2 + 4x - 2)(2x+1)^{-1} \Rightarrow f'(x) = (2x+1)^{-1} \times (Ax + B) - C(6x^2 + 4x - 2)(2x+1)^{-2}$ where A, B and C are > 0

A1: Correct (unsimplified) expression for $f'(x)$. Look for

- $f'(x) = \frac{(2x+1) \times (12x+4) - 2(6x^2 + 4x - 2)}{(2x+1)^2}$ via the quotient rule
- $f'(x) = (2x+1)^{-1} \times (12x+4) + (6x^2 + 4x - 2) \times -2(2x+1)^{-2}$ via the product rule
- $3 + 5(2x+1)^{-2}$ following division. Condone an incorrect value for A.

A1: Simplified answers such as $\frac{4(3x^2 + 3x + 2)}{(2x+1)^2}$, $\frac{12x^2 + 12x + 8}{4x^2 + 4x + 1}$ or $3 + 5(2x+1)^{-2}$ following correct A.

ISW following a correct answer

(b)

M1: Substitutes $x = 2$ into their attempt at $f'(x)$.

If $f'(x)$ is incorrect then look for correctly embedded 2's or a correct value.

dM1: Full method for finding the equation of the normal at $(2, 6)$. Look for $y - 6 = -\frac{1}{f'(2)}(x - 2)$

A1*: $y - 6 = -\frac{5}{16}(x - 2)$ o.e with at least one correct intermediate line leading to $16y + 5x = 106$

If $y = mx + c$ is used look for $6 = -\frac{5}{16} \times 2 + c \Rightarrow c = \frac{106}{16}$ You can allow $y = -\frac{5}{16}x + \frac{106}{16}$ going straight to the given answer. ISW following sight of correct answer. It must follow a correct derivative in (a)

(c)

B1: Any correct value for $A = 3$, $B = \frac{1}{2}$ or $D = -\frac{5}{2}$.

They may be embedded within an expression e.g. $3x$ in the quotient of a division sum

M1: Correct method for finding all three constants.

Via comparison/inspection look for $6x^2 + 4x - 2 = (Ax + B)(2x + 1) + D \Rightarrow$ Values of A , B and D

Via division look for a linear quotient and a constant remainder

A1: $3x + \frac{1}{2} + \frac{-5/2}{2x+1}$ o.e such as $3x + \frac{1}{2} - \frac{5}{2(2x+1)}$ which may be seen within an integral in (d).

It is not just for values of A , B and D .

ISW following a correct answer (e.g. they may go on to $\times 2$ but then the A1 fit in (d) would be unavailable)

(d)

M1: Attempts to integrate their $Ax + B + \frac{D}{2x+1}$. Look for two terms in the correct form, one of which must

be the ln term. Award for $\dots x^2 + \dots + k \ln(2x+1)$ or $\dots + \dots x + k \ln(2x+1)$

Also note that $k \ln(4x+2)$ and any multiple of $(2x+1)$ within the ln is also correct

A1ft: $\frac{3}{2}x^2 + \frac{1}{2}x - \frac{5}{4} \ln(2x+1)$ o.e. but follow through on their values of A, B and D

dM1: A correct method of finding the area under the curve using limits of $\frac{1}{3}$ and 2 either way around.

Dependent upon the previous M mark. Award if the intention is clear.

M1: For a correct method of finding the area of R . It is scored for adding the area under the curve to the correct calculation for the area of the triangle. The integration of $f(x)$ need not be correct but the limits must be applied the correct way around. The area of the triangle may be done by integration. Look for

$$\int_{\frac{1}{3}}^2 f(x) dx + \frac{1}{2} \times 6 \times \left(\frac{106}{5} - 2 \right) = [I(x)]_{\frac{1}{3}}^2 + \frac{1}{2} \times 6 \times \left(\frac{106}{5} - 2 \right) = I(2) - I\left(\frac{1}{3}\right) + \frac{1}{2} \times 6 \times \left(\frac{106}{5} - 2 \right) = \dots$$

where $I(x)$ is their attempt at $\int f(x) dx$

A1: $\frac{964}{15} - \frac{5}{4} \ln 3$

