

Write your name here	
Surname	Other names
Pearson Edexcel	Centre Number
International GCSE	Candidate Number
<h1 style="margin: 0;">Further Pure Mathematics</h1> <h2 style="margin: 0;">Paper 2</h2>	
Thursday 23 January 2014 – Morning	Paper Reference
Time: 2 hours	4PM0/02
Calculators may be used.	Total Marks

Instructions

- Use **black** ink or ball-point pen.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions.
- Without sufficient working, correct answers may be awarded no marks.
- Answer the questions in the spaces provided
– *there may be more space than you need.*

Information

- The total mark for this paper is 100.
- The marks for **each** question are shown in brackets
– *use this as a guide as to how much time to spend on each question.*

Advice

- Read each question carefully before you start to answer it.
- Check your answers if you have time at the end.

Turn over ►

P44027RA

©2014 Pearson Education Ltd.

6/6/1/1/1/



PEARSON

- 2 The volume of a right circular cone is increasing at a constant rate of $12 \text{ cm}^3/\text{s}$. The radius of the base of the cone is always half the height of the cone. Find, in cm/s , the exact value of the rate of increase of the height of the cone when the height is 4 cm.

(5)

(Total for Question 2 is 5 marks)



4




Diagram **NOT** accurately drawn

Figure 1

Figure 1 shows a sector of a circle of radius 10 cm and centre O . The area of triangle OAB is 20 cm^2 and the size of angle AOB is θ radians.

Find, to 3 significant figures,

- (a) the value of θ , (2)

- (b) the length of the arc AB , (2)

- (c) the area of the shaded segment. (3)

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

6

Question 7 continued

A large rectangular area containing 25 horizontal dotted lines for writing answers.



8




Figure 4

Figure 4 shows a hexagon $OABCDE$. Each internal angle of the hexagon is 120° .

$$OA = OE, AB = ED = 2 \times OA \text{ and } OC = 3 \times OA$$

$$\vec{OA} = \mathbf{a} \text{ and } \vec{OE} = \mathbf{e}.$$

Find as simplified expressions in terms of \mathbf{a} and \mathbf{e}

(a) \vec{AB} , (2)

(b) \vec{BE} . (2)

The point P divides AB internally in the ratio 2:3

(c) Find \vec{PC} as a simplified expression in terms of \mathbf{a} and \mathbf{e} . (3)

The point Q lies on ED produced so that the points P , C and Q are collinear.

(d) Find \vec{OQ} in the form $\lambda\mathbf{a} + \mu\mathbf{e}$, stating the value of λ and the value of μ . (6)

.....

.....

.....

.....

.....

.....

.....

10 The sum of the second and third terms of a convergent geometric series is 7.5

The sum to infinity, S , of the series is 20

The common ratio of the series is r .

(a) Show that r is a root of the equation

$$8r^3 - 8r + 3 = 0 \quad (4)$$

(b) Show that $r = \frac{1}{2}$ is a root of this equation. (1)

Given that $r < 0.6$

(c) show that $\frac{1}{2}$ is the only possible value of r . (4)

(d) Find the first term of the series. (2)

The sum of the first n terms of the series is S_n

(e) Find the least value of n for which S_n exceeds 99% of S . (6)



