Please check the examination details below before entering your candidate information

Candidate surname	Other names

Centre Number | Candidate Number

Pearson Edexcel International GCSE

Tuesday 20 May 2025

Afternoon (Time: 2 hours)

Paper reference **4PM1/01**

Further Pure Mathematics

PAPER 1

Calculators may be used.

Total Marks

Instructions

- Use **black** ink or ball-point pen.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions.
- Without sufficient working, correct answers may be awarded no marks.
- Answer the questions in the spaces provided
  – *there may be more space than you need.*
- You must **NOT** write anything on the formulae page.
  Anything you write on the formulae page will gain NO credit.

Information

- The total mark for this paper is 100.
- The marks for **each** question are shown in brackets
  – *use this as a guide as to how much time to spend on each question.*

Advice

- Read each question carefully before you start to answer it.
- Check your answers if you have time at the end.

Turn over ▶

P72865A

Y:1/1/1/1/1/

P 7 2 8 6 5 A 0 1 3 6

Pearson

International GCSE in Further Pure Mathematics Formulae sheet

Mensuration

Surface area of sphere = $4\pi r^2$

Curved surface area of cone = $\pi r \times$ slant height

Volume of sphere = $\frac{4}{3}\pi r^3$

Series

Arithmetic series

Sum to n terms, $S_n = \frac{n}{2}[2a + (n-1)d]$

Geometric series

Sum to n terms, $S_n = \frac{a(1-r^n)}{(1-r)}$

Sum to infinity, $S_\infty = \frac{a}{1-r} \quad |r| < 1$

Binomial series

$$(1+x)^n = 1 + nx + \frac{n(n-1)}{2!}x^2 + \ldots + \frac{n(n-1)\ldots(n-r+1)}{r!}x^r + \ldots \quad \text{for} \quad |x| < 1, n \in \mathbb{Q}$$

Calculus

Quotient rule (differentiation)

$$\frac{\mathrm{d}}{\mathrm{d}x}\left(\frac{\mathrm{f}(x)}{\mathrm{g}(x)}\right) = \frac{\mathrm{f}'(x)\mathrm{g}(x) - \mathrm{f}(x)\mathrm{g}'(x)}{[\mathrm{g}(x)]^2}$$

Trigonometry

Cosine rule

In triangle ABC: $a^2 = b^2 + c^2 - 2bc\cos A$

$\tan\theta = \frac{\sin\theta}{\cos\theta}$

$\sin(A+B) = \sin A \cos B + \cos A \sin B$ $\quad$ $\sin(A-B) = \sin A \cos B - \cos A \sin B$

$\cos(A+B) = \cos A \cos B - \sin A \sin B$ $\quad$ $\cos(A-B) = \cos A \cos B + \sin A \sin B$

$\tan(A+B) = \frac{\tan A + \tan B}{1 - \tan A \tan B}$ $\quad$ $\tan(A-B) = \frac{\tan A - \tan B}{1 + \tan A \tan B}$

Logarithms

$\log_a x = \frac{\log_b x}{\log_b a}$

There are no questions on this page.

Answer all ELEVEN questions.

Write your answers in the spaces provided.

You must write down all the stages in your working.

1 A particle P moves in a straight line. At time t seconds, the velocity, v m/s, of P is given by

$$v = 3t^2 - 9t + 7$$

(a) Show that P never comes to rest.

(2)

(b) Find the acceleration of P, in m/s^2, when $t = 4$

(2)

Using algebra

(c) find the distance, in m, that P travels in the interval $0 \leqslant t \leqslant 5$

(3)

Question 1 continued

(Total for Question 1 is 7 marks)

2 The point A has coordinates $(3, 2)$, the point B has coordinates $(8, 3)$ and the point C has coordinates $(4, 7)$

(a) Show that ABC is an isosceles triangle.

(2)

The midpoint of BC is M

(b) Find an equation of the line that passes through A and M

Give your answer in the form $y = mx + c$

(3)

The points A, C and D are collinear such that $AD = kAC \quad (k > 1)$

Given that $\angle ABD = 90^o$

(c) find the coordinates of D
Show your working clearly.

(8)

Question 2 continued

Question 2 continued

Question 2 continued

(Total for Question 2 is 13 marks)

3



Diagram **NOT** accurately drawn

Figure 1

Figure 1 shows circle C_1 with radius 7 cm and circle C_2 with centre O and radius 7 cm

The circles C_1 and C_2 intersect at the point A and at the point B

The size of angle AOB is θ radians

The perimeter of the region R shown shaded in Figure 1 is $\frac{28\pi}{9}$ cm

(a) Find the exact value of θ

(2)

(b) Find the area, in cm^2 to 3 significant figures, of the region R

(4)

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 3 continued

(Total for Question 3 is 6 marks)

4 (a) On the grid opposite, draw the graph of the line with equation

(i) $3x + 2y = 18$ (ii) $x + 3y - 6 = 0$ (iii) $y = 3x$

(3)

(b) Show, by shading on the grid, the region R defined by the inequalities

$3x + 2y \leqslant 18$ $x + 3y - 6 \geqslant 0$ $y \leqslant 3x$

(1)

For all points in R, with coordinates (x, y)

$$P = 2x - y$$

(c) find the least value of P

(4)

Question 4 continued



Turn over for a spare grid if you need to redraw your graph.

Question 4 continued

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 4 continued

Only use this grid if you need to redraw your graph.



(Total for Question 4 is 8 marks)

5 (a) Show that $(\alpha+\beta)^3-3\alpha\beta(\alpha+\beta)=\alpha^3+\beta^3$

(2)

The quadratic equation $2x^2-6x-7=0$ has roots α and β

Without solving the equation

(b) form a quadratic equation, with integer coefficients, which has roots $\frac{\alpha^2}{\beta}$ and $\frac{\beta^2}{\alpha}$

(6)

Question 5 continued

(Total for Question 5 is 8 marks)

6 (a) Show that $\sqrt{4-x} = A\left(1-\frac{x}{B}\right)^{\frac{1}{2}}$ where A and B are integers to be found.

(2)

(b) Hence expand $\sqrt{4-x}$ in ascending powers of x up to and including the term in x^3
Give each coefficient as an exact fraction in its lowest terms.

(3)

(c) Use your expansion with a suitable value of x to obtain an estimate of $\frac{\sqrt{305}}{10}$

Give your answer correct to 5 decimal places.

(3)

Question 6 continued

(Total for Question 6 is 8 marks)

7



Diagram **NOT** accurately drawn

Figure 2

Figure 2 shows a hollow hemisphere with radius 20 cm

The hemisphere contains liquid, which is dripping out of a small hole at the lowest point A at a constant rate of k cm^3/s

At time t seconds after the liquid starts to drip from the hemisphere, the height of the liquid is h cm above A

The volume V cm^3 of liquid in the hemisphere is given by

$$V = \frac{\pi}{3}h^2(60 - h)$$

When $h = 12$, the height of the liquid is decreasing at a rate of $\frac{1}{60}$ cm/s

Find the value of k
Give your answer in terms of π

(6)

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 7 continued

Question 7 continued

Question 7 continued

(Total for Question 7 is 6 marks)

8



Diagram **NOT** accurately drawn

Figure 3

Figure 3 shows a right pyramid with vertex V and square base, $ABCD$, of side 12 cm

The size of the angle between the plane VAB and the base $ABCD$ is $30°$

(a) Show that the height of the pyramid is $2\sqrt{3}$ cm

(2)

(b) Find, in cm, the exact length of VA

(3)

(c) Find, in cm^2, the exact area of triangle VAD

(3)

(d) Find, in cm, the exact length of the perpendicular from D to VA

(2)

(e) Find, in degrees to one decimal place, the size of the obtuse angle between the plane VAB and the plane VAD

(3)

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Question 8 continued

Question 8 continued

Question 8 continued

(Total for Question 8 is 13 marks)

9



Figure 4

Figure 4 shows a sketch of part of the curve C with equation $y = \mathrm{f}(x)$ where

$$\mathrm{f}(x) = 2x^3 + ax^2 + bx + c$$

The curve C has a maximum at the point A with coordinates $\left(-\frac{1}{3}, \frac{100}{27}\right)$ and a minimum at the point B with coordinates $(2, -9)$

Given that a, b and c are integers

(a) show that $a = -5, b = -4$ and $c = 3$

(5)

(b) (i) Show that $(x+1)$ is a factor of $\mathrm{f}(x)$

(1)

(ii) Hence, or otherwise, use algebra to factorise $\mathrm{f}(x)$ completely.

(3)

The curve C crosses the x-axis at the points M, N and P
The finite regions shown shaded in Figure 4 are bounded by the curve C and parts of the x-axis from M to N and from N to P

(c) Use algebraic integration to determine the total area of the shaded regions. Give your answer as an exact fraction.

(4)

Question 9 continued

Question 9 continued

Question 9 continued

(Total for Question 9 is 13 marks)

10 Solve the equation

(i) $\log_4(6y-5)=3$

(2)

(ii) $\log_4(4-3x)^2-\log_2(x^2-5)-3=0$
Show clear algebraic working.
Give your answer to 3 significant figures.

(9)

Question 10 continued

(Total for Question 10 is 11 marks)

11



Figure 5

Figure 5 shows part of the curve C with equation $y = \sqrt{4x - 8}$ and the line l with equation $x = b$ where $b > 0$

The finite region bounded by the curve C, the x-axis and the line l, shown shaded in Figure 5, is rotated through $360°$ about the x-axis.

Given that the volume of the solid formed is 50π units^3

find the value of b

(7)

Question 11 continued

Question 11 continued

(Total for Question 11 is 7 marks)

TOTAL FOR PAPER IS 100 MARKS