



Pearson

# Mark Scheme (Results)

## Summer 2018

Pearson Edexcel International GCSE  
In Mathematics B (4MB1)  
Paper 01R

## Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at [www.edexcel.com](http://www.edexcel.com) or [www.btec.co.uk](http://www.btec.co.uk). Alternatively, you can get in touch with us using the details on our contact us page at [www.edexcel.com/contactus](http://www.edexcel.com/contactus).

## Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: [www.pearson.com/uk](http://www.pearson.com/uk)

Summer 2018

Publications Code 4MB1\_01R\_1806\_MS

All the material in this publication is copyright

© Pearson Education Ltd 2018

## General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.
- **Types of mark**
  - M marks: method marks
  - A marks: accuracy marks
  - B marks: unconditional accuracy marks (independent of M marks)
- **Abbreviations**
  - cao – correct answer only
  - ft – follow through
  - isw – ignore subsequent working
  - SC - special case
  - oe – or equivalent (and appropriate)
  - dep – dependent
  - indep – independent
  - awrt – answer which rounds to
  - eoo – each error or omission

- **No working**

If no working is shown then correct answers normally score full marks

If no working is shown then incorrect (even though nearly correct) answers score no marks.
- **With working**

If there is a wrong answer indicated on the answer line always check the working in the body of the script (and on any diagrams), and award any marks appropriate from the mark scheme.

If it is clear from the working that the “correct” answer has been obtained from incorrect working, award 0 marks.

Any case of suspected misread loses A (and B) marks on that part, but can gain the M marks.

If working is crossed out and still legible, then it should be given any appropriate marks, as long as it has not been replaced by alternative work.

If there is a choice of methods shown, then no marks should be awarded, unless the answer on the answer line makes clear the method that has been used.

If there is no answer on the answer line then check the working for an obvious answer.
- **Ignoring subsequent work**

It is appropriate to ignore subsequent work when the additional work does not change the answer in a way that is inappropriate for the question: eg. Incorrect cancelling of a fraction that would otherwise be correct.

It is not appropriate to ignore subsequent work when the additional work essentially makes the answer incorrect eg algebra.


Transcription errors occur when candidates present a correct answer in working, and write it incorrectly on the answer line; mark the correct answer.
- **Parts of questions**

Unless allowed by the mark scheme, the marks allocated to one part of the question CANNOT be awarded in another part.

Question	Working	Answer	Mark	Notes	Sub-Total	Total
1	$\frac{23}{8} \times \frac{8}{5}$		M1	Need to see $\frac{23}{8} \times \frac{8}{5}$ and $\frac{23}{5}$ or $\frac{184}{40}$		2
		$4\frac{3}{5}$	A1	<b>NB</b> no marks for an answer without any working. Must be the mixed fraction in its simplest form		
2	$360 \div 12$ or $180(n-2) = 168n$ oe		M1	$360 \div (180-168)$ <b>NB</b> $180(n-2) = 168$ is M0		2
		30	A1			
3	$(1.7 \times 10^7) \div (1.5 \times 10^3)$		M1	for $1.1(3) \times 10^n$ <b>or</b> correct value to 2 or more significant figures. Eg 11333.33... 11000, 11300		2
		$1.1 \times 10^4$	A1	cao $1.1 \times 10^4$		
4	$-3 \times 7x^{-4} - 5x^4$		M1	for one correct term (allow $-3 \times 7x^{-4}$ )		2
		$-21x^{-4} - 5x^4$	A1	oe e.g. $-\frac{21}{x^4} - 5x^4$		
5	$\frac{12}{3-\sqrt{5}} \times \frac{3+\sqrt{5}}{3+\sqrt{5}}$ or $12 = 3a + 3b\sqrt{5} - a\sqrt{5} - 5b$ and $3a - 5b = 12, 3b - a = 0$ oe		M1			2
	$\frac{36+12\sqrt{5}}{9-5}$ or $4b = 12$ or $4a = 36$	$9+3\sqrt{5}$	A1	Correct expansion/correct method for solving simultaneous equations <b>with</b> a correct answer and no errors. <b>NB</b> no marks for answer without any working.		



Question	Working	Answer	Mark	Notes	Sub-Total	Total
10	19.45 or 19.35 or 2.35 or 2.45		B1			3
	$(b =) 19.45 - 2 \times 2.35$		M1	Or for $UB_1 - 2 \times LB_2$ or $UB_1 = 2 \times LB_2 + b$ where $19.4 < UB_1 \leq 19.5$ & $2.3 \leq LB_2 < 2.4$		
		14.75	A1			
11	$3(x^3 + a) = 4(c - x^3)$ oe		M1			3
	$3x^3 + 4x^3 = 4c - 3a$ or $3a - 4c = -4x^3 - 3x^3$		M1	Collecting $x$ terms on one side and other terms on the opposite side oe. Do not ISW		
		$\sqrt[3]{\frac{4c-3a}{7}}$	A1	<b>NB</b> A0 for $\pm\sqrt[3]{\frac{4c-3a}{7}}$ $3\sqrt[3]{\frac{4c-3a}{7}}$		
12	$5^{3k+4} = 125$		M1	Allow $\frac{750}{6}$		3
	$3k + 4 = 3$		M1	Dep first M1 Writing "125" as a power of 5 and equating powers, 0.33(0.33...)		
		$-\frac{1}{3}$	A1	cao		
13	$\left[\frac{BE^2}{9.6^2} = \right] \left(\frac{9}{16}\right) \text{ or } \left(\frac{27}{21+27}\right)$ oe		M1	For $\frac{9}{16}$ or $\frac{27}{21+27}$ <b>Alternate</b> $h = 10$ , $0.5BE \times x = 27$		3
	$[BE =] \sqrt{\frac{9}{16}} \times 9.6$		M1	<b>Alternate</b> $(9.6 + BE)(10 - x) = 42$		
		7.2	A1			

Question	Working	Answer	Mark	Notes	Sub-Total	Total
14 (a)(i) (ii) (iii)	$y = 2$		B1	correct line	1	
	$x + y = 5$		B1	correct line (condone incorrect labelling)	1	
	$y = 2x + 1$		B1	correct line	1	
(b)		<b>R</b> correctly placed	B1	Do not award if lines incorrect Ignore labelling of lines	1	4
15	$\frac{1}{5} \times \left( \frac{120}{5} \times 3 \right) (=14.4(0))$		M1	or (Barry:) $\frac{3}{5} \times \frac{1}{5} (= \frac{3}{25})$		
	$0.35 \times \left( \frac{120}{5} \times 2 \right) (=16.8(0))$		M1	or (Carlos:) $\frac{35}{100} \times \frac{2}{5} (= \frac{14}{100} = \frac{7}{50})$		
	$\frac{'14.4'+ '16.8'}{120} = \frac{'31.2'}{120}$		M1	Dep on M2 or for $'\frac{3}{25}'+ '\frac{7}{50}'$		
		$\frac{13}{50}$ or 0.26	A1			

Question	Working	Answer	Mark	Notes	Sub-Total	Total
16 (a)		$6w^5y^8$	B2	B1 for 2 terms correct as part of a product. Do not ISW	2	4
(b)		$3a^2c$	B2	B1 for 2 terms correct as part of a product, allow $3a^2c^1$ . Do not ISW	2	
17	$OBA = 52^\circ$		M1	may be marked on diagram		4
	$AOB = 76^\circ$ or $BAC = 128^\circ$		M1	may be marked on diagram must be identified as correct angles		
		14	A1			
	e.g. angle between <b>tangent</b> and <b>radius</b> = $90^\circ$ base angles/radii equal / isosceles triangle <b>Angle sum</b> of <b>triangle</b> Angle sum of <b>triangle</b> = <b>180</b> <b>Angle sum</b> of <b>straight line</b> Angle sum of <b>straight line</b> = <b>180</b>		B1	for 2 correct reasons for method used		
18 (a)	$\begin{pmatrix} -4 \\ 2 \end{pmatrix} + \begin{pmatrix} -2 \\ 6 \end{pmatrix}$ or $\begin{pmatrix} -2 \\ 6 \end{pmatrix} - \begin{pmatrix} 4 \\ -2 \end{pmatrix}$	$\begin{pmatrix} -6 \\ 8 \end{pmatrix}$	M1	oe	2	
			A1			
(b)	$\sqrt{(-6)^2 + 8^2}$		M1ft	ft part(a). Condone missing minus.	2	4
		10	A1ft	ft part (a)		

Question	Working	Answer	Mark	Notes	Sub-Total	Total
19	$(3x+2) \times \frac{5}{3x^2-7x-6} \left[ -\frac{5}{x+3} \right]$		M1	For $\times$ by reciprocal condone missing bracket round $3x+2$		
	$(3x+2) \times \frac{5}{(3x+2)(x-3)} \left[ -\frac{5}{x+3} \right]$		M1	Factorising correctly		
	$\frac{5(x+3)-5(x-3)}{(x-3)(x+3)}$		M1	Correct method for combining into a single fraction		
	$\frac{5x+15-5x+15}{(x+3)(x-3)}$					
		$\frac{30}{x^2-9}$	A1	or $\frac{30}{(x+3)(x-3)}$		
20	$\overrightarrow{AP} = -\mathbf{a} + \frac{5}{6}(\mathbf{a} + 3\mathbf{b}) \left[ = -\frac{1}{6}\mathbf{a} + \frac{5}{2}\mathbf{b} \right]$		M1	For correct vector for $\overrightarrow{AP}$		
	$\overrightarrow{AD} = -\mathbf{a} + n\mathbf{b}$ or $-\mathbf{a} + (5+n)\mathbf{b}$		M1	indep allow $\overrightarrow{OD} = \mathbf{a} + n\overrightarrow{AP}$		
	$\overrightarrow{AD} = 6\left(-\frac{1}{6}\mathbf{a} + \frac{5}{2}\mathbf{b}\right) \left[ = -\mathbf{a} + 15\mathbf{b} \right]$		M1	or $AD = 6AP$ or $1 - \frac{1}{6}n = 0$ and $\overrightarrow{OD} = 15\mathbf{b}$		
	$OB : OD = 5 : 15$	$1 : 3$	A1	Seeing $5 : 15$ or $5\mathbf{b} : 15\mathbf{b}$ equals $1 : 3$ from correct working		

Question	Working	Answer	Mark	Notes	Sub-Total	Total
21	$\sqrt{8^2 + 15^2} (=17)$		M1	Using Pythagoras correctly		
	$10 \times 9 + 18 \times 9 + 15 \times 9$		M1	correct areas of the 3 rectangles		
	$\frac{18+10}{2} \times 15$ or $10 \times 15 + \frac{8 \times 15}{2}$ [=210]		M1	Attempt at area of trapezium		
	$2 \times "210" + 10 \times 9 + 18 \times 9 + 15 \times 9 + "17" \times 9$		M1	dep on previous method marks – for adding the six areas together		
		960	A1			
22 (a)	$[T =] \frac{k}{y^2}$		M1	For $\frac{k}{y^2}$		
	$0.32 = \frac{k}{5^2}$		M1	Subst 0.32 for $T$ and 5 for $y$		
		$T = \frac{8}{y^2}$	A1	<b>NB</b> SCB1 for $0.32 = \frac{k}{\sqrt{5}}$		
(b)	$200 = \frac{"8"}{y^2}$		M1			
		0.2	A1	oe		

Question	Working	Answer	Mark	Notes	Sub-Total	Total
<b>23</b> (a)		$(x+5)^2 - 32$	B2	or for $p = 5$ and $q = -32$ B1 for $(x + 5)^2$ , B1 for $-32$	2	5
(b)	$(x + '5')^2 = '32'$		M1	ft from (a)		
	$x + '5' = \pm\sqrt{'32'}$		M1	ft		
		$-5 \pm \sqrt{32}$	A1	$-5 \pm 4\sqrt{2}$ gets A0 SCB1 for use of formula with correct answers although $-5 \pm 4\sqrt{2}$ is B0	3	
<b>24</b> (a)	$(-2)^3 - 3 \times (-2)^2 - 2a + 12 (=0)$ Or $(-2)^3 - 3 \times (-2)^2 - 2 \times -4 + 12$ $2a = -8$ or $-2a = 8$ or $2a = -20 + 12$ , $a = -4$ Or $-8 - 12 + 8 + 12 = 0$ so $a = -4$		M1		2	
			A1	no working gains zero marks		
(b)	$(x + 2)(x^2 - 5x + 6)$ or $x^2(x - 3) - 4(x - 3)$		M1	Allow a sign error		5
	$x^2 - 5x + 6 = (x - 2)(x - 3)$		M1	For factorising any 3 term quadratic which when expanded, the result gives at least 2 of the 3 terms from their trinomial, e.g. $(x - 6)(x - 1)(=0)$ will give $x^2$ and $+6$ terms or $(x^2 - 4)(x - 3)$ Indep of previous M mark		
		$(x+2)(x-3)(x-2)$	A1		3	

Question	Working	Answer	Mark	Notes	Sub-Total	Total
25 (a)	$1 - (0.15+0.13+0.2+0.32)$		M1		2	
		0.2	A1			
(b)	$\frac{0.32}{0.13} \times 39$ or $0.32 \times (39 \div 0.13)$ or $0.32 \times 300$		M1	oe	2	
		96	A1			
(c)	$0.15 \times 360$		M1	oe	2	6
		54	A1			
26(a)		$\begin{pmatrix} 16 & 3 \\ -4 & 11 \end{pmatrix}$	B2	-1eeoo	2	
(b)		$\begin{pmatrix} 11 & 0 \\ -3 & 10 \end{pmatrix}$	B2	-1eeoo	2	
(c)		$(12 \quad -7)$	B1	for matrix of correct order or for 12 and -7 seen. Allow $10 + 2$ , $-15 + 8$ or $5 \times 2 + 2 \times 1$ and $5 \times -3 + 2 \times 4$	2	6
			B1	fully correct including brackets		

Question	Working	Answer	Mark	Notes	Sub-Total	Total
27	$\angle EOD = \frac{360 \times 3.5\pi}{12\pi} \text{ oe}(= 105)$		M1	method to find angle $EOD$		
	$ED = \sqrt{6^2 + 6^2 - 2 \times 6 \times 6 \times \cos 105}$ or $ED = 2 \times 6 \sin 52.5 \text{ oe } (=9.52) \text{ oe}$		M1	method to find length $ED$		
	$DC \times (DC + "9.52") = 9 \times 16$		M1	correct equation for $DC$ Allow 9.5 or better for $ED$		
	$DC^2 + 9.52DC - 144 (= 0) \text{ oe}$		M1	Dep on previous M mark. correct equation in form to solve		
	$DC = \frac{-9.52 + \sqrt{9.52^2 - 4 \times 1 \times -144}}{2 \times 1}$		M1	Dep on previous two M marks fully correct method to find $DC$		
		8.15	A1			