



Mark Scheme (Results)

Summer 2019

Pearson Edexcel International GCSE
In Mathematics B (4MB1)
Paper 01R

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

Summer 2019

Publications Code 4MB1_01R_2019_MS

All the material in this publication is copyright

© Pearson Education Ltd 2019

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme.

Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.

- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

- **Types of mark**

- o M marks: method marks
- o A marks: accuracy marks
- o B marks: unconditional accuracy marks (independent of M marks)

- **Abbreviations**

- o cao – correct answer only
- o ft – follow through
- o isw – ignore subsequent working
- o SC - special case
- o oe – or equivalent (and appropriate)

- o dep – dependent
- o indep – independent
- o awrt – answer which rounds to
- o eooo – each error or omission

- **No working**

If no working is shown then correct answers normally score full marks

If no working is shown then incorrect (even though nearly correct) answers score no marks.

- **With working**

If there is a wrong answer indicated on the answer line always check the working in the body of the script (and on any diagrams), and award any marks appropriate from the mark scheme.

If it is clear from the working that the “correct” answer has been obtained from incorrect working, award 0 marks.

If a candidate misreads a number from the question. Eg. Uses 252 instead of 255; method marks may be awarded provided the question has not been simplified. Examiners should send any instance of a suspected misread to review. If there is a choice of methods shown, then no marks should be awarded, unless the answer on the answer line makes clear the method that has been used.

If there is no answer on the answer line then check the working for an obvious answer.

- **Ignoring subsequent work**

It is appropriate to ignore subsequent work when the additional work does not change the answer in a way that is inappropriate for the question: eg. Incorrect cancelling of a fraction that would otherwise be correct.

It is not appropriate to ignore subsequent work when the additional work essentially makes the answer incorrect eg algebra. Transcription errors occur when candidates present a correct answer in working, and write it incorrectly on the answer line; mark the correct answer.

- **Parts of questions**

Unless allowed by the mark scheme, the marks allocated to one part of the question CANNOT be awarded to another.

Question		Working	Answer	Mark	Notes																								
1		60, 120, 180, 240, 300, 360, 420, 480, 540, ... 135, 270, 405, 540, ... or 60 = 2×2×3×5 or 15×2×2 135 = 3×3×3×5 or 15×3×3 or <table border="1"><tr><td>5</td><td>60</td><td>135</td></tr><tr><td>3</td><td>12</td><td>27</td></tr><tr><td></td><td>4</td><td>9</td></tr></table>	5	60	135	3	12	27		4	9			M1 for a correct list of multiples up to 540 or 60 and 135 written as a correct product of primes - factors may be on ends of trees or in ladder diagrams (so expect to see 3, 3, 3, 4 and 5 or equivalent e.g. 3, 4, 5, 9) or correct factor grid The following is common: <table><tr><td>5</td><td>60</td><td>135</td></tr><tr><td>3</td><td>12</td><td>27</td></tr><tr><td>4</td><td>4</td><td>9</td></tr><tr><td>9</td><td>1</td><td>9</td></tr><tr><td></td><td>1</td><td>1</td></tr></table>	5	60	135	3	12	27	4	4	9	9	1	9		1	1
5	60	135																											
3	12	27																											
	4	9																											
5	60	135																											
3	12	27																											
4	4	9																											
9	1	9																											
	1	1																											
			540	2	A1																								
					Total 2 marks																								
2		9n – 7 = 214			M1 or for working out the 24 th and 25 th terms																								
			No and n = 24.55... or 9n = 221 and 221 is not a multiple of 9 or 24th term = 209, 25th term = 218	2	A1 oe (e.g. No with either 24.5 or 24.6 (or better)) – for A1 must see ‘No’ + appropriate values A0 if No and $n = \frac{221}{9}$ only For A1 it must be explicitly clear that $n = \frac{221}{9}$ is not an integer e.g. No and $n = \frac{221}{9}$ is not an integer/whole number is A1																								
					Total 2 marks																								

3					M1 for $24 - 18 (= 6)$ or $\frac{7+x}{2} = 10$ oe or one value correctly stated
			$w = 6$ $x = 13$	2	A1 SC B1 for $w = 13, x = 6$
					Total 2 marks
4		$\pi \times 8^2 \times 12$			M1 (or for 768π)
			2413	2	A1 accept answer in the range 2412 – 2413
					Total 2 marks
5		$2x = 5 \times 9 + 3$ or $\frac{2x}{5} = 9 + \frac{3}{5}$			M1 (oe)
			24	2	A1
					Total 2 marks
6		$(-6)^2 - 4 \times -6$ oe e.g. $36 - - 24$ or $-6(-6 - 4)$			M1 (must include a bracket around the -6 unless recovered later) or for $(+)36$ or $(+)24$ seen in their working M0 if -12 seen with no working M0 if -12 comes from $-6^2 - 4 \times -6$ without seeing $(+)24$
			60	2	A1
					Total 2 marks
7		$\frac{11}{4} \times \frac{12}{11}$ or $\frac{33}{12} \div \frac{11}{12} = \frac{33}{11}$			M1
		$\frac{132}{44} = 3$ or $\frac{1}{4} \times \frac{12}{1} = \frac{12}{4} = 3$ or $\frac{1}{1} \times \frac{3}{1} = 3$ oe (cancelling of 11s and 4 and 12 seen) or $\frac{33}{12} \div \frac{11}{12} = \frac{33}{11} = 3$ or $\frac{11}{4} \times \frac{12}{11} = \frac{12}{4} = 3$	3	2	A1 dependent on all working seen $\frac{11}{4} \times \frac{12}{11} = 3$ or $\frac{33}{12} \times \frac{12}{11} = 3$ is A0 unless explicit cancelling seen
					Total 2 marks
8					M1 for $ABC = 48^\circ$ or reflex $AOC = 264^\circ$ stated or marked on diagram
			96	2	A1
					Total 2 marks

9					M1 for $12x^2$ or $14x^{-3}$ oe (e.g. $\frac{14}{x^3}$)
			$12x^2 + 14x^{-3}$	2	A1 oe (e.g. $12x^2 + \frac{14}{x^3}$) must be + in between the two terms – mark final answer
					Total 2 marks
10					M1 for writing $\sqrt{100a^2} + a\sqrt{25a^2}$ or $\sqrt{20a} = 2\sqrt{5a}$ or for $10a$ or $5a^2$
			$10a + 5a^2$	2	A1 or equivalent e.g. $5a(2 + a)$ but A0 if correct answer replaced with $2a + a^2$ (or other incorrect simplification)
					Total 2 marks
11		$(7 \times 5) \div 4$ or $5 \times 7 = 4 \times CP$ oe e.g. $4CP = 35$, $\frac{5}{4} \times 7 = CP$ etc			M1
			8.75	2	A1
					Total 2 marks
12			$x \geq 1$		B1oe accept use of $>$ oe (but B0 for $1 \leq x \leq 4$)
			$y \geq \frac{1}{2}x$		B1oe accept use of $>$ oe (e.g. $2y \geq x$ or $y - \frac{1}{2}x \geq 0$ etc.)
			$x + y \leq 6$	3	B1oe accept use of $>$ oe (e.g. $y \leq 6 - x$ etc.) - condone for full marks inconsistent use of $>$ or $\geq$
					Total 3 marks

13		$0.2 \times 8600 (=1720) \text{ or } 0.8 \times 8600 (=6880)$			M1 – award M1 only for 5848 (from $8600 \times 0.8 \times 0.85$) M0 for 4300 (reducing by 50%) unless first year explicitly seen (e.g. either 1720 or 6880 seen in calculation)
		$0.85^2 \times "6880"$			M1ft sight of multiplying their '6880' (which cannot be 8600 but allow for say 1720 or a different changed value) by 0.85^2 – so a correct method for years 2 and 3 following on from their possibly incorrect year 1) or M2 for $0.8 \times 0.85^2 \times 8600$
			4970.8(0) or 4971	3	A1 - award M2 only for 4970
					Total 3 marks
14		$\begin{pmatrix} 20 & 15 \\ 10 & -5 \end{pmatrix} + n \begin{pmatrix} 4 & x \\ 2y & 7 \end{pmatrix} \text{ oe}$			M1 for correct matrix sum or $20 + 4n = 8$ or $-5 + 7n = -26$ or $n\mathbf{B} = \begin{pmatrix} -12 & 12 \\ -9 & -21 \end{pmatrix}$
			$n = -3, \quad x = -4, \quad y = 1.5$	3	A2 for all 3 values correct, A1 for 2 correct values
					Total 3 marks
15	(a)	$(x \times 10^5 + y \times 10^3) \div 10^5 \text{ oe}$			M1 for dividing a correct expression in x and y by 10^5
			$x + y \times 10^{-2}$	2	A1 oe (e.g. $x + \frac{y}{10^2}$ or $x + \frac{y}{100}$ or $\frac{100x+y}{100}$ or $\frac{10^2x+y}{10^2}$ or $x + 0.01y$) but A0 for $(x + y) \times 10^{-2}$ - mark answer line or final answer if no expression given on answer line
15	(b)				M1 for 34×10^{132}
			3.4×10^{133}	2	A1
					Total 4 marks

16		7.15, 7.25, 9.25, 9.35, 15.35, 15.45			B1 for one of the given values, accept 7.149 etc.
		$\left(\frac{9.35}{7.15}\right)^2 \times 15.45$			M1 or for $\left(\frac{UB(AE)}{LB(BD)}\right)^2 \times (UB(BCD))$ where $9.3 < UB(AE) \leq 9.35$ $7.15 \leq LB(BD) < 7.2$ $15.4 < UB(BCD) \leq 15.45$
			26.4	3	A1 (26.4204142...) – must come from correct bounds (9.35, 7.15 and 15.45 only) – allow 26.5
					Total 3 marks
17		67 + (180 – 123) or 180 – [180 – (67 + (180 – 123))] or 360 – 123 – (180 – 67) or for one of EBC = 67° BEC = 57° BCE = 56°			M1 – angle(s) could be shown on the diagram
			124		A1
				4	B2 (dependent on M1 A1) for all correct reasons for method used. e.g. angles on a <u>straight line</u> , <u>opposite angles</u> & <u>exterior angle</u> (is equal to the sum of the 2 opposite interior angles) (so the word 'angle' only needs to be used once) allow symbols for 'angles' and 'triangle' and allow abbreviations for the underlined words (e.g. VOA or OA for opposite angles) or angles on a <u>straight line</u> (mentioned twice), <u>opposite angles</u> , <u>angles</u> in a <u>triangle</u> oe (condone 'adjacent angles on a straight line' but not 'alternate angles on a straight line') - reasons must be given alongside the corresponding working (and not just appearing somewhere in their response) B1 (dependent on M1) for 2 correct reasons for method used
					Total 4 marks
18	(a)	$x + 3x + 3x - 7$ oe			M1

		$x + 3x + 3x - 7 = 56$ oe		2	A1 – award if seen in working – condone incorrect simplification
18	(b)	$7x = 56 + 7$ or $x = 9$			M1ft dep on 3 people's crayons (so must have three terms in x) – must isolate x terms correctly - award this mark if $x = 9$ stated in part (a)
			20	2	A1
					Total 4 marks
19		e.g. $\frac{1}{2}(\frac{2700}{90} + 4)$ or $180(n - 2) = 2700$ or $(2700 + 360) \div 180$ oe			M1 a fully correct method to find the number of sides
		17 (sides)			A1
			158.8 or better	3	A1 – note that 159 without 158.8 (or better) seen is A0
					Total 3 marks
20		$\frac{3(5\sqrt{2} - 2)}{2 - \sqrt{2}}$ oe			M1
		$\frac{3(5\sqrt{2} - 2)}{2 - \sqrt{2}} \times \frac{2 + \sqrt{2}}{2 + \sqrt{2}}$ oe			M1
			$9 + 12\sqrt{2}$	3	A1 dep on M2
					Total 3 marks
20	ALT	$(a + b\sqrt{2})(2 - \sqrt{2}) = 3(5\sqrt{2} - 2)$ $(2a - 2b) + (-a + 2b)\sqrt{2} = -6 + 15\sqrt{2}$ $2a - 2b = -6, -a + 2b = 15$ $a - b = -3, -a + 2b = 15$			M1 (allow one slip) – getting to simultaneous equations in a and b
					M1dep solve simultaneous equations (see Qu21 first M mark for applying the method mark for solving simultaneous equations)
			$a = 9, b = 12$		A1
21		$6x + 8y = 9$ OR $9x + 12y = 13.5$ oe			M1 for coefficient of x or y the same in both equations and correct operation to

		$\frac{6x - 9y = 34.5}{17y = -25.5} \quad \frac{8x - 12y = 46}{17x = 59.5}$ OR $x = \frac{11.5 + 3y}{2} \text{ and } 2\left(\frac{4.5 - 4y}{3}\right) - 3y = 11$ OR $y = \frac{4.5 - 3x}{4} \text{ and } 2x - 3\left(\frac{4.5 - 3x}{4}\right) = 11.5$			eliminate selected variable (condone one arithmetic error) or for correct rearrangement of one equation followed by correct substitution into the other
		e.g. $3 \times 3.5 + 4y = 4.5$ or $3x + 4 \times -1.5 = 4.5$			M1dep for substituting their found value correctly into one of the correct equations or correct use of elimination or substitution again.
			$x = 3.5,$ $y = -1.5$	4	First A1 dep on first M1 (so for their first value) Second A1 dep on second M1 (so for their second value)
					Total 4 marks
21	ALT	$\begin{pmatrix} 3 & 4 \\ 2 & -3 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 4.5 \\ 11.5 \end{pmatrix}$ $\begin{pmatrix} 3 & 4 \\ 2 & -3 \end{pmatrix}^{-1} = \frac{1}{3(-3) - 2(4)} \begin{pmatrix} -3 & -4 \\ -2 & 3 \end{pmatrix}$			M1 for inverse of correct 2 by 2 matrix (allow one slip only)
		$-\frac{1}{17} \begin{pmatrix} -3 & -4 \\ -2 & 3 \end{pmatrix} \begin{pmatrix} 4.5 \\ 11.5 \end{pmatrix} = -\frac{1}{17} \begin{pmatrix} -59.5 \\ 25.5 \end{pmatrix}$			M1dep multiplying their inverse by $\begin{pmatrix} 4.5 \\ 11.5 \end{pmatrix}$ to obtain at least a 2 by 1 matrix
			$x = 3.5,$ $y = -1.5$		A2 (A1 for one correct value)
22					M1 for $EC = DB$ together with D and E are <u>midpoints</u> (oe e.g. middle but must be in words and not just shown algebraically) and one of: <u>equal sides</u> or

					mention <u>isosceles</u> or $AB = AC$
					M1 for BC <u>common</u> side or $BC = BC$ or $BC = CB$
					M1 for $\angle DBC = \angle ECB$ (oe e.g. $\angle ABC = \angle ACB$) together with base angles in an <u>isosceles</u> triangle are equal All M marks are independent of each other but reasons must be given alongside the corresponding result (and not just appearing somewhere in their response)
			SAS	4	A1 for all correct reasons stated and SAS (dependent on all previous M marks)
Total 4 marks					
23		$\frac{(3^3)^{3x}}{(3^2)^y} = 3^{2x+x+1}$			M1 for one of $(3^3)^{3x}$ or $(3^2)^y$ or 3^{2x+x+1} (or better)
		$\frac{(3^3)^{3x}}{(3^2)^y} = 3^{2x+x+1}$			M1 for all of three of $(3^3)^{3x}$, $(3^2)^y$ and 3^{2x+x+1} (or better)
		$9x - 2y = 3x + 1$			M1 for $\alpha x - \beta y = 2x + x + 1$ (oe) with either $\alpha = 9$ or $\beta = 2$ (so dependent on first M mark only)
			$y = \frac{6x-1}{2}$	4	A1 oe
Total 4 marks					
24	(a)		Correct bisector	2	B2 correct perpendicular bisector of AB , with 2 sets of arcs and bisector drawn B1 for 2 sets of arcs with no line drawn or for a bisector drawn without arcs
					M1 for a correct bearing of 250° drawn
24	(b)		C correctly	2	A1 for C in the correct position

			positioned		
24	(c)		210 – 230 Inclusive	1	B1
					Total 5 marks
25		$y = -\frac{5}{4}x(+4)$ or gradient = $-\frac{5}{4}$			M1 or for $5 \times 8 + 4 \times 15 (=100)$
		$y = -\frac{5}{4}x + 25$ or $5x + 4y = 100$ or $y - 15 = -\frac{5}{4}(x - 8)$ (oe)			A1 for any correct form of the equation of the parallel line – or for one correct value (20 or 25) seen (if working shown then it must be correct)
		A(20, 0) and B(0, 25)			A1 for 20 and 25 (condone if 25 stated as A and 20 as B) – A and B do not need to be given as coordinates
		$\sqrt{20^2 + 25^2}$			M1 – correct method for finding their AB from their found values – using their equation but not from $5x + 4y = 16$ (independent of previous M mark)
			32	5	A1 (awrt)
					Total 5 marks
26	(a)	Use of 0.5 in the factor theorem			M1
		$6 \times 0.5^3 + 23 \times 0.5^2 - 5 \times 0.5 - 4 = 0$	Shown	2	A1 – must have = 0
26	(b)	$(3x^2 \dots)$			M1
		$(3x^2 + 13x + 4)$			A1 for correct 3TQ (correct 3TQ implies previous M mark)
		$(3x + 1)(x + 4)(=0)$			M1 a correct method to solve their 3TQ (that is must have $(3x + a)(x + b)$)

					with $ab = 4$ or $3b + a = 13$ for their 4 and 13) – or correct substitution into correct formula - dependent on first M mark
			$-\frac{1}{3}, -4$ only	4	A1dep both solutions and no extra solutions (e.g. A0 if $\frac{1}{2}$ included) Condone for the first three marks if a factor of $(2x - 1)$ is retained throughout
					Total 6 marks
27		$(AC^2 = 7.2^2 + 9^2 - 2 \times 7.2 \times 9 \times \cos(128^\circ))$			M1 (correct application of cosine rule – condone if stated as AC)
		$\sqrt{212.6...} (=14.58...)$			A1
		$\frac{\sin CAD}{6.4} = \frac{\sin 85^\circ}{14.58...}$ ($CAD = 25.9...$)			M1 or for $AC^2 = AD^2 + 6.4^2 - 2(AD)(6.4) \cos 85^\circ$ (oe complete method with AD in cosine rule) with their AC substituted $(AD^2 - 1.1155903507AD - 171.6697272 = 0$ $AD = (13.6(719426)))$ dependent on previous M mark
		$0.5 \times 7.2 \times 9 \times \sin(128^\circ) (=25.5...)$			M1 (correct application of $0.5ab \sin C$ for triangle ABC) – independent of previous M marks
		$0.5 \times 7.2 \times 9 \times \sin(128^\circ) +$ $0.5 \times 6.4 \times "14.58..." \times \sin(180 - 85 - "25.9...")$ $25.5... + 43.5.....$			M1 for adding the areas of their two triangles together – dependent on all previous M marks (oe for triangle ACD e.g. $0.5 \times 6.4 \times "13.6" \times \sin 85^\circ$)
			69.1	6	A1 accept answers in range 69.0 – 69.1
					Total 6 marks
28	(a)	$2.5 \times 8 + 12.5 \times 10 + 25 \times 15 + 35 \times 17 + 50 \times 25$ $20 + 125 + 375 + 595 + 1250 (= 2365)$ $"2365" \div 75$			M2 (Condone any one error only) if not M2 then award M1 for multiplication of all midpoints by frequencies (without addition) or multiplication of frequencies by one end of interval consistently and addition M1ft dep on M1
			31.5	4	A1 (31.5 or better)
28	(b)				M1 for $30 \times 1\text{cm}^2 = 75$ cars or correct frequency densities (2.4, 2.7 and 0.9) either stated by table or labelled on histogram or one correct value in table

			Correct values in table		A1 correct values of 30 and 45 in table
		2.4, 2.7, 0.9			M1 two correct bars in histogram – must be using 2 cm = a frequency density of 1 if vertical axis not labelled (however, if vertical axis is labelled it must be correct)
			Correctly completed histogram	4	A1 including frequency density axis labelled correctly (2 cm is equivalent to 1 unit) – note that a single value on the vertical axis is sufficient - heights of the bars are 2.4 (between 0 and 5), 2.7 (between 30 and 40) and 0.9 (between 40 and 60)
					Total 8 marks
29	(a)	$P = \frac{k}{w^2}$			M1
		$16 = \frac{k}{5^2}$ or $k = 400$			A1
			2.56	3	A1
29	(b)	$P = \frac{k}{y}, 75 = \frac{k}{4} \rightarrow k = 300$			M1 – obtaining correct value of k
		$P = \frac{'400'}{4} = 100$ and $'300' \div '100'$ or $\frac{w^2}{'400'} = \frac{y}{'300'}$ and $'300' \div '100'$ or			M1 dependent on both previous M marks (in (a) and (b))
			3	3	A1
					Total 6 marks

Pearson Education Limited. Registered company number 872828
with its registered office at 80 Strand, London, WC2R 0RL, United Kingdom