

Write your name here

Surname

Other names

**Pearson Edexcel Certificate**

Centre Number

Candidate Number

**Pearson Edexcel  
International GCSE**

--	--	--	--	--

--	--	--	--

# Chemistry

**Unit: KCH0/4CH0**

**Paper: 2C**

Wednesday 14 June 2017 – Morning

**Time: 1 hour**

Paper Reference

**KCH0/2C  
4CH0/2C**

**You must have:**

Calculator

Total Marks

## Instructions

- Use **black** ink or ball-point pen.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions.
- Answer the questions in the spaces provided  
– *there may be more space than you need.*
- Show all the steps in any calculations and state the units.
- Some questions must be answered with a cross in a box ☒. If you change your mind about an answer, put a line through the box ☒ and then mark your new answer with a cross ☒.

## Information

- The total mark for this paper is 60.
- The marks for **each** question are shown in brackets  
– *use this as a guide as to how much time to spend on each question.*

## Advice

- Read each question carefully before you start to answer it.
- Write your answers neatly and in good English.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ►

P48083A

©2017 Pearson Education Ltd.

1/1/1/1/



  
Pearson

# THE PERIODIC TABLE

Period 1 2 3 4 5 6 7 0

Group

4	He	Helium	2
---	----	--------	---

1	H	Hydrogen	1
---	---	----------	---

7	Li	Lithium	3	9	Be	Beryllium	4	20	Ne	Neon	10
23	Na	Sodium	11	24	Mg	Magnesium	12	31	P	Phosphorus	15
39	K	Potassium	19	40	Ca	Calcium	20	73	Ge	Germanium	32
86	Rb	Rubidium	37	88	Sr	Strontium	38	115	In	Indium	49
133	Cs	Caesium	55	137	Ba	Barium	56	204	Tl	Thallium	81
223	Fr	Francium	87	226	Ra	Radium	88	207	Pb	Lead	82
227	Ac	Actinium	89	227	Fr	Francium	87	207	Pb	Lead	82
45	Sc	Scandium	21	46	Ti	Titanium	22	59	Ni	Nickel	28
89	Y	Yttrium	39	90	Zr	Zirconium	40	106	Pd	Palladium	46
139	La	Lanthanum	57	140	Hf	Hafnium	72	195	Pt	Platinum	78
227	Ac	Actinium	89	227	Fr	Francium	87	207	Pb	Lead	82
51	V	Vanadium	23	52	Cr	Chromium	24	63.5	Cu	Copper	29
93	Nb	Niobium	41	94	Mo	Molybdenum	42	108	Ag	Silver	47
181	Ta	Tantalum	73	182	W	Tungsten	74	197	Au	Gold	79
48	Ti	Titanium	22	49	V	Vanadium	23	59	Ni	Nickel	28
91	Zr	Zirconium	40	92	Nb	Niobium	41	106	Pd	Palladium	46
179	Hf	Hafnium	72	180	Ta	Tantalum	73	197	Au	Gold	79
56	Fe	Iron	26	57	Mn	Manganese	25	59	Ni	Nickel	28
101	Ru	Ruthenium	44	102	Rh	Rhodium	45	106	Pd	Palladium	46
190	Os	Osmium	76	191	Ir	Iridium	77	197	Au	Gold	79
112	Cd	Cadmium	48	113	In	Indium	49	115	Tl	Thallium	81
201	Hg	Mercury	80	202	Tl	Thallium	81	207	Pb	Lead	82
127	I	Iodine	53	128	Xe	Xenon	54	129	Ba	Barium	56
210	At	Astatine	85	211	Rn	Radon	86	210	Po	Polonium	84

Key

Relative atomic mass
Symbol
Name
Atomic number

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



**Answer ALL questions.**

1 Sodium sulfate is a compound with many uses.

(a) The formula of the main compound used as the source of sodium sulfate is  $\text{Na}_2\text{SO}_4 \cdot 10\text{H}_2\text{O}$

How many different elements are shown in this formula?

(1)

A 2

B 3


C 4

D 5

(b) Sodium sulfate can be made from sodium hydroxide and sulfuric acid.

Balance the equation for the reaction between sodium hydroxide and sulfuric acid.

(1)



(c) Sodium hydroxide is manufactured by the electrolysis of brine in the diaphragm cell.

Sulfuric acid is manufactured using the contact process.

The table contains some statements about these two processes.

Place ticks (✓) in the boxes to show the two correct statements.

(2)

brine is a solution of sodium chloride in water	
the temperature used in the contact process is greater than 1000 °C	
an equation for the contact process is $\text{SO}_2 + \text{H}_2\text{O} \rightarrow \text{H}_2\text{SO}_4$	
the reactions in the diaphragm cell are displacement reactions	
the catalyst used in the contact process is vanadium(V) oxide	

**(Total for Question 1 = 4 marks)**



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

**BLANK PAGE**



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

2 The diagram shows the positions of some elements in part of the Periodic Table.

							He
Li		B		N		F	
	Mg		Si		S		Ar

(a) How many periods and groups are shown in this diagram?

(1)

	Periods	Groups
<input type="checkbox"/> A	2	4
<input type="checkbox"/> B	3	4
<input type="checkbox"/> C	2	8
<input type="checkbox"/> D	3	8

(b) How many elements shown in the diagram are noble gases?

(1)

- A 1
- B 2
- C 3
- D 4

(c) What is the formula of the compound formed between magnesium and fluorine?

(1)

- A MgF
- B Mg<sub>2</sub>F
- C MgF<sub>2</sub>
- D Mg<sub>2</sub>F<sub>2</sub>



(d) The table shows the percentage composition by mass of a sample of silicon.

Isotope	$^{28}\text{Si}$	$^{29}\text{Si}$	$^{30}\text{Si}$
Percentage (%)	92.2	4.70	3.10

Calculate the relative atomic mass of this sample of silicon.

Give your answer to one decimal place.


(2)

relative atomic mass = .....

(e) A molecule of silicon tetrafluoride ( $\text{SiF}_4$ ) contains covalent bonds.

Draw a dot and cross diagram to show the outer electrons in this molecule.

(2)



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(f) The table shows the boiling points of some compounds containing silicon. All of these compounds contain covalent bonds.

Compound	Boiling point in °C
SiF <sub>4</sub>	-86
SiCl <sub>4</sub>	58
SiO <sub>2</sub>	2950

SiF<sub>4</sub> and SiCl<sub>4</sub> have simple molecular structures.

SiO<sub>2</sub> has a giant covalent structure.

(i) Explain why the boiling point of SiCl<sub>4</sub> is greater than the boiling point of SiF<sub>4</sub> (2)

.....

.....

.....

.....

.....

.....

(ii) Explain why the boiling point of SiO<sub>2</sub> is very much greater than the boiling point of SiCl<sub>4</sub> (2)

.....

.....

.....

.....


.....

.....

**(Total for Question 2 = 11 marks)**



3 This apparatus can be used to investigate electrolysis.



(a) Name the particles that move through the connecting wires to form an electric current. (1)

(b) The electrodes are made of platinum, which is an inert metal.

State what is meant by the term **inert**. (1)

(c) Explain why the electrolytic cell containing  $\text{PbBr}_2$  needs to be heated before electrolysis can occur. (2)



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(d) When NaCl(aq) is electrolysed, two gases form at the positive electrode and one gas forms at the negative electrode.

The formulae of the species in NaCl(aq) are Na<sup>+</sup>, Cl<sup>-</sup>, H<sup>+</sup>, OH<sup>-</sup> and H<sub>2</sub>O.

(i) Name the gases formed at each electrode. (2)

positive electrode ..... and .....

negative electrode.....

(ii) Give ionic half-equations to show the formation of each gas. (3)

.....  
.....  
.....  
.....  
.....  
.....  
.....

(e) The ionic half-equation for one of the reactions in the cell containing copper(II) sulfate solution is



During the electrolysis, a charge of 0.040 faradays passes through this cell.

Calculate the mass of copper metal formed. (2)

mass of copper = ..... g

**(Total for Question 3 = 11 marks)**



- 4 Compounds containing C=C double bonds are used to manufacture alcohols such as ethanol and addition polymers such as PVC.

The table shows the formulae of some compounds containing C=C bonds.

<b>A</b>  $\text{CH}_2=\text{CH}_2$	<b>B</b>  $\begin{array}{c} \text{H} \quad \text{H} \quad \text{H} \\   \quad   \quad   \\ \text{H}-\text{C}=\text{C}-\text{C}-\text{H} \\   \\ \text{H} \end{array}$
<b>C</b>  $\text{C}_4\text{H}_8$	<b>D</b>  $\text{C}_2\text{H}_3\text{Cl}$

- (a) (i) Explain which three of these compounds are hydrocarbons.

(2)

.....

.....

.....

.....

- (ii) Explain which compound is shown as a displayed formula.

(2)

.....

.....

.....

.....



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

- (b) Ethanol can be manufactured from compound A using reaction 1.  
Ethanol can also be manufactured from glucose using reaction 2.

The equations for these reactions are



Give two advantages of using each reaction to manufacture ethanol.

(4)

reaction 1 .....

.....

.....

.....

.....

reaction 2 .....


.....

.....

.....

.....

- (c) Compound C reacts with bromine to form a product with this formula.



- (i) Use this formula to determine the name of compound C.

(2)

.....

- (ii) State the colour of the product formed when compound C reacts with bromine.

(1)

.....




(d) Compound X is an isomer of compound C.

(i) Explain what is meant by the term **isomers**.

(2)

(ii) The equation for the conversion of compound X to an alcohol is



Place ticks (✓) in the boxes to show the two correct descriptions of this reaction.

(2)

addition	
dehydration	
hydration	
oxidation	
reduction	

(e) Compound D, chloroethene, can be used to manufacture the polymer PVC.

(i) State the full name of PVC.

(1)

(ii) Write an equation, containing displayed formulae, to represent the reaction that occurs in the manufacture of PVC.

(3)

(Total for Question 4 = 19 marks)

DO NOT WRITE IN THIS AREA


DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

**BLANK PAGE**



- 5 A student uses this apparatus to find the increase in temperature of water when methanol,  $\text{CH}_3\text{OH}$ , is burned.



(a) There are several reasons why the increase in temperature is less than expected.

- (i) One reason is the incomplete combustion of methanol to form only carbon monoxide and water.

Write the chemical equation for this incomplete combustion.

(2)

- (ii) State another reason why the increase in temperature is less than expected.

(1)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(b) The student records these results.

mass of burner and methanol before combustion	84.7 g
mass of burner and methanol after combustion	83.2 g
mass of water	125 g
temperature of water at start	22 °C
temperature of water at end	58 °C

(i) Calculate the heat energy change ( $Q$ ), in joules, in this experiment using the expression

$$Q = m \times 4.2 \times \Delta T$$

where  $m$  is the mass of water in grams and  $\Delta T$  represents the increase in temperature.

(2)

$Q = \dots\dots\dots$  J

(ii) The relative molecular mass of methanol is 32

Use this information and your value for  $Q$  to calculate the molar enthalpy change,  $\Delta H$ , for the combustion of methanol.


Give your answer in kJ/mol.

(4)

$\Delta H = \dots\dots\dots$  kJ/mol



(iii) The student draws an energy level diagram for the complete combustion of methanol.



Identify the two mistakes in his diagram.

(2)

1 .....


2 .....

DO NOT WRITE IN THIS AREA

(c) The student is given this table of average (mean) bond energies.

Bond	C—H	C—O	O—H	O=O	C=O
Average bond energy in kJ/mol	412	360	463	496	743

The equation for the complete combustion of methanol is



Use this equation and the information in the table to calculate another value for the molar enthalpy change,  $\Delta H$ , for the combustion of methanol.

(4)

$$\Delta H = \dots\dots\dots \text{kJ/mol}$$

**(Total for Question 5 = 15 marks)**

**TOTAL FOR PAPER = 60 MARKS**



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

**BLANK PAGE**

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

**BLANK PAGE**



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

**BLANK PAGE**

Every effort has been made to contact copyright holders to obtain their permission for the use of copyright material. Pearson Education Ltd. will, if notified, be happy to rectify any errors or omissions and include any such rectifications in future editions.

