



# Mark Scheme (Results)

January 2020

Pearson Edexcel International GCSE in  
Chemistry (4CH1)  
Paper 2CR

## **Edexcel and BTEC Qualifications**

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at [www.edexcel.com](http://www.edexcel.com) or [www.btec.co.uk](http://www.btec.co.uk). Alternatively, you can get in touch with us using the details on our contact us page at [www.edexcel.com/contactus](http://www.edexcel.com/contactus).

## **Pearson: helping people progress, everywhere**

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: [www.pearson.com/uk](http://www.pearson.com/uk)

January 2020

Publications Code 4CH1\_2CR\_msc\_20200305

All the material in this publication is copyright

© Pearson Education Ltd 2020

## General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

Question number	Answer	Notes	Marks
1 (a) (i)	<p>B box 2 The only correct answer is B because box 2 contains two different particles in the same space that are not chemically joined</p> <p>A is not correct because box A shows an element C is not correct because box C shows an element D is not correct because box 4 shows a compound</p>		1
(ii)	<p>C Boxes 1 and 3 The only correct answer is C because boxes 1 and 2 contain one type of atom only. A is not correct because box 2 shows a mixture B is not correct because box 2 shows a mixture D is not correct because box 4 shows a compound</p>		1
(iii)	<p>M1 (box 5 shows) two (different) elements</p> <p>M2 (chemically) bonded (together)</p>	<p><b>ALLOW</b> two (different) types of atoms</p> <p><b>REJECT</b> mixture for M1</p> <p><b>ALLOW</b> (chemically) combined/joined (together)</p> <p><b>M2 DEP</b> on mention of elements/atoms in M1</p>	2
(b) (i)	<p>C number of protons</p> <p>The only correct answer is B because the elements in the periodic table are arranged in order of proton number A is not correct because elements are not arranged in order of mass number B is not correct because elements are not arranged in order of the number of neutrons D is not correct because elements are not arranged in order of reactivity</p>		1
(ii)	<p>A electrons in the outer shell</p> <p>The only correct answer is A because elements in the same group of the periodic table have the same number of electrons in the outer shell. B is not the correct answer because elements in the same period have the same number of shells C elements in the same group do not have the same number of neutrons D is not the correct answer because elements in the same group do not have the same number of protons</p>		1

Question number	Answer	Notes	Marks
2 (a)	<p>M1 level of the water must be below the dyes/start line</p> <p>M2 the start line must be drawn in pencil</p>	ACCEPT dyes/start line should be above water level	2
2 (b) (i)	<p>2 marks for any two conclusions from (the green food colouring)</p> <p>M1 contains (dye) B and (dye) D</p> <p>M2 contains an unknown dye</p> <p>M3 does not contain A or C</p> <p>M4 contains three dyes</p> <p>and 1 mark for a correct explanation of given conclusion for the green food colouring</p> <p>eg (explanation for M1) because has spots at same level (as B and D)</p> <p>eg (explanation for M2) because has a spot at different level (from A B C D)</p> <p>eg (explanation for M3) because has no spots at same level (as A and C)</p> <p>eg (explanation for M4) because has three spots</p>	ALLOW is not pure	3
(ii)	<p>M1 (distance moved by solvent correctly measured) = 9.5 (cm)</p> <p>M2 use of</p> $R_f = \frac{\text{distance moved by the dye C}}{\text{distance moved by the solvent}}$ <p>M3 evaluation of <math>R_f</math></p>	<p>ALLOW a tolerance of <math>\pm 2\text{mm}</math></p> <p>eg <math>\frac{6.2}{9.5}</math></p> <p>ALLOW ECF from M1</p> <p>eg <math>\frac{(6.2)}{(9.5)} = 0.65(3)</math></p> <p>ALLOW 1-4 sig fig</p> <p>ALLOW ECF from M2</p>	3
(iii)	(dye A) is not soluble in water	ALLOW solvent for water	1

Question number	Answer	Notes	Marks
3 (a)	<p>B Precipitation</p> <p>The only correct answer is B the reaction of two solutions to produce an insoluble solid is precipitation.</p> <p>A is not correct because this reaction is not neutralisation</p> <p>C is not correct because this reaction is not a redox reaction</p> <p>D is not correct because this reaction is not thermal decomposition</p>		1
3 (b)	<p>M1 wash the solid with (deionised) water</p> <p>M2 suitable method of drying solid eg dry between filter papers/on paper towel/in (warm) oven/in a desiccator</p>	<p><b>ALLOW</b> leave to dry</p> <p><b>ALLOW</b> leave in a warm place</p> <p><b>ALLOW</b> leave for the water to evaporate</p> <p><b>IGNORE</b> dry it alone</p> <p><b>REJECT</b> hot oven or any method of direct heating eg Bunsen burner</p> <p><b>REJECT</b> references (direct or inferred) to silver chloride solution or crystallisation for M1 and M2</p> <p>No M2 if solid washed after drying</p>	2
3 (c)	<p>Any one from:</p> <p>M1 (hydrochloric acid/it) contains chloride ions</p> <p>M2 (hydrochloric acid/it) produces a (white) precipitate with silver nitrate</p> <p>M3 (hydrochloric acid/it) reacts with silver nitrate</p>	<b>ALLOW</b> contains Cl <sup>-</sup>	1
3 (d)	<p>M1 <math>n(\text{AgNO}_3 \text{ or } \text{AgCl}) = 0.0025</math></p> <p>M2 <math>(\text{mass AgCl}) = 0.0025 \times 143.5 = 0.35(9)\text{g}</math></p>	<p><b>ALLOW</b> ECF from M1</p> <p><b>ALLOW</b> one or more sig fig</p> <p>Correct answer without working scores 2 marks.</p>	2


Question number	Answer	Notes	Marks
4 (a)	M1 layers of atoms/positive ions  M2 can slide over one another	IGNORE layers unqualified REJECT layers of molecules  M2 DEP on mention of layers/atoms/ions in M1	2
4 (b) (i)	ions cannot move	ALLOW ions are in fixed positions/in a lattice  IGNORE no free ions REJECT any reference to electrons	1
(ii)	$2\text{Br}^- \rightarrow \text{Br}_2 + 2\text{e}^-$		1
(iii)	M1 lead ions (are positive and) are attracted to the negative electrode / $\text{Pb}^{2+}$ (ions) are attracted to the negative electrode  M2 lead ions gain electrons / $\text{Pb}^{2+}$ (ions) gain electrons (to form lead)	ALLOW cathode for negative electrode  ALLOW a correct half equation for M2 IGNORE references to redox ALLOW lead ions get discharged (to form lead)	2
(iv)	metal or lead connects the electrodes or completes the circuit OWTTE	ALLOW metal or lead conducts electricity ALLOW metal or lead allows electrons to flow	1

Question number	Answer	Notes	Marks
5 (a) (i)	Any two from: M1 (lithium) moves (on the surface) M2 (lithium) gets smaller/disappears M3 colourless solution forms	ALLOW floats ALLOW dissolves IGNORE white trail forms REJECT melts / turns into a ball ALLOW temperature increases/heat given off	2
(ii)	(when mixed with air) lit spill/splint or flame gives (squeaky) pop	must refer to test and result IGNORE squeaky pop test alone ALLOW burns with (squeaky) pop REJECT glowing spill/splint and pop	1
5 (b) (i)	Any one from: M1 more rapid bubbles/fizzing/effervescence M2 turns into a ball M3 moves more quickly M4 catches alight / burns / produces a flame	ALLOW potassium melts ALLOW gets smaller/disappears more quickly IGNORE flame colour	1
(ii)	M1 potassium has more shells than lithium  M2 (therefore) there is less attraction between the outer shell/electron and the nucleus  M3 so the electron in the outer shell is more easily lost	ALLOW potassium atom is bigger than lithium ALLOW outer shell/electron is further from nucleus ALLOW more repulsion (from inner shells) or more shielding (from the nuclear attraction) ALLOW nuclear pull for the outer shell/electron is weaker ACCEPT answers in terms of lithium for M1, M2 and M3	3

Question number	Answer	Notes	Marks
5 (c) (i)	<p>M1 <math>n(\text{Li}) = \frac{0.500}{7}</math> OR 0.0714</p> <p>M2 <math>n(\text{H}_2) = \frac{0.0714}{2}</math> OR 0.0357</p> <p>M3 volume of <math>\text{H}_2</math> (<math>= 0.0357 \times 24000</math>) = 857 (<math>\text{cm}^3</math>)</p>	<p>ALLOW ECF from M1</p> <p>ALLOW ECF M2 x 24 000</p> <p>M3 must be to 3 sig fig</p> <p>Correct answer to 3 sig fig without working scores 3 marks.</p>	3
(ii)	<p>M1 <math>n(\text{H}_2\text{SO}_4) = 0.02485 \times 0.1</math> OR 0.002485</p> <p>M2 <math>n(\text{LiOH}) = 2 \times 0.002485</math> OR 0.00497</p> <p>M3 concentration of LiOH = 0.0331 (<math>\text{mol}/\text{dm}^3</math>)</p>	<p>ALLOW ECF from M1</p> <p>ALLOW ECF from M2 (M2 <math>\div</math> 0.150)</p> <p>ALLOW any number of sig fig except one for M1 M2 and M3</p> <p>Correct answer without working scores 3 marks.</p>	3

Question number	Answer	Notes	Marks
6 (a)	<p>A description including the following points:</p> <p>M1 (use) fractional distillation / fractionating column / fractionating tower</p> <p>M2 (crude oil) heated / vaporised</p> <p>M3 column is cooler at top / hotter at the bottom / idea of temperature gradient</p> <p>M4 fractions condense/collected at different heights OWTTE</p> <p>OR fractions have different boiling point (ranges)</p>	<p>ALLOW boiled</p> <p>ALLOW fractions with high(er) boiling points/large(r) molecules condense/collected near bottom ORA</p>	4
6 (b) (i)	<p>M1 particles/molecules are closer together</p> <p>M2 therefore more (successful) collisions per unit time</p>	<p>ALLOW more particles/molecules per unit volume</p> <p>ALLOW more frequent (successful) collisions If reference to particles have more energy/move faster, then 0/2</p>	2
(ii)	<p>M1 (a catalyst provides) an alternative pathway</p> <p>M2 of lower activation energy</p>	<p>ALLOW a catalyst provides a surface for the reaction to take place on</p> <p>ALLOW so more collisions/particles have energy greater than the activation energy.</p>	2
(iii)	increase the temperature	<p>ALLOW heat it up IGNORE references to concentration REJECT increase the surface area</p>	1

6 (b) (iv)



M1 C<sub>2</sub>H<sub>6</sub> on horizontal line below the reactants

M2 activation energy correctly shown and labelled

M3  $\Delta H$  correctly shown and labelled

ACCEPT arrow pointing upwards or double headed arrow

ACCEPT arrow pointing from reactants level to product level or double headed arrow

ALLOW max 2 for a correctly labelled endothermic reaction

(c)

- find energy needed to break bonds
- find energy released when bonds form
- correct evaluation of  $\Delta H$

Example calculation

M1  $(4 \times 412) + 612 + 436$  OR 2696

M2  $(6 \times 412) + 348$  OR 2820

M3 -124 if M1 M2 correct

IGNORE sign  
ALLOW 1048

IGNORE sign  
ALLOW 1172

-124 without working scores 3  
(+)-124 scores 2

ALLOW M3 ECF M1 M2

3

3

Question number	Answer	Notes	Marks
7 (a) (i)	CO <sub>2</sub>		1
(ii)	(otherwise) ethanoic acid will form	<b>ALLOW</b> (otherwise) ethanol will be oxidised or react with oxygen <b>ALLOW</b> fermentation/reaction/respiration needs to be anaerobic <b>ALLOW</b> (otherwise) ethanol would not be formed /CO <sub>2</sub> and H <sub>2</sub> O would be formed	1
(iii)	M1 (reaction is catalysed by) enzymes (in yeast) M2 which will denature (above 40°C)	<b>IGNORE</b> yeast unqualified <b>ALLOW</b> enzymes do not work above 40°C	2
(iv)	M1 maximum mass of ethanol = 8 x 46 = 368 (g) M2 $\frac{55.2}{368} \times 100$ (=15%)  Alternative method: M1 actual yield of ethanol in mol = $\frac{55.2}{46} = 1.2$ M2 $\frac{1.2}{8} \times 100$ (=15%)		2
(b) (i)	M1 rate of the forwards reaction = the rate of the backwards reaction  M2 the concentrations of reactants and products remain constant	<b>IGNORE</b> it is reversible reaction  <b>REJECT</b> concentrations of reactants and products are equal or are the same	2
(b) (ii)	M1 an increase in temperature shifts the (position of) equilibrium in the endothermic direction (so backwards reaction is endothermic)  M2 so forward reaction is exothermic	<b>IGNORE</b> references to Le Chatelier's Principle <b>ALLOW</b> heating for increase in temperature  M2 DEP M1 or near miss	2

7 (c) (i)	<p>M1 (displayed formula of A - propanoic acid)</p> $  \begin{array}{c}  \text{H} \quad \text{H} \quad \quad \text{O} \\    \quad   \quad \quad // \\  \text{H}-\text{C}-\text{C}-\text{C} \\    \quad   \quad \quad \backslash \\  \text{H} \quad \text{H} \quad \quad \text{O}-\text{H}  \end{array}  $ <p>M2 (displayed formula of B - butan-1-ol)</p> $  \begin{array}{c}  \text{H} \quad \text{H} \quad \text{H} \quad \text{H} \\    \quad   \quad   \quad   \\  \text{H}-\text{C}-\text{C}-\text{C}-\text{C}-\text{O}-\text{H} \\    \quad   \quad   \quad   \\  \text{H} \quad \text{H} \quad \text{H} \quad \text{H}  \end{array}  $	<p><b>ALLOW</b> 1 mark if both OH but otherwise correct</p>	2
(c) (ii)	<p>M1 add a named carbonate or hydrogencarbonate</p> <p>M2 effervescence/bubbles/fizzing</p> <p>OR</p> <p>M1 add a suitable named metal e.g. magnesium, aluminium, zinc, iron</p> <p>M2 effervescence/bubbles/fizzing</p> <p>OR</p> <p>M1 add a named alcohol (and some concentrated sulfuric acid and warm)</p> <p>M2 sweet smell (of an ester)</p>	<p><b>ALLOW</b> correct formula</p> <p>M2 DEP M1 or near miss <b>ALLOW</b> carbon dioxide/CO<sub>2</sub> produced <b>REJECT</b> incorrect gas</p> <p><b>ALLOW</b> correct symbol <b>REJECT</b> a metal that is too reactive e.g. potassium or too unreactive e.g. copper</p> <p>M2 DEP M1 or near miss <b>ALLOW</b> hydrogen/H<sub>2</sub> produced</p> <p><b>REJECT</b> incorrect gas</p> <p><b>ALLOW</b> correct formula</p> <p>M2 DEP M1 or near miss</p>	2

Pearson Education Limited. Registered company number 872828  
with its registered office at 80 Strand, London, WC2R 0RL, United Kingdom