



Mark Scheme (Results)

November 2024

Pearson Edexcel International GCSE
In Chemistry (4CH1) Paper 1C

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

November 2024

Question Paper Log Number P75946A

Publications Code 4CH1_1C_2411_MS

All the material in this publication is copyright

© Pearson Education Ltd 2024

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

Question number	Answer	Notes	Marks
1 (a)	from liquid to solid freezing from gas to liquid condensing from solid to gas sublimation	ALLOW condensation ALLOW subliming	3
(b) (i)	6 circles similar size randomly arranged none touching	At least one circle in top/bottom of box	1
(ii)	A (the atoms move randomly in the gas state) B is not correct since atoms do not move randomly in the solid state C is not correct since atoms are not in a fixed position in the gas state D is not correct since atoms are not in a fixed position in the liquid state		1
(c)	$\text{H}_2\text{O}(\text{l}) \rightarrow \text{H}_2\text{O}(\text{s})$	ALLOW upper case/lower case	1
		Total = 6	

Question number	Answer	Notes	Marks
2 (a)	(i) oxygen	ALLOW O ₂	1
	(ii) nitrogen	ALLOW N ₂	1
	(iii) chlorine	ALLOW Cl ₂	1
(b)	M1 (hydrogen chloride) has (atoms of) two / different elements M2 (chemically) bonded/joined / (chemically) combined together	ALLOW two different atoms	2
(c)	D (chlorine has the strongest forces of attraction between its molecules) A is not the correct answer because covalent bonds are not broken when chlorine boils B is not the correct answer because covalent bonds do not occur between molecules C is not the correct answer because chlorine does not have ionic bonds		1
		Total = 6	

Question number	Answer	Notes	Marks
3 (a)	(i) M1 oxygen	ALLOW air ALLOW O ₂	2
	M2 water	ALLOW moisture ALLOW H ₂ O	
	(ii) (hydrated) iron (III) oxide	ALLOW ferric oxide ALLOW Fe ₂ O ₃	1
(b)	(i) M1 paint acts as a barrier / (protective) layer OWTTE	NOT galvanising NOT coating/covering	2
	M2 which prevents water/oxygen/air getting to the iron/reacting with iron		
	(ii) galvanising	ALLOW sacrificial protection	1
	(iii) M1 zinc is more reactive than iron OR zinc has a greater tendency to lose electrons	ALLOW zinc reacts instead of iron NOT zinc reacts more rapidly/faster	2
	M2 zinc oxidises / forms zinc oxide/reacts before iron	REJECT references to zinc rusting REJECT zinc reacts with iron	
		Total = 8	



Question number	Answer	Notes	Marks
4 (a)	<p>M1 draw a line in pencil (just above the bottom of the paper)</p> <p>M2 put a spot of each ink on the line (before contact with solvent)</p> <p>M3 pour some solvent in the beaker</p> <p>M4 place the paper in the beaker so the spots are above the solvent</p> <p>M5 leave until the solvent has risen up the paper (nearly to the top)</p>	<p>ALLOW water for solvent</p> <p>ALLOW water for solvent</p> <p>ALL marks can be scored/supported from a labelled diagram</p>	5
(b) (i)	<p>M1 E</p> <p>M2 because it stayed on the start line/did not travel up paper</p>	<p>ALLOW didn't move/ Rf value =0</p> <p>M2 dep on M1</p>	2
(ii)	<p>M1 A and C</p> <p>M2 because they both (have a spot) at the same height OWTTE</p>	<p>ALLOW travelled same distance/same Rf value</p> <p>M2 dep on M1</p>	2
(iii)	<p>M1 measure the distance from the start line to the spot and the distance from the start line to the solvent front</p> <p>M2 distance moved by the spot ÷ distance moved by the solvent</p>	<p>M2 subsumes M1</p> <p>Allow 2 marks for a correct calculation method</p>	2
		Total = 11	

Question number	Answer	Notes	Marks
6 (a) (i)	AlCl ₃ ZnSO ₄ (NH ₄) ₃ N	ALLOW formula in reverse NOT molecular formula Penalise symbol letters/size of subscripts once only	3
(ii)	aluminium sulfate	ALLOW aluminium sulphate	1
(b)	M1 magnesium loses electrons M2 chlorine gains electrons M3 magnesium loses two electrons and two chlorines each gain one electron	ALLOW magnesium gives/transfers electrons to chlorine for M1,M2 NOT chloride gains electrons M3 assumes M1,M2 ALLOW correct ionic equations	3
(c) (i)	M1 two electrons between each nitrogen and hydrogen atom M2 two non-bonding electrons	M2 dep on M1	2
(ii)	M1 (electrostatic) forces of attraction between shared pair(s) of electrons M2 and the nuclei	REJECT nucleus (must be plural) REJECT intermolecular forces for both marks	2
		Total = 11	

Question number	Answer	Notes	Marks
7 (a) (i)	any one from: M1 to condense the water vapour M2 to ensure all the water collects in the tube (as a liquid)	ALLOW condense steam/condense gas NOT cools water NOT stops water evaporating	1
(ii)	When the mass doesn't change / is constant/stops increasing	Accept: the last two results are the same Accept: balance reading stays the same	1
(b)	M1 add anhydrous/white copper(II) sulfate/sulphate M2 which turns (from white to) blue	ALLOW anhydrous/white copper sulfate/sulphate ALLOW add anhydrous/blue cobalt chloride ALLOW which turns (from blue to) pink M2 dep on M1	2
(c)	M1 (mass of water) = 6.3g M2 (moles of MgSO_4) = 0.05 M3 (moles of H_2O) 0.35 M4 $x=7$	Ecf for incorrect mass of water M1 can be awarded from moles of H_2O in M3 Answer of 7 on its own scores 4 marks	4
		Total = 8	

Question number	Answer	Notes	Marks
8 (a)	carbon dioxide/a gas escapes/is lost/released (through the cotton wool)	NOT carbon dioxide/gas is given off/produced NOT wrong named gas	1
(b)	M1 the concentration (of hydrochloric acid) is highest M2 so there are more collisions per unit time	ALLOW there is a greater surface area of marble chips ALLOW greater amount of hydrochloric acid/reactants ALLOW more particles ALLOW more frequent collisions REJECT references to greater (kinetic) energy for both marks	2
(c)	the hydrochloric acid has been used up OWTTE	NOT acid is saturated IGNORE acid is a limiting factor	1
(d) (i)	any two from: (same) mass of marble chips (same) surface area of marble chips (same) concentration of hydrochloric acid (same) volume of hydrochloric acid	ALLOW (same) amount of marble chips ALLOW (same) size marble chips NOT same amount of acid	2 3
(ii)	M1 rate of reaction increases M2 particles have more energy OR more particles have energy greater than (or equal to) the activation energy M3 so more successful collisions per unit time	ALLOW particles move faster ALLOW more frequent successful collisions	
		Total = 9	

Question number	Answer	Notes	Marks
9 (a)	aluminium is a better conductor (of heat) than glass (comparison needed)	REJECT insulation references	1
(b) (i)	carbon / soot/ C		1
(ii)	incomplete combustion occurs OR the supply of oxygen/air is limited		1
(c) (i)	M1 $1100 \times 4.2 \times 50$ M2 21000(J)	ALLOW ecf for M2 if answer close to 20000J	2
(ii)	M1 21 kJ M2 $1.84 \div 46$ OR 0.04 moles M3 $21 \div 0.04$ OR 525 (kJ/mol) M4 -525 (kJ/mol)	ALLOW 20kJ ALLOW $21 \div M2$ ALLOW 500 (kJ/mol) if 20kJ used M4 is for the - sign. ALLOW ecf from M3	4
(d) (i)	$5O_2$	ALLOW multiples if the rest of the balancing numbers have been multiplied	1
(ii)	M1 (M_r of butanol) 74 M2 (moles of butanol) $3.7 \div 74$ OR 0.05 M3 0.45 moles	ALLOW $3.7 \div M1$ if attempted M_r shown ALLOW $M2 \times 9$ Answer of 0.45 scores 3	3
		Total = 13	

Question number	Answer	Notes	Marks
10 (a) (i)	any one from: M1 ethane is saturated M2 ethane has no double bonds M3 ethane has single bonds only		1
(ii)	M1 products C_2H_5Br and HBr M2 condition ultra violet radiation / ultra violet light/UV	In either order ALLOW balanced equations with a polysubstituted halogenoalkane	2
(iii)	orange/yellow/brown to colourless/decolourises	NOT red/red-brown	1
(b) (i)	any two from: M1 same functional group M2 the same/similar chemical properties OR undergo same/similar chemical reactions M3 trend in physical properties M4 differ by CH_2	NOT similar reactivity ALLOW a named physical property e.g. boiling point NOT similar physical properties	2
(ii)	M1 same molecular formula M2 different displayed/structural formulae	NOT same empirical/general formula ALLOW different structures/arrangements	2
(iii)		ALLOW E/trans isomer 	1

(c)	M1 chain length longer in poly(ethene) M2 polymer contains only single (covalent) bonds /no double bond M3 ethene is a gas and poly(ethene) is a solid	ALLOW monomer contains double C=C bonds ALLOW reactant/ethene unsaturated ALLOW product/polyethene saturated	3
		Total = 12	

Question number	Answer	Notes	Marks
11 (a) (i)	$2\text{PbS} + 3\text{O}_2 \rightarrow 2\text{PbO} + 2\text{SO}_2$ M1 formulae of O_2 and SO_2 M2 rest of equation correctly balanced	M2 dep on M1 ALLOW multiples/fractions	2
(ii)	(sulfur dioxide causes) acid rain / breathing problems	ALLOW named breathing problems such as asthma ALLOW other effects of acid rain such as killing fish, damage to stonework, killing plants	1
(iii)	M1 (moles lead(II) oxide) = $892\,000\,000 \div 223$ OR $4\,000\,000$ moles M2 (moles of carbon dioxide) = $2\,000\,000$ M3(mass of carbon dioxide) = 88 (tonnes)	ALLOW calculations done in megamoles throughout ALLOW $\text{M1} \div 2$ 88 (tonnes) scores 3 marks	3
(iv)	any 5 from: lead(II) sulfide M1 giant ionic structure/lattice M2 strong (ionic) bonds OR strong electrostatic forces (between oppositely charged) ions M3 which take a lot of energy to break / overcome sulfur dioxide M4 simple molecular/covalent structure M5 weak intermolecular forces OR weak forces between molecules M6 which take little energy to overcome	REJECT molecules/covalent bonds/ intermolecular forces for all three marks M3 dep on M2 REJECT ions/ionic bonds for all 3 marks ALLOW molecules NOT particles/atoms NOT weak IMF between atoms M6 dep on M5	5
(b)	M1 $90.7 \div 207$ and $9.30 \div 16$ M2 0.438 (moles of lead) and 0.581 (moles of	NOT atomic numbers ALLOW $9.30 \div 32$ for ecf	4

	oxygen) M3 ratio of moles = 1:1.33 M4 empirical formula is Pb_3O_4	Answer must be 2sf or more ALLOW 1.3 ALLOW ecf from ratio shown to produce formula	
		Total = 15	

