



Pearson  
Edexcel

# Mark Scheme (Results)

January 2019

Pearson Edexcel International GCSE

In Human Biology (4HB0) Paper 2

## **Edexcel and BTEC Qualifications**

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at [www.edexcel.com](http://www.edexcel.com) or [www.btec.co.uk](http://www.btec.co.uk). Alternatively, you can get in touch with us using the details on our contact us page at [www.edexcel.com/contactus](http://www.edexcel.com/contactus).

### **Pearson: helping people progress, everywhere**

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: [www.pearson.com/uk](http://www.pearson.com/uk)

January 2019

Publications Code 4HB0\_02\_1901\_MS

All the material in this publication is copyright

© Pearson Education Ltd 2019

## General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

Question number	Answer	Notes	Marks
1		<p>All correct = 6 marks  5 or 6 correct = 5 marks</p>	<p>6</p> <p>Total 6 marks</p>

Question number	Answer	Notes	Marks
2 (a) (i)	<ul style="list-style-type: none"> <li>• otherwise would see food/ identify food by sight/would not need to taste or smell food/to ensure food is identified by taste/smell only;</li> </ul>		1
(ii)	<ul style="list-style-type: none"> <li>• 160;</li> <li>• 200;</li> <li>• 80%;</li> </ul>	Full marks for correct final answer	3
(iii)	<p>THREE OF</p> <ul style="list-style-type: none"> <li>• more people correctly identified food when nostrils not closed/when only eyes covered/ easier to identify food when nostrils not covered;</li> <li>• smell increases ability to identify food;</li> <li>• taste increases ability to identify food;</li> <li>• variations/differences between people;</li> </ul>	Accept description of differences	max 3
(iv)	<ul style="list-style-type: none"> <li>• reference to guesswork;</li> <li>• anomalous result for Person 3</li> </ul>		
(b)	<p>TWO OF</p> <ul style="list-style-type: none"> <li>• size/volume/mass/type of food;</li> <li>• time left on tongue/same area of tongue;</li> <li>• same people/age/gender;</li> </ul>		3
(c)	<p>TWO OF</p> <ul style="list-style-type: none"> <li>• reduce/eliminate effect of anomalous result/to identify anomalies;</li> <li>• to compare results;</li> <li>• calculate mean;</li> <li>• more reliable;</li> </ul>		max 2
(c)	touch/texture/sight;	ACCEPT description that implies sense or valid alternative	1
			Total 13 marks

Question number	Answer	Notes	Marks
3 (a) (i)	<ul style="list-style-type: none"> <li>• <u>deamination</u>;</li> <li>• (<u>excess</u>) amino acids;</li> <li>• in liver;</li> </ul>	Allow 1 mark for breakdown of amino acids for marking points 1 AND 2	3
(ii)	<ul style="list-style-type: none"> <li>• not very toxic/medium toxicity;</li> <li>• highly soluble;</li> <li>• little water needed to remove it/enables water to be conserved;</li> <li>• not much energy/<u>only</u> 4ATP required to remove each molecule/little energy used;</li> </ul>		4
(iii)	<ul style="list-style-type: none"> <li>• urine is a solution/fluid/urea is a solid;</li> <li>• urine is a mixture/urea not a mixture;</li> <li>• (urine) contains dissolved salts/urea;</li> <li>• urea is an excretory substance/nitrogen containing molecule;</li> </ul>		3
(b) (i)	plasma;		1
(ii)	<ul style="list-style-type: none"> <li>• kidneys keep urea concentration constant/at 100 au/remove urea from the blood;</li> <li>• concentration keeps on rising when kidneys not working;</li> </ul>		2
(c)	<ul style="list-style-type: none"> <li>• dialysis;</li> <li>• transplant;</li> </ul>		2
			Total 15 marks

Question number	Answer	Notes	Marks
4	<p>8 of</p> <ul style="list-style-type: none"> <li>• diffusion</li> <li>• movement from high to low concentration/down a concentration gradient;</li> <li>• osmosis;</li> <li>• movement of water molecules;</li> <li>• from high to low water potential/from a dilute to solution to a (more) concentrated solution;</li> <li>• through selectively permeable membrane;</li> </ul> <p>(Factors affecting diffusion/osmosis):</p> <ul style="list-style-type: none"> <li>• surface area;</li> <li>• temperature;</li> <li>• distance to travel;</li> <li>• ref to steepness of gradient;</li> <li>• active transport/uptake;</li> <li>• uses energy/ATP;</li> <li>• against a concentration gradient;</li> </ul>	<p>Ignore references to concentrations of water ALLOW once with ref to diffusion/osmosis</p> <p>MP's 7-10 allow once with ref to either diffusion or osmosis</p>	<p>8</p> <p>Total 8 marks</p>

Question number	Answer	Notes	Marks										
5 (a) (i)	<table border="1" data-bbox="384 342 1015 533"> <thead> <tr> <th data-bbox="384 342 700 383">Function</th> <th data-bbox="700 342 1015 383">Letter of chamber</th> </tr> </thead> <tbody> <tr> <td data-bbox="384 383 700 423"><i>receives oxygenated blood</i></td> <td data-bbox="700 383 1015 423">Y;</td> </tr> <tr> <td data-bbox="384 423 700 463"><i>sends blood to lungs</i></td> <td data-bbox="700 423 1015 463">W;</td> </tr> <tr> <td data-bbox="384 463 700 504"><i>receives deoxygenated blood</i></td> <td data-bbox="700 463 1015 504">X;</td> </tr> <tr> <td data-bbox="384 504 700 533"><i>sends blood to body</i></td> <td data-bbox="700 504 1015 533">Z;</td> </tr> </tbody> </table>	Function	Letter of chamber	<i>receives oxygenated blood</i>	Y;	<i>sends blood to lungs</i>	W;	<i>receives deoxygenated blood</i>	X;	<i>sends blood to body</i>	Z;	Accept from Z to Y	4
Function	Letter of chamber												
<i>receives oxygenated blood</i>	Y;												
<i>sends blood to lungs</i>	W;												
<i>receives deoxygenated blood</i>	X;												
<i>sends blood to body</i>	Z;												
(ii)	<ul style="list-style-type: none"> <li>• prevents backflow of blood;</li> <li>• from left ventricle to left atrium;</li> </ul>	2											
(b)	<p>FOUR OF</p> <ul style="list-style-type: none"> <li>• blood can pass between ventricles/oxygenated and deoxygenated blood mix;</li> <li>• blood doesn't travel to lungs/body effectively;</li> <li>• poor oxygenation/less oxygen in blood/to body cells;</li> <li>• less aerobic respiration/less energy/less growth;</li> <li>• reference to anaerobic respiration/build-up of lactic acid;</li> <li>• reference to drop in blood pressure/increase in heart rate;</li> </ul>		<p>Max 4</p> <p>Total 10 marks</p>										

Question number	Answer	Notes	Marks
6 (a)	temperature;		1
(b)	<ul style="list-style-type: none"> <li>• maximum rate of activity between 60-70 ° /optimum temperature for this enzyme is between 60-70 °C ;</li> <li>• optimum temperature for human enzymes 37-40°C;</li> <li>• this enzyme would be less active/inactive at body temperature/37-40 °C;</li> <li>• human enzymes denature/breakdown above optimum/over 40 °C;</li> </ul>		4
(c)	<ul style="list-style-type: none"> <li>• enzyme denatures;</li> <li>• shape of active site lost/disrupted;</li> <li>• no enzyme-substrate complexes formed/no binding of substrate possible;</li> </ul>	Reject dies	3  Total 8 marks

Pearson Education Limited. Registered company number 872828  
with its registered office at 80 Strand, London, WC2R 0RL, United Kingdom

