



Pearson

Mark Scheme (Results)

Summer 2018

Pearson Edexcel International GCSE
In Mathematics B (4MB1)
Paper 02

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

Summer 2018

Publications Code 4MB1_01_1806_MS

All the material in this publication is copyright

© Pearson Education Ltd 2018

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.
- **Types of mark**
 - M marks: method marks
 - A marks: accuracy marks
 - B marks: unconditional accuracy marks (independent of M marks)
- **Abbreviations**
 - cao – correct answer only
 - ft – follow through
 - isw – ignore subsequent working
 - SC - special case
 - oe – or equivalent (and appropriate)
 - dep – dependent
 - indep – independent
 - awrt – answer which rounds to
 - eeoo – each error or omission

- **No working**

If no working is shown then correct answers normally score full marks

If no working is shown then incorrect (even though nearly correct) answers score no marks.

- **With working**

If there is a wrong answer indicated on the answer line always check the working in the body of the script (and on any diagrams), and award any marks appropriate from the mark scheme.

If it is clear from the working that the "correct" answer has been obtained from incorrect working, award 0 marks.

Any case of suspected misread loses A (and B) marks on that part, but can gain the M marks.

If working is crossed out and still legible, then it should be given any appropriate marks, as long as it has not been replaced by alternative work.

If there is a choice of methods shown, then no marks should be awarded, unless the answer on the answer line makes clear the method that has been used.

If there is no answer on the answer line then check the working for an obvious answer.

- **Ignoring subsequent work**

It is appropriate to ignore subsequent work when the additional work does not change the answer in a way that is inappropriate for the question: eg. Incorrect cancelling of a fraction that would otherwise be correct.

It is not appropriate to ignore subsequent work when the additional work essentially makes the answer incorrect eg algebra.

Transcription errors occur when candidates present a correct answer in working, and write it incorrectly on the answer line; mark the correct answer.

- **Parts of questions**

Unless allowed by the mark scheme, the marks allocated to one part of the question CANNOT be awarded in another part.

| Question | Working | Answer | Notes | Mark | Sub-Total | Total |
|----------|--|-----------|---|------|-----------|-------|
| 1(a) | $\frac{600}{4} \times (12 + 9 + 4)$ oe | | Full method to find total number of seats | M1 | | |
| | | 3750 | | A1 | 2 | |
| 1(b) | Stalls: 1800 Dress Circle: 1350 | | | B1 | | |
| | $(1800 \times 65) + (0.9 \times (1350 \times 40)) + (0.25 \times (600 \times 25))$ | | | M1 | | |
| | | \$169,350 | | A1 | 3 | |
| 1(c) | $\frac{175000 - 169350}{25}$ (= 226) or $\frac{175000 - 117000 - 48600}{25}$ | | | M1 | | |
| | | 376 | | A1 | 2 | 7 |

| Question | Working | Answer | Notes | Mark | Sub-Total | Total |
|----------|--|-----------------------------|--|------|-----------|-------|
| 2 | $AC^2 = 5^2 + 12^2$ or $AC = 13$ or $6^2 + 2.5^2 (=42.25)$ $(E \text{ to middle of } BC)^2 = 11^2 - 2.5^2$ or $E\text{-mid}BC = \frac{3\sqrt{51}}{2}$ $(E \text{ to middle of } AB)^2 = 11^2 - 6^2$ or $E\text{-mid}AB = \sqrt{85}$ | | Correct method to find side of triangle in first stage to find height of pyramid | M1 | | |
| | $OE^2 = 11^2 - ((0.5 \times "13")^2)$ or $11^2 - 2.5^2 - 6^2$ oe | | Correct statement for $(OE)^2$ oe | M1 | | |
| | | $OE = \frac{3\sqrt{35}}{2}$ | Allow 8.87(41...) | A1 | | |
| | $V = \frac{1}{3}(12)(5)(OE)$ | | (dep on M2) or for an answer of 177.(4823...) | M1 | | |
| | | $30\sqrt{35}$ | Or $p = 30, q = 35$ | A1 | | 5 |

| Question | Working | Answer | Notes | Mark | Sub-Total | Total |
|----------|--|--------------------------|--|----------------|-----------|-------|
| 3 | $\mathbf{A}^2 = \begin{pmatrix} a^2 + 12 & 3a - 3 \\ 4a - 4 & 13 \end{pmatrix}$ | | | B2 (-1 eeo) | | |
| | $\begin{pmatrix} a^2 + 12 & 3a - 3 \\ 4a - 4 & 13 \end{pmatrix} - \begin{pmatrix} 13 & 3 \\ 4 & 10 \end{pmatrix} = \lambda \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ | | Their $\mathbf{A}^2 - \mathbf{B} = \lambda \mathbf{I}$ | M1 | | |
| | | $a = 2$ $\lambda = 3$ | | A1 A1 | | 5 |

| Question | Working | Answer | Notes | Mark | Sub-Total | Total |
|----------|---------|------------------------|---|------|-----------|-------|
| 4(a) | | $\{h, l\}$ | | B1 | 1 | |
| 4(b) | | $\{a, b, c, f, g, h\}$ | | B1 | 1 | |
| 4(c) | | 4 | | B1 | 1 | |
| 4(d) | | 10 | | B1 | 1 | |
| 4(e) | | \subset | | B1 | 1 | |
| 4(f) | | $\frac{2}{7}$ | Decimals or % to 2dp truncated or rounded 0.28(5714...) | B1 | 1 | 6 |

| Question | Working | | Answer | Notes | Mark | Sub-Total | Total |
|----------|--|---|--|---|------|-----------|-------|
| 5(a) | $3 + 5a = 0$ | | | | M1 | | |
| | | | $a = -0.6$ | oe | A1 | 2 | |
| 5(b) | | | $f(x) < 3$ | Accept any equivalent notation but not $x < 3$ | B1 | 1 | |
| 5(c) | | | $x = \frac{2}{3}$ | | B1 | 1 | |
| 5(d) | $(3 + 5x)(3x - 2) = 5 - 4x$ | | | Removing denominator in a correct equation | M1oe | | |
| | $15x^2 + 3x - 11 = 0$ | | | | A1 | | |
| | $\frac{-3 \pm \sqrt{3^2 - 4(15)(-11)}}{2(15)} = \frac{-3 \pm \sqrt{69}}{30}$ | | | Solving their 3 term quadratic | M1oe | | |
| | 0.762... and/or -0.962... | | | Or | A1 | | |
| | awrt -0.962 | | -0.962 | Rejection of positive root | A1 | 5 | |
| 5(e) | $y = \frac{5 - 4x}{3x - 2}$ $y(3x - 2) = 5 - 4x$ | $x = \frac{5 - 4y}{3y - 2}$ $x(3y - 2) = 5 - 4y$ | | Write in x or y in terms of y or x and multiply | M1 | | |
| | $3xy - 2y = 5 - 4x$ $3xy + 4x = 5 + 2y$ | $3xy - 2x = 5 - 4y$ $3xy + 4y = 5 + 2x$ | | expand and isolate terms in x or y | M1 | | |
| | $x(3y + 4) = 5 + 2y$ $x = \frac{5 + 2y}{3y + 4}$ | $y(3x + 4) = 5 + 2x$ $y = \frac{5 + 2x}{3x + 4}$ | | Factorise out x or y and divide. | M1 | | |
| | | | $g^{-1} : x \rightarrow \frac{5 + 2x}{3x + 4}$ | Correct answer written in fully correct form. | A1 | 4 | 13 |

| Question | Working | Answer | Notes | Mark | Sub-Total | Total |
|----------|---|----------------|---|---------|-----------|-------|
| 6(a) | $x + y = 12$ | | | B1indep | | |
| | $\frac{x + y + y + 12 + (x + y) + 12}{6} = 9.5$ | | | M1indep | | |
| | eg $2x + 3y = 33$ | | Terms in x and y and numerical terms simplified | A1 | | |
| | $2x + 3(12 - x) = 33$ | | Solve simultaneous equations (dependent on previous M mark) | M1 dep | | |
| | | $x = 3, y = 9$ | | A1 | 5 | |
| 6(b) | Median = $\frac{"9"+12}{2}$ | | Correct calculation with their value of y | M1ft | | |
| | | 10.5 | cao | A1 | 2 | 7 |

| Question | Working | Answer | Notes | Mark | Sub-Total | Total |
|----------|---|---|--|----------------|-----------|-------|
| 7(a) | | Triangle <i>A</i> drawn | Penalise labelling ONCE only | B1 | 1 | |
| 7(b) | | $x = -2$ drawn | Implied by correct triangle or two correct coordinates | B1 | | |
| | | Triangle <i>B</i> drawn | $(-5,1), (-5,3), (-4,3)$ | B1 | 2 | |
| 7(c) | $(-4,-2), (-4,0), (-3,0)$ | Triangle <i>C</i> drawn | ft from their <i>B</i> | B2 (-1 ee) | 2 | |
| 7(d) | $\begin{pmatrix} -2 & 0 \\ 0 & -2 \end{pmatrix} \begin{pmatrix} -4 & -4 & -3 \\ -2 & 0 & 0 \end{pmatrix}$ | | NB: coordinates may be in any order | M1ft | | |
| | | $\begin{pmatrix} 8 & 8 & 6 \\ 4 & 0 & 0 \end{pmatrix}$ | | A1ft | | |
| | | Triangle <i>D</i> drawn | cao | A1 | 3 | |
| 7(e) | $\begin{pmatrix} -0.5 & 0 \\ 0 & -0.5 \end{pmatrix}$ | $\frac{1}{4} \begin{pmatrix} -2 & 0 \\ 0 & -2 \end{pmatrix}$ oe | B1 for one correct row or column & correct expression for det. | B2 | 2 | |
| 7(f) | | enlargement scale factor -0.5 oe centre $(0, 0)$ oe | | B1 B1 B1 | 3 | |

| Question | Working | Answer | Notes | Mark | Sub-Total | Total |
|----------|--|---------------------------|--|------|-----------|-------|
| 8(a) | $y = \frac{1}{2}x^4 - \frac{1}{2}x^2 - \frac{5}{3}x^3 + 11x + \frac{1}{3}$ | | Expanding(at least 1 term correct) and differentiating (at least one term correct) | M1 | | |
| | $\frac{dy}{dx} = 2x^3 - x - 5x^2 + 11$ and $\frac{dy}{dx} = 5$ when $x = p$ giving $2p^3 - p - 5p^2 + 11 = 5$ | | Sets their derivative = 5 with at least two terms differentiated correctly | M1 | | |
| | | $2p^3 - 5p^2 - p + 6 = 0$ | Dep on M2 NB: answer given | A1 | 3 | |
| 8(b) | $2(1.5)^3 - 5(1.5)^2 - (1.5) + 6$ | | Substitute 1.5 into cubic or by algebraic division | M1 | | |
| | $2(1.5)^3 - 5(1.5)^2 - (1.5) + 6 = 0$ | | Showing that the remainder is zero | A1 | 2 | |
| 8(c) | $p^2 \dots\dots$ | | Showing division with at least p^2 in quotient | M1 | | |
| | | $p^2 - p - 2$ | Fully correct quotient | A1 | | |
| | $(p+1)(p-2)$ | | $p^2 - p - 2$ correctly factorised | M1 | | |
| | | $(2p-3)(p+1)(p-2)$ | Fully factorised expression | A1 | 4 | |
| 8(d) | $q = \frac{1}{2}(2)^2(2^2 - 1) - \frac{5}{3}(2)^3 + 11(2) + \frac{1}{3}$ | | substitution of 2 into y to give q | M1 | | |
| | | 15 | | A1 | | |

| Question | Working | Answer | Notes | Mark | Sub-Total | Total |
|----------|---|---|---------------------------------------|------|-----------|-------|
| 9(a)(i) | | $\mathbf{b - a}$ | Must be lower case a, b | B1 | | |
| 9(a)(ii) | | $-2\mathbf{a}$ | Must be lower case a | B1 | 2 | |
| 9(b)(i) | $\overline{CD} = '-2\mathbf{a}' + \lambda\mathbf{b} + \mu\mathbf{a}$ or $\overline{AE} = 3'(\mathbf{b} - \mathbf{a})' + \lambda\mathbf{b}$ | | | M1ft | | |
| | $\overline{CD} = '-2\mathbf{a}' + \lambda\mathbf{b} + \mu\mathbf{a}$ and $\overline{AE} = 3'(\mathbf{b} - \mathbf{a})' + \lambda\mathbf{b}$ | Expressions for \overline{CD} and \overline{AE} | | A1ft | | |
| 9(b)(ii) | | $(\mu - 2)\mathbf{a} + \lambda\mathbf{b}$ and $-3\mathbf{a} + (3 + \lambda)\mathbf{b}$ | Both expressions simplified | A1ft | 3 | |
| 9(c) | $4(\mu - 2) = -3$ | | Forming equation for a | M1ft | | |
| | | $\mu = \frac{5}{4}$ oe | | A1 | | |
| | $4\lambda = 3 + \lambda$ | | Forming equation for b | M1ft | | |
| | | $\lambda = 1$ | | A1 | 4 | |
| 9(d) | $\frac{1}{2}(4\mu + -2\mathbf{a})h = 5$ oe eg $\frac{1}{2}(5 + 8)h = 5$ | | Using their \overline{CF} and μ | M1 | | |
| | | $h = \frac{10}{13}$ | | A1 | | |
| | $\sin \angle CFE = \frac{h}{\lambda} = \frac{10}{13}$ | | | M1 | | |
| | | 50.3° | | A1 | 4 | 13 |

| Question | Working | Answer | Notes | Mark | Sub-Total | Total |
|----------|--|---|--|-------------------|-----------|-------|
| 10(a) | | 6.64, -3.33, 1.6 | Penalise rounding once | B1,B1,B1 | 3 | |
| 10(b) | <p>Curve -1 mark for straight line segments</p> <p>Each point missed</p> <p>Each missed segment</p> <p>Each point incorrectly plotted</p> <p>Tramlines</p> <p>Very poor curve</p> <p>N.B. Accuracy for both plotting and drawing is $\pm \frac{1}{2}$ss</p> | A correct curve | | B3 ft (-1 eeo) | 3 | |
| 10(c) | $x^2 - 5x + \frac{8}{x} = ax + b$ or $x^2 + \frac{8}{x} < \frac{21}{4}x + 3$ oe | | | M1 | | |
| | $4x^3 + 4(-5-a)x^2 - 4bx + 32 = 0$ oe Or $x^2 - 5x + \frac{8}{x} < \frac{1}{4}x + 3$ | | Re-arrange and attempt to solve for a and b | M1 | | |
| | $a = \frac{1}{4}$ and $b = 3$ | <p>Draw line</p> $y = \frac{1}{4}x + 3$ | | A1 | | |
| | 1.1 and 5.5 | | | A1 | | |
| | | $1.1 < x < 5.5$ | Dep on previous marks ft their graph. Must be correct inequality signs | A1 | 5 | 11 |
| | | | | | | |
| | | | | | | |

| Question | Working | Answer | Notes | Mark | Sub-Total | Total |
|----------|---|--|--|------------------------|-----------|-------|
| 11(a) | | $\frac{5}{14}, \frac{9}{14}$ $\frac{3}{7}, \frac{4}{7}$ $\frac{4}{13}, \frac{9}{13}$ | Ignore extra branches completed oe | B1 B1 B1 | 3 | |
| 11(b) | $\frac{2}{5} \times \frac{9}{14}$, or $\frac{3}{5} \times \frac{3}{7}$, or $\frac{3}{5} \times \frac{4}{7}$, or $\frac{3}{5} \times 1$ or $\frac{6}{15} \times \frac{5}{14}$ | | ft their tree diagram | M1 | | |
| | $\frac{2}{5} \times \frac{9}{14} + \frac{3}{5} \times \frac{3}{7} + \frac{3}{5} \times \frac{4}{7}$, or $1 - \frac{6}{15} \times \frac{5}{14}$ | | ft their tree diagram | M1dep | | |
| | | $\frac{6}{7}$ oe | 0.85(714...) decimals and % to 2dp truncated or rounded | A1 | 3 | |
| 11(c) | $P(A): \left(\frac{2}{5} \times \frac{9}{14}\right) + \left(\frac{2}{5} \times \frac{5}{14} \times \frac{4}{13}\right) + \left(\frac{3}{5} \times \frac{3}{7}\right)$ oe $= \frac{254}{455}$ | | Complete method to calculate either 15 (A) or 20 (B) cents removed 0.55(824...) | M1ft | | |
| | $P(B): \left(\frac{2}{5} \times \frac{5}{14} \times \frac{9}{13}\right) + \left(\frac{3}{5} \times \frac{4}{7}\right)$ oe = $\frac{201}{455}$ | | Correct probability for either 15 or 20 ft 0.44(175...) | A1ft | | |
| | | A is more likely as $\frac{254}{455} > \frac{201}{455}$ | Correct conclusion with correct reason dep on both A and B correct | A1 cao | 3 | |
| | | | | | | |

