



Pearson
Edexcel

Mark Scheme (Results)

November 2020

Pearson Edexcel International GCSE
In Mathematics B (4MB1)
Paper 01R

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

November 2020

Publications Code 4MB1_01R_2011_MS

All the material in this publication is copyright

© Pearson Education Ltd 2020

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme.

Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.

- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

- **Types of mark**

- M marks: method marks
- A marks: accuracy marks
- B marks: unconditional accuracy marks (independent of M marks)

- **Abbreviations**

- cao – correct answer only
- ft – follow through
- isw – ignore subsequent working
- SC – special case
- oe – or equivalent (and appropriate)
- dep – dependent

- indep – independent
- awrt – answer which rounds to
- eeo – each error or omission

- **No working**

If no working is shown then correct answers normally score full marks
 If no working is shown then incorrect (even though nearly correct) answers score no marks.

- **With working**

If there is a wrong answer indicated on the answer line always check the working in the body of the script (and on any diagrams), and award any marks appropriate from the mark scheme.

If it is clear from the working that the “correct” answer has been obtained from incorrect working, award 0 marks.

If a candidate misreads a number from the question. Eg. Uses 252 instead of 255; method marks may be awarded provided the question has not been simplified. Examiners should send any instance of a suspected misread to review. If there is a choice of methods shown, then no marks should be awarded, unless the answer on the answer line makes clear the method that has been used.

If there is no answer on the answer line then check the working for an obvious answer.

- **Ignoring subsequent work**

It is appropriate to ignore subsequent work when the additional work does not change the answer in a way that is inappropriate for the question: eg. Incorrect cancelling of a fraction that would otherwise be correct.

It is not appropriate to ignore subsequent work when the additional work essentially makes the answer incorrect eg algebra.

Transcription errors occur when candidates present a correct answer in working, and write it incorrectly on the answer line; mark the correct answer.

- **Parts of questions**

Unless allowed by the mark scheme, the marks allocated to one part of the question CANNOT be awarded to another.

Question	Working	Answer	Mark	Notes
1	$\frac{2563}{11} \times 13$ and $\frac{2563}{11} \times 8$ or $\frac{2563}{11} \times 5$		M1	
		1165	A1	
Total 2 marks				
2	$\frac{8(2x+3)^2}{4(2x+3)}$ or $\frac{2(2x+3)^2}{(2x+3)}$ or $\frac{8(2x+3)}{4}$		M1	Factorising the denominator or dividing by 4 or cancelling a bracket. Implied by a correct answer
		4x + 6	A1	or equivalent, for example 2(2x + 3)
Total 2 marks				
3	$\frac{16}{3}$ and $\frac{13}{5}$ or $\frac{16}{3}$ and $\frac{5}{13}$ oe		M1	
	$\frac{16}{3} \times \frac{5}{13} = \frac{80}{39}$ or e.g. $\frac{80}{15} \div \frac{39}{15} = \frac{80}{39}$ Allow $\frac{16}{3} \div \frac{13}{5} = \frac{80}{39}$		M1	Must multiply both correctly. Or for $\frac{16}{3} \times \dots = \dots$, $\dots \times \frac{5}{13} = \dots$ Note that $\frac{16}{3} \times \frac{5}{13} = 2\frac{2}{39}$ is M1M1A0
		$2\frac{2}{39}$	A1	Must be given as a mixed number
Total 3 marks				
4	$\frac{36}{60}$ or 3.6 or $3\frac{36}{60}$ or $3 \times 60 + 36 [=216]$		M1	
	$\frac{11.52}{"3.6"}$ or $\frac{11.52}{"216"} \times 60$		M1	For using distance \div time. Allow 11520 instead of 11.52, e.g. $\frac{11520}{"3.6"} \div 1000$
		3.2	A1	
Total 3 marks				

5		$45500 \times 1.055 [= 48002.5]$		M1	
		"48002.5" $\times 1.0125^2$		M1	M2 only if after 49210 working out a third year e.g. 49825.1...
			49 210	A1	Awrt (For reference: 49210.06289...)
Total 3 marks					
6	(a)	$y = \frac{3x+5}{2}$		M1	Attempt to re-arrange to make y the subject – allow sign errors only
			$\frac{3}{2}$	A1	Or equivalent (e.g. 1.5) – M1A0 for $\frac{3}{2}x$ Condone for full marks those that have the correct gradient but incorrect constant term $\left(\text{e.g. } y = \frac{3x-5}{2} \Rightarrow m = \frac{3}{2} \right)$
	(b)		$\left(0, \frac{5}{2} \right)$	B1	
Total 3 marks					
7		$\frac{1}{10} \times \frac{1}{a} \times \frac{2}{b}$ where $a = 9$ or 10 and $b = 8$ or 10		M1	Condone if other terms seen (e.g. from considering other possible orders of the letters P, A and L) but not if divided by 6
		$\frac{1}{10} \times \frac{1}{9} \times \frac{2}{8}$		A1	Only (so no other terms seen)
			$\frac{1}{360}$	A1	Accept awrt 0.00278
Total 3 marks					

8		$3x+4 \quad 2x-3 \quad (>0)$		M1	For solving given quadratic. When factorising must give at least 2 correct terms when multiplied out
		$x < "-\frac{4}{3}", x > "\frac{3}{2}"$		M1	Selecting outside regions for their critical values. Condone non-strict inequalities or 'incorrect' notation (e.g. $-\frac{4}{3} > x > \frac{3}{2}$)
			$x < -\frac{4}{3}, x > \frac{3}{2}$	A1	Must be 2 separate inequalities. Do not accept $-\frac{4}{3} > x > \frac{3}{2}$ but condone 'and' separating the 2 inequalities. Mark final answer.
					Total 3 marks
9		$y = \frac{3x}{2x+2}$ and an attempt to clear the fraction $2y \quad x+2 = 3x$		M1	
		$x \quad 2y-3 = -4y$		M1	Expand, rearranging and factorise out x (allow sign errors only)
			$\frac{-4x}{2x-3}$ or $\frac{4x}{3-2x}$	A1	
					Total 3 marks
10		$\frac{6-\sqrt{8}}{2+\sqrt{8}} \times \frac{2-\sqrt{8}}{2-\sqrt{8}}$ or $\frac{3-\sqrt{2}}{1+\sqrt{2}} \times \frac{1-\sqrt{2}}{1-\sqrt{2}}$		M1	Or equivalent, for example, $\frac{6-\sqrt{8}}{2+\sqrt{8}} \times \frac{-2+\sqrt{8}}{-2+\sqrt{8}}$
		$\frac{20-8\sqrt{8}}{-4}$ or $-5+2\sqrt{8}$ or $-5+4\sqrt{2}$		M1dep	For multiplying out correctly and simplifying
			$-5+\sqrt{32}$	A1	If $-5+4\sqrt{2}$ with no working, then no marks If $-5+\sqrt{32}$ seen, then SC B1 only
					Total 3 marks

11		$-3(x-...)^2 - ...$		M1	Or $a = -3$
		$-3(x-1)^2 - ...$		M1	Or $b = -1$
			$a = -3$ $b = -1$ $c = 5$	A1	Allow $-3(x-1)^2 + 5$. For those that consider $3x^2 - 6x - 2$ and conclude $3(x-1)^2 - 5$ then M0M1A0 but allow full marks if then followed by correct answer
Total 3 marks					
12		$\frac{n}{440} = \frac{144}{360}$		M1	Or equivalent
			176, 55	A1	
		$\frac{209}{440} \times 360 [= 171]$ or $\frac{55}{440} \times 360 [= 45]$ oe		M1	Implied by 45 or 171
			Milk 171 or White 45	A1	Drawn accurately $\pm 1^\circ$ at least 1 labelled correctly (angle and type)
Total 4 marks					
13		$b = \frac{b}{2} + a$ or $\frac{b}{2} = a$ oe		M1	
		$\frac{4a + 5.5b + 8}{5} = 10.6$ oe		M1	Forming an equation in a and b using the mean e.g. $a + b + \frac{1}{2}b + a + 2b + a + 3 + 2b + a + 5 = 10.6$
		$7.5b = 45$ or $15a = 45$		M1dep	Forming an equation in a or b only – dependent on both previous M marks
			$a = 3, b = 6$	A1	Correct answers from two correct equations with no working to solve these two equations scores 3 marks. SC B1 both correct answers with no working
Total 4 marks					

14		$BD = 2\tan 64$ [4.10...]		M1	Accept any equivalent form (so BD does not need to be the subject) e.g. $BD = \frac{2}{\tan 26}$ or $\frac{BD}{\sin 64} = \frac{2}{\sin 26}$
		$\sin 28 = \frac{BD}{BC}$		M1	This mark can be implied by a correct expression for BC Or equivalent e.g. $\frac{BC}{\sin 90} = \frac{BD}{\sin 28}$
		$BC = \frac{"2 \tan 64"}{"\sin 28"}$		M1dep	Dependent on both previous M marks
			$BC = 8.7$	A1	Accept AWRT to 8.7 (for reference: 8.7345177...)
Total 4 marks					
15	(a)	$\begin{pmatrix} 9 & -3 \\ -12 & 6 \end{pmatrix} - \begin{pmatrix} 4 & -1 \\ 4 & -4 \end{pmatrix}$		M1	Correct implied subtraction or three correct numbers in the final answer in correct places.
			$\begin{pmatrix} 5 & -2 \\ -16 & 10 \end{pmatrix}$	A1	
	(b)	$\mathbf{Q} = \begin{pmatrix} 0 & 1 \\ -10 & -17 \end{pmatrix} \begin{pmatrix} 5 & -3 \\ -3 & 2 \end{pmatrix}$		M1	Or general \mathbf{Q} and \mathbf{P} inverse to form simultaneous equations, e.g. $\mathbf{Q} = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \Rightarrow \begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} 2 & 3 \\ 3 & 5 \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ -10 & -17 \end{pmatrix}$ $\Rightarrow 2a+3b=0, 3a+5b=1, 2c+3d=-10, 3c+5d=-17$ (must have correct inverse for this alternative method)
			$\begin{pmatrix} -3 & 2 \\ 1 & -4 \end{pmatrix}$	A1	
Total 4 marks					

16		$PK.PJ = PM.PL$ or $5.5 \times 2.5 = 2 \times PL$		M1	
		$PL = 6.875$ or $\frac{5.5 \times 2.5}{2}$		A1	
		$\text{Area} = \frac{1}{2} \times 6.875 \times 2.5 \times \sin 74$		M1	Correct use of $0.5absinC$ for their PL M0 if angle LKP assumed to be right-angled
			8.26	A1	Accept AWRT to 8.26 (For reference: 8.260842...)
Total 4 marks					
17	(a)		2.34×10^{-2}	B1	
	(b)		240 000	B1	
	©	0.5×10^{199} or 5×10^n $n = 198$ or 200		M1	
			5×10^{198}	A1	
Total 4 marks					
18	(a)	$12w^9y^0$		M1	Two of the three parts correct
			$12w^9$	A1	
	(b)	125 or x^6		M1	
			$125x^6$	A1	
Total 4 marks					
19		$\frac{FE}{3} = \frac{FC}{AC}$, $\frac{FE}{7} = \frac{AF}{AC}$		M1 M1	
		$FC = \frac{7}{3} AF$			
		$AC = AF + \frac{7}{3} AF \left[= \frac{10}{3} AF \right]$		M1	Using $AC = AF + FC$
		$FE = 3 \times \frac{\frac{7}{3} AF}{\frac{10}{3} AF}$		M1	
			2.1	A1	Or equivalent
Total 5 marks					

21		$180 - \frac{360}{11} [= 147.27\dots]$ or $\frac{9 \times 180}{11}$		M1	
		$5 \times 180 [= 900]$		M1	Could be seen in working
		"900" - $5 \times$ "147.2..." [= 163.6... or $\frac{1800}{11}$]		M1dep	Oe e.g. $2x + 5 \times 147.2 = 900$ - this mark is dependent on both previous M marks
		$x = \frac{163.6}{2}$		M1dep	Dependent on all previous M marks
			81.8	A1	Accept exact $\left(\frac{900}{11}\right)$ or awrt to 81.8 (81.81818...) Oe e.g. $\frac{180 - \left(\frac{360}{11}\right) \times \frac{1}{2}}{2}$
					Total 5 marks
22	(a)	$\frac{(360 - 76)}{360} \times \pi \times 16$ or $16\pi - \frac{76}{360} \times \pi \times 16$		M1	
			39.7	A1	Awrt to 39.7 or 39.6 (For reference: $39.65388\dots$ or $\frac{568}{45}\pi$)
	(b)	$\frac{360 - 76}{360} \times 2 \times \pi \times 4 [= 19.8\dots]$		M1	Or equivalent, e.g. $2\pi(4) - \frac{76}{360} \times 2\pi(4)$ (For reference if correct: $19.82694\dots$ or $\frac{284}{45}\pi$)
		$19.8\dots + 4 + 4$		M1dep	
			27.8	A1	Awrt to 27.8 (For reference: $27.826940\dots$)
					Total 5 marks

23		$y = k_1 x^3$		M1	
		$x = \frac{k_2}{\sqrt{w}}$		M1	Condone if the same letter is used for the two constants of proportionality
		$729 = k_1 \times 4.5^3$ or $25 = \frac{k_2}{\sqrt{0.16}}$		M1dep	Dependent on one of the previous M marks – both equations imply the previous two marks
		$y = 8x^3$ and $x = \frac{10}{\sqrt{w}}$		A1	Or equivalent
			$y = 8000w^{-\frac{3}{2}}$	A1	Or equivalent e.g. $y = 8 \left(\frac{10}{w^{\frac{1}{2}}} \right)^3$
					Total 5 marks
24		$\frac{10x-10}{2x+1} = \frac{35x-5}{10x-10}$ or $\frac{10x-10}{2x+1} = \frac{7x-1}{2x-2}$		M1	Forming a correct equation any form
		$100x^2 - 200x + 100 = 70x^2 + 25x - 5$ or $20x^2 - 40x + 20 = 14x^2 + 5x - 1$		M1dep	Removing the fractions and one side multiplied out correctly
		$30x^2 - 225x + 105 = 0$ or $6x^2 - 45x + 21 = 0$ or $2x^2 - 15x + 7 = 0$		A1	A correct three term quadratic in x (any equivalent form) = 0
		$= (2x-1)(x-7)$		M1	Correct method for solving their three-term quadratic in x , implied by $x = \frac{1}{2}$ or 7 Dependent on both previous M marks
			4	A1	Only (A0 if given as the answer -2.5 too)
					Total 5 marks

25		$\vec{AC} = \begin{pmatrix} 3+x \\ -2+5x \end{pmatrix}$		B1	
		$(3+x)^2 + (-2+5x)^2 = 25$		M1	Correct method for finding the modulus of \vec{AC} (must contain x) and = 5 to form an equation – not from $\vec{AC} = \begin{pmatrix} 3x \\ -7x \end{pmatrix}$
		$13x^2 - 7x - 6 = 0$		M1dep	For correct method for multiplying out the brackets and simplifying to make a quadratic equation = 0
		$(x-1)(13x+6) = 0$ $\Rightarrow \vec{AC} = \begin{pmatrix} 3+'1' \\ -2+5('1') \end{pmatrix}$		M1	Correct method for solving 3-term quadratic and substituting their positive value of x into \vec{AC}
			$\vec{AC} = \begin{pmatrix} 4 \\ 3 \end{pmatrix}$	A1	Answer with no working gains no marks
					Total 5 marks

26	(a)		$v = k - 10t$	B1	Allow mark if seen in part (b) – give bod if just right-hand side stated
	(b)	" $k - 10t = 0$ " leading to $t = \frac{k}{10}$ or $k = 10t$		M1	Calculating the time from projection to maximum height (or equivalent) from their (a)
		Maximum height reached by the ball is $(h-1) + h = 161 \Rightarrow h = 81$		B1	Correctly deriving the maximum height reached by the ball (possibly implied in later working)
		$f(k) = 1 + \frac{k^2}{10} - \frac{5k^2}{100}$ or $f(t) = 1 + 10t^2 - 5t^2$		M1	Eliminating either k or t using their $k = 10t$ to obtain an expression in a single variable only (bod if the +1 is absent)
		$81 = 1 + \frac{k^2}{10} - \frac{5k^2}{100}$ or $81 = 1 + 10t^2 - 5t^2$		M1dep	Setting up a correct equation in terms of k or t only with their $k = 10t$ (dependent on both previous M marks and B mark)
			40	A1	A0 if final answer is ± 40 Note that $1 + k\left(\frac{k}{10}\right) - 5\left(\frac{k}{10}\right)^2$ equal to either 161, 160, 80.5 or 80 (or equivalent) will score M1B0M1M0A0 or possibly M1B1M1M0A0 (if max. height found but subsequently not used)
					Total 6 marks